

ESSENTIALS

HOFFBRAND'S ESSENTIAL HAEMATOLOGY

A. VICTOR HOFFBRAND | PAUL A.H. MOSS

SEVENTH EDITION



WILEY Blackwell

Hoffbrand's Essential Haematology

This title is also available as an e-book.

For more details, please see

www.wiley.com/buy/9781118408674

or scan this QR code:



Hoffbrand's Essential Haematology

A. Victor Hoffbrand

MA DM FRCP FRCPATH FRCP(Edin) DSc FMedSci

Emeritus Professor of Haematology

University College London

London, UK

Paul A. H. Moss

PhD MRCP FRCPATH

Professor of Haematology

University of Birmingham

Birmingham, UK

Seventh Edition

WILEY Blackwell

This edition first published 2016 © 2016 by John Wiley & Sons Ltd
Previous editions: 1980, 1984, 1993, 2001, 2006, 2011

Registered office: John Wiley & Sons, Ltd, The Atrium, Southern Gate, Chichester, West Sussex, PO19 8SQ, UK

Editorial offices: 9600 Garsington Road, Oxford, OX4 2DQ, UK
The Atrium, Southern Gate, Chichester, West Sussex, PO19 8SQ, UK
111 River Street, Hoboken, NJ 07030-5774, USA

For details of our global editorial offices, for customer services and for information about how to apply for permission to reuse the copyright material in this book please see our website at www.wiley.com/wiley-blackwell

The right of the author to be identified as the author of this work has been asserted in accordance with the UK Copyright, Designs and Patents Act 1988.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, except as permitted by the UK Copyright, Designs and Patents Act 1988, without the prior permission of the publisher.

Designations used by companies to distinguish their products are often claimed as trademarks. All brand names and product names used in this book are trade names, service marks, trademarks or registered trademarks of their respective owners. The publisher is not associated with any product or vendor mentioned in this book. It is sold on the understanding that the publisher is not engaged in rendering professional services. If professional advice or other expert assistance is required, the services of a competent professional should be sought.

The contents of this work are intended to further general scientific research, understanding, and discussion only and are not intended and should not be relied upon as recommending or promoting a specific method, diagnosis, or treatment by health science practitioners for any particular patient. The publisher and the author make no representations or warranties with respect to the accuracy or completeness of the contents of this work and specifically disclaim all warranties, including without limitation any implied warranties of fitness for a particular purpose. In view of ongoing research, equipment modifications, changes in governmental regulations, and the constant flow of information relating to the use of medicines, equipment, and devices, the reader is urged to review and evaluate the information provided in the package insert or instructions for each medicine, equipment, or device for, among other things, any changes in the instructions or indication of usage and for added warnings and precautions. Readers should consult with a specialist where appropriate. The fact that an organization or Website is referred to in this work as a citation and/or a potential source of further information does not mean that the author or the publisher endorses the information the organization or Website may provide or recommendations it may make. Further, readers should be aware that Internet Websites listed in this work may have changed or disappeared between when this work was written and when it is read. No warranty may be created or extended by any promotional statements for this work. Neither the publisher nor the author shall be liable for any damages arising herefrom.

Library of Congress Cataloging-in-Publication Data

Hoffbrand, A. V., author.

[Essential haematology]

Hoffbrand's essential haematology / A. Victor Hoffbrand, Paul A. H. Moss. — Seventh edition.

p. ; cm. — (Essentials)

Includes index.

ISBN 978-1-118-40867-4 (pbk.)

I. Moss, P. A. H., author. II. Title. III. Series: Essentials (Wiley-Blackwell (Firm)).

[DNLM: 1. Hematologic Diseases. WH 120]

RC633

616.1'5—dc23


2015019097

A catalogue record for this book is available from the British Library.

Wiley also publishes its books in a variety of electronic formats. Some content that appears in print may not be available in electronic books.

Cover image: P242/0207. Blood cells, SEM. NATIONAL CANCER INSTITUTE/SCIENCE PHOTO LIBRARY. Blood cells and platelets. Coloured Scanning Elec-tron micrograph (SEM) of human blood showing red and white cells and platelets. Red blood cells (erythrocytes) have a characteristic biconcave- disc shape and are numerous. These large cells contain haemoglobin, a red pigment by which oxygen is transported around the body. They are more numerous than white blood cells (yellow). White blood cells (leucocytes) are rounded cells with microvilli projections from the cell surface. Leucocytes play an important role in the immune response of the body. Platelets are smaller cells (pink) that play a major role in blood clotting.

Set in 10/12 Adobe Garamond Pro by Aptara Inc., New Delhi, India



Contents

Preface to the Seventh Edition	vi	16 Myelodysplasia	177
Preface to the First Edition	vii	17 Acute lymphoblastic leukaemia	186
How to use your textbook	viii	18 The chronic lymphoid leukaemias	197
About the companion website	x	19 Hodgkin lymphoma	205
1 Haemopoiesis	1	20 Non-Hodgkin lymphoma	213
2 Erythropoiesis and general aspects of anaemia	11	21 Multiple myeloma and related disorders	228
3 Hypochromic anaemias	27	22 Aplastic anaemia and bone marrow failure	242
4 Iron overload	41	23 Stem cell transplantation	250
5 Megaloblastic anaemias and other macrocytic anaemias	48	24 Platelets, blood coagulation and haemostasis	264
6 Haemolytic anaemias	60	25 Bleeding disorders caused by vascular and platelet abnormalities	278
7 Genetic disorders of haemoglobin	72	26 Coagulation disorders	290
8 The white cells 1: granulocytes, monocytes and their benign disorders	87	27 Thrombosis 1: pathogenesis and diagnosis	302
9 The white cells 2: lymphocytes and their benign disorders	102	28 Thrombosis 2: treatment	311
10 The spleen	116	29 Haematological changes in systemic disease	321
11 The aetiology and genetics of haematological malignancies	122	30 Blood transfusion	333
12 Management of haematological malignancy	135	31 Pregnancy and neonatal haematology	346
13 Acute myeloid leukaemia	145	Appendix: World Health Organization classification of tumours of the haematopoietic and lymphoid tissues	352
14 Chronic myeloid leukaemia	156	Index	357
15 Myeloproliferative disease	165		

Preface to the Seventh Edition

There have been remarkable advances in the understanding of the pathogenesis of diseases of the blood and lymphatic system and in the treatment of these diseases, since the 6th Edition of *Essential Haematology* was published in 2011. This new knowledge is due largely to the application of next generation sequencing of DNA which has enabled the detection of the genetic mutations, inherited or acquired, that underlie these diseases. As examples, sequencing has revealed the *CALR* mutation underlying a substantial proportion of patients with myeloproliferative diseases and the *MYD88* mutation present in almost all cases of Waldenström's macroglobulinaemia. Multiple 'driver' gene mutations affecting signalling pathways and epigenetic reactions involved in cell proliferation and survival have been discovered which underlie myelodysplasia, acute myeloid and lymphoblastic leukaemias, chronic lymphocytic leukaemia and the lymphomas. The complexity of the molecular changes underlying the malignant diseases and the relevance of this to their sensitivity or resistance to therapy is becoming apparent.

This new knowledge has been accompanied by spectacular improvements in therapy. Inhibition of the B cell receptor signalling pathway has transformed the life expectancy in many patients with resistant chronic lymphocytic leukaemia and some of the B cell lymphomas resistant to other therapy. *JAK2* inhibitors are improving the quality of life and survival in primary myelofibrosis. Survival in myeloma is improving remarkably with new proteasome inhibitory and immunomodulatory drugs. Life expectancy has also improved for patients with diseases such as thalassaemia major receiving multiple transfusions with the worldwide introduction of orally active iron chelating agents. New anticoagulants which directly inhibit at a single point in the coagulation cascade and rarely need monitoring are now used commonly in preference to warfarin for the treatment and prevention of arterial and venous thrombosis.

These advances in knowledge have been incorporated as new text, diagrams and tables for this seventh edition. New multiple choice questions have been added to the website and short summary boxes are included at the end of each chapter.

We thank Dr Trevor Baglin for his helpful suggestions for the coagulation section of the book. We wish to thank our publishers Wiley-Blackwell and the staff who have helped us with the production of this 7th Edition. We also thank Jane Fallows for once more producing clear, expertly drawn scientific diagrams. We hope it will be widely used both by undergraduates and by postgraduates in medicine and related sciences wishing to gain a grounding in one of the most exciting and advanced fields of medicine.

Victor Hoffbrand
Paul Moss

Preface to the First Edition

The major changes that have occurred in all fields of medicine over the last decade have been accompanied by an increased understanding of the biochemical, physiological and immunological processes involved in normal blood cell formation and function and the disturbances that may occur in different diseases. At the same time, the range of treatment available for patients with diseases of the blood and blood-forming organs has widened and improved substantially as understanding of the disease processes has increased and new drugs and means of support care have been introduced.


We hope the present book will enable the medical student of the 1980s to grasp the essential features of modern clinical and laboratory haematology and to achieve an understanding of how many of the manifestations of blood diseases can be explained with this new knowledge of the disease processes.

We would like to thank many colleagues and assistants who have helped with the preparation of the book. In particular, Dr H.G. Prentice cared for the patients whose haematological responses are illustrated in Figs 5.3 and 7.8 and Dr J. McLaughlin supplied Fig. 8.6. Dr S. Knowles reviewed critically the final manuscript and made many helpful suggestions. Any remaining errors are, however, our own. We also thank Mr J.B. Irwin and R.W. McPhee who drew many excellent diagrams, Mr Cedric Gilson for expert photomicrography, Mrs T. Charalambos, Mrs B. Elliot, Mrs M. Evans and Miss J. Allaway for typing the manuscript, and Mr Tony Russell of Blackwell Scientific Publications for his invaluable help and patience.

**AVH, JEP
1980**

How to use your textbook

Features contained within your textbook



◀ Every chapter begins with a list of **Key topics** of the chapter.

▶ Every chapter ends with a **Summary** that can be used for study and revision purposes.

10 / Chapter 1: Haemopoiesis

proteins that are involved in mediating apoptosis following DNA damage, such as p53 and ATM, are also frequently mutated and therefore inactivated in haemopoietic malignancies.

Necrosis is death of cells and adjacent cells due to ischaemia, chemical trauma or hyperthermia. The cells swell, the plasma

membrane loses integrity. There is usually an inflammatory infiltrate in response to spillage of cell contents. Autophagy is the digestion of cell organelles by lysosomes. It may be involved in cell death but in some situations also in maintaining cell survival by recycling nutrients.

SUMMARY

- Haemopoiesis (blood cell formation) arises from pluripotent stem cells in the bone marrow. Stem cells give rise to progenitor cells which, after cell divisions and differentiation, form red cells, granulocytes (neutrophils, eosinophils and basophils), monocytes, platelets and B and T lymphocytes.
- Haemopoietic tissue occupies about 50% of the marrow space in normal adult marrow. Haemopoiesis in adults is confined to the central skeleton but in infants and young children haemopoietic tissue extends down the long bones of the arms and legs.
- Stem cells reside in the bone marrow in niches formed by stromal cells and circulate in the blood.
- Growth factors attach to specific cell receptors and produce a cascade of phosphorylation events to the cell nucleus. Transcription factors carry the message to those genes that are to be 'switched on', to stimulate cell division, differentiation, functional activity or suppress apoptosis.
- Adhesion molecules are a large family of glycoproteins that mediate attachment of marrow precursors and mature leucocytes and platelets to extracellular matrix, endothelium and to each other.
- Epigenetics refers to changes in DNA and chromatin that affect gene expression other than those that affect DNA sequence. Histone modification and DNA methylation are two important examples relevant to haemopoiesis and haematological malignancies.
- Transcription factors are molecules that bind to DNA and control the transcription of specific genes or gene families.
- Apoptosis is a physiological process of cell death resulting from activation of caspases. The intracellular ratio of pro-apoptotic proteins (e.g. BAX) to anti-apoptotic proteins (e.g. BCL-2) determines the cell susceptibility to apoptosis.




Now visit www.wileyessential.com/haematology to test yourself on this chapter.


► Your textbook is full of **photographs, illustrations and tables.**

CD8 molecules, whereas the CD8 molecule recognizes the HLA-A, B and C molecules (see Fig. 2.15). The antigen recognition site of the TCR is joined to several other subunits in the CD3 complex which together mediate signal transduction. Depending on their cytokine production, CD4⁺ T cells can be further subdivided into T helper type 1 (Th1) and Th2 cells. Th1 cells produce mainly IL-2, TNF- β and interferon (IFN- γ), and are important in boosting cell-mediated immunity (and granuloma formation), whereas Th2 cells produce IL-4 and IL-6 and are mainly responsible for providing help for antibody production.

Antigen-specific immune responses are generated in so-called **lymphoid organs** and centres, where antigen is presented via a lymph node (Fig. 9.10) or dendritic cells. B cells recognize antigen through their surface immunoglobulin and although most antibody responses require help from antigen-specific T cells, some antigens such as polysaccharides can lead to T cell-independent B cell antibody production. In the full body, germinal centres arise as a result of continuing exposure to antigenic stimulation (Fig. 9.11). These consist of follicular dendritic cells (FDCs), which are loaded with antigen, B cells and activated T cells which have migrated up from the T zone.



(a)



(b)

Figure 9.10 (a) Structure of a lymph node. (b) Lymph node showing germinal follicles surrounded by a darker mantle zone rim and lighter, more diffuse marginal and T zone areas.




Figure 9.11 Generation of a germinal centre. B cells activated by antigen migrate from the T zone to the follicle where they undergo massive proliferation. Cells enter the dark zone as centroblasts and accumulate mutations in their immunoglobulin V genes. Cells then pass back into the light zone (Fig. 9.10) as centrocytes. Only those cells that can interact with antigen on follicular dendritic cells and receive signals from antigen-specific T cells (Fig. 9.8) are selected and migrate out as plasma cells and memory cells. Cells not selected die by apoptosis.

Proliferating B cells move to the dark zone of the germinal centre as **centroblasts** where they undergo somatic mutation of their immunoglobulin variable-region genes (Fig. 9.11). Their progeny are known as **centrocytes** and these must be selected for survival by antigen on FDCs, otherwise they undergo apoptosis. If selected they become memory B cells or plasma cells (Fig. 9.11). Plasma cells originate in the bone marrow and either stay in the RE system and produce high-affinity antibody.

Lymphocytosis

Lymphocytosis often occurs in infants and young children in response to infections that produce a neutrophil reaction in adults. Conditions particularly associated with lymphocytosis are listed in Table 9.3.

Chlaudius fever is a general term for a disease characterized by fever, sore throat, lymphadenopathy and capillary lymphocytes in the blood. It may be caused by primary infection with Epstein-Barr virus (EBV), cytomegalovirus, human immunodeficiency virus (HIV) or *Toxoplasma*. EBV infection, otherwise known as infectious mononucleosis, is the most common cause.

Table 9.3 Causes of lymphocytosis.

Infections
Acute infectious mononucleosis, rubella, pertussis, mumps, acute infectious lymphocytosis, infectious hepatitis, cytomegalovirus, HIV, herpes simplex or zoster, chronic tuberculosis, toxoplasmosis, brucellosis, syphilis
Chronic lymphoid leukaemias (see Chapter 18)
Acute lymphoblastic leukaemia (Chapter 17)
Non-Hodgkin lymphoma (see Chapter 23)
Thymomas
Anti-tumour immunosuppressive virus

Infectious mononucleosis

This is caused by primary infection with EBV and occurs only in a minority of infected individuals – in most cases infection is subclinical. The disease is characterized by a lymphocytosis caused by clonal expansion of T cells reacting against B lymphocytes infected with EBV. The disease is associated with a

Differential diagnosis of macrocytic anaemias

The clinical history and physical examination may suggest B_{12} or folate deficiency as the cause. Diet, drugs, alcohol intake, family history, history suggestive of malabsorption, presence of autoimmune diseases or other associations with pernicious anaemia (Table 5.4), previous gastrointestinal disease or operations are all important. The presence of jaundice, glossitis or neuropathy are also important indications of megaloblastic anaemia.


The laboratory features of particular importance are the shape of macrocytes (oval in megaloblastic anaemia), the presence of hypersegmented neutrophils, of leucopenia and thrombocytopenia in megaloblastic anaemia, and the bone marrow appearance. Assay of serum B_{12} and folate is essential. Exclusion of alcoholism (particularly if the patient is not anaemic), liver and thyroid function tests, and bone marrow examination for myelodysplasia, aplasia or myeloma are important in the investigation of macrocytosis not caused by B_{12} or folate deficiency.

- Macrocytic anaemias show an increased size of circulating red cells (MCV >98 fL). Causes include vitamin B_{12} (B_{12} , cobalamin) or folate deficiency, alcohol, liver disease, hypothyroidism, myelodysplasia, paraproteinaemia, cytotoxic drugs, aplastic anaemia, pregnancy and the neonatal period.
- B_{12} or folate deficiency cause megaloblastic anaemia, in which the bone marrow erythroblasts have a typical abnormal appearance.
- Folates take part in biochemical reactions in DNA synthesis. B_{12} has an indirect role by its involvement in folate metabolism.
- B_{12} deficiency may also cause a neuropathy due to damage to the spinal cord and peripheral nerves.
- B_{12} deficiency is usually caused by B_{12} malabsorption brought about by pernicious anaemia in which there

is autoimmune gastritis, resulting in severe deficiency of intrinsic factor, a glycoprotein made in the stomach which facilitates B_{12} absorption by the ileum.

- Other gastrointestinal diseases as well as a vegan diet may cause B_{12} deficiency.
- Folate deficiency may be caused by a poor diet, malabsorption (e.g. gluten-induced enteropathy) or excess cell turnover (e.g. pregnancy, haemolytic anaemias, malignancy).
- Treatment of B_{12} deficiency is usually with injections of hydroxocobalamin and of folate deficiency with oral folic (pteroylglutamic) acid.
- Rare causes of megaloblastic anaemia include inborn errors of B_{12} or folate transport or metabolism, and defects of DNA synthesis not related to B_{12} or folate.

SUMMARY



The website icon

indicates that you can find accompanying multiple choice questions and answers on the book's companion website.



Now visit www.wileyessential.com/haematology to test yourself on this chapter.

About the companion website

Don't forget to visit the companion website for this book:

www.wileyessential.com/haematology




There you will find valuable material designed to enhance your learning, including:

- Interactive multiple choice questions
- Figures and tables from the book

Scan this QR code to visit the companion website






CHAPTER 1

Haemopoiesis

Key topics

■ Site of haemopoiesis	2
■ Haemopoietic stem and progenitor cells	2
■ Bone marrow stroma	4
■ The regulation of haemopoiesis	4
■ Haemopoietic growth factors	4
■ Growth factor receptors and signal transduction	6
■ Adhesion molecules	8
■ The cell cycle	8
■ Transcription factors	8
■ Epigenetics	8
■ Apoptosis	9



This first chapter is concerned with the general aspects of blood cell formation (haemopoiesis). The processes that regulate haemopoiesis and the early stages of formation of red cells (erythropoiesis), granulocytes and monocytes (myelopoiesis) and platelets (thrombopoiesis) are also discussed.

Site of haemopoiesis

In the first few weeks of gestation the yolk sac is a transient site of haemopoiesis. However, definitive haemopoiesis derives from a population of stem cells first observed on the AGM (aorta-gonads-mesonephros) region. These common precursors of endothelial and haemopoietic cells (haemangioblasts) are believed to seed the liver, spleen and bone marrow. From 6 weeks until 6–7 months of fetal life, the liver and spleen are the major haemopoietic organs and continue to produce blood cells until about 2 weeks after birth (Table 1.1; see Fig. 7.1b). The placenta also contributes to fetal haemopoiesis. The bone marrow is the most important site from 6–7 months of fetal life. During normal childhood and adult life the marrow is the only source of new blood cells. The developing cells are situated outside the bone marrow sinuses; mature cells are released into the sinus spaces, the marrow microcirculation and so into the general circulation.

In infancy all the bone marrow is haemopoietic but during childhood there is progressive fatty replacement of marrow throughout the long bones so that in adult life haemopoietic marrow is confined to the central skeleton and proximal ends of the femurs and humeri (Table 1.1). Even in these haemopoietic areas, approximately 50% of the marrow consists of fat (Fig. 1.1). The remaining fatty marrow is capable of reversion to haemopoiesis and in many diseases there is also expansion of haemopoiesis down the long bones. Moreover, the liver and spleen can resume their fetal haemopoietic role ('extramedullary haemopoiesis').

Haemopoietic stem and progenitor cells

Haemopoiesis starts with a pluripotential stem cell that can by asymmetric cell division self-renew but also give rise to the separate cell lineages. These cells are able to repopulate a bone marrow from which all stem cells have been eliminated by lethal irradiation or chemotherapy. This **haemopoietic stem cell** (HSC) is rare, perhaps 1 in every 20 million nucleated cells




Figure 1.1 Normal bone marrow trephine biopsy (posterior iliac crest). Haematoxylin and eosin stain; approximately 50% of the intertrabecular tissue is haemopoietic tissue and 50% is fat.

in bone marrow. Many of the cells are dormant and in mice it has been estimated that they enter cell cycle approximately every 20 weeks. Although its exact phenotype is unknown, on immunological testing the HSC is CD34⁺ CD38⁻ and negative for lineage markers (Lin⁻) and has the appearance of a small or medium-sized lymphocyte (see Fig. 23.3). The cells reside in specialized osteoblastic or vascular 'niches'.

Cell differentiation occurs from the stem cell via committed **haemopoietic progenitors** which are restricted in their developmental potential (Fig. 1.2). The existence of the separate progenitor cells can be demonstrated by *in vitro* culture techniques. Very early progenitors are assayed by culture on bone marrow stroma as long-term culture initiating cells, whereas late progenitors are generally assayed in semi-solid media. An example is the earliest detectable mixed myeloid precursor which gives rise to granulocytes, erythrocytes, monocytes and megakaryocytes and is termed CFU (colony-forming unit)-GEMM (Fig. 1.2). The bone marrow is also the primary site of origin of lymphocytes, which differentiate from a common lymphoid precursor. The spleen, lymph nodes and thymus are secondary sites of lymphocyte production (see Chapter 9).

The stem cell has the capability for **self-renewal** (Fig. 1.3) so that marrow cellularity remains constant in a normal healthy steady state. There is considerable amplification in the system: one stem cell is capable of producing about 10⁶ mature blood cells after 20 cell divisions (Fig. 1.3). In humans HSCs are capable of about 50 cell divisions, telomere shortening affecting viability. Under normal conditions most are dormant. With aging, the number of stem cells falls and the relative proportion giving rise to lymphoid rather than myeloid progenitors also falls. Stem cells also accumulate genetic mutations with age, an average of 8 at age 60, and these, either passenger or driver, may be present in tumours arising from these stem cells (see Chapter 11). The precursor cells are capable of responding to haemopoietic growth factors with increased production of one

Table 1.1 Sites of haemopoiesis.

Fetus	0–2 months (yolk sac)
	2–7 months (liver, spleen)
	5–9 months (bone marrow)
Infants	Bone marrow (practically all bones)
Adults	Vertebrae, ribs, sternum, skull, sacrum and pelvis, proximal ends of femur




Figure 1.2 Diagrammatic representation of the bone marrow pluripotent stem cell and the cell lines that arise from it. Various progenitor cells can be identified by culture in semi-solid medium by the type of colony they form. It is possible that an erythroid/megakaryocytic progenitor may be formed before the common lymphoid progenitor diverges from the mixed granulocytic/monocyte/eosinophil myeloid progenitor. Baso, basophil; BFU, burst-forming unit; CFU, colony-forming unit; E, erythroid; Eo, eosinophil; GEMM, granulocyte, erythroid, monocyte and megakaryocyte; GM, granulocyte, monocyte; Meg, megakaryocyte; NK, natural killer.




Figure 1.3 (a) Bone marrow cells are increasingly differentiated and lose the capacity for self-renewal as they mature. **(b)** A single stem cell gives rise, after multiple cell divisions (shown by vertical lines), to $>10^6$ mature cells.

or other cell line when the need arises. The development of the **mature cells** (red cells, granulocytes, monocytes, megakaryocytes and lymphocytes) is considered further in other sections of this book.

Bone marrow stroma

The bone marrow forms a suitable environment for stem cell survival, self-renewal and formation of differentiated progenitor cells. It is composed of stromal cells and a microvascular network (Fig. 1.4). The stromal cells include mesenchymal stem cells, adipocytes, fibroblasts, osteoblasts, endothelial cells and macrophages and they secrete extracellular molecules such as collagen, glycoproteins (fibronectin and thrombospondin) and glycosaminoglycans (hyaluronic acid and chondroitin derivatives) to form an extracellular matrix. In addition, stromal cells secrete several growth factors necessary for stem cell survival.

Mesenchymal stem cells are critical in stromal cell formation. Together with osteoblasts or endothelial cells they form niches and provide the growth factors, adhesion molecules and cytokines which support stem cells, e.g. the protein jagged, on stromal cells, binds to a receptor NOTCH1 on stem cells which then becomes a transcription factor involved in the cell cycle.

Stem cells are able to traffic around the body and are found in peripheral blood in low numbers. In order to exit the bone marrow, cells must cross the blood vessel endothelium and this process of **mobilization** is enhanced by administration of

growth factors such as granulocyte colony-stimulating factor (G-CSF) (see p. 91). The reverse process of stem cell **homing** appears to depend on a chemokine gradient in which the stromal-derived factor 1 (SDF-1) which binds to its receptor CXCR4 on HSC is critical. Several critical interactions maintain stem cell viability and production in the stroma including stem cell factor (SCF) and jagged proteins expressed on stroma and their respective receptors KIT and NOTCH expressed on stem cells.

The regulation of haemopoiesis

Haemopoiesis starts with stem cell division in which one cell replaces the stem cell (*self-renewal*) and the other is committed to differentiation. These early committed progenitors express low levels of transcription factors that may commit them to discrete cell lineages. Which cell lineage is selected for differentiation may depend both on chance and on the external signals received by progenitor cells. Several transcription factors (see p. 8) regulate survival of stem cells (e.g. SCL, GATA-2, NOTCH-1), whereas others are involved in differentiation along the major cell lineages. For instance, PU.1 and the CEBP family commit cells to the myeloid lineage, whereas GATA-2 and then GATA-1 and FOG-1 have essential roles in erythropoietic and megakaryocytic differentiation. These transcription factors interact so that reinforcement of one transcription programme may suppress that of another lineage. The transcription factors induce synthesis of proteins specific to a cell lineage. For example, the erythroid-specific genes for globin and haem synthesis have binding motifs for GATA-1.

Haemopoietic growth factors

The haemopoietic growth factors are glycoprotein hormones that regulate the proliferation and differentiation of haemopoietic progenitor cells and the function of mature blood cells. They may act locally at the site where they are produced by cell–cell contact or circulate in plasma. They also bind to the extracellular matrix to form niches to which stem and progenitor cells adhere. The growth factors may cause cell proliferation but can also stimulate differentiation, maturation, prevent apoptosis and affect the function of mature cells (Fig. 1.5).

They share a number of common properties (Table 1.2) and act at different stages of haemopoiesis (Table 1.3; Fig. 1.6). Stromal cells are the major source of growth factors except for erythropoietin, 90% of which is synthesized in the kidney, and thrombopoietin, made largely in the liver. An important feature of growth factor action is that two or more factors may synergize in stimulating a particular cell to proliferate or differentiate. Moreover, the action of one growth factor on a cell may stimulate production of another growth factor or growth factor receptor. SCF and FLT3 ligand (FLT3-L) act locally on the pluripotent stem cells and on early myeloid and lymphoid progenitors (Fig. 1.6). Interleukin-3 (IL-3) and




Figure 1.4 Haemopoiesis occurs in a suitable microenvironment ('niche') provided by a stromal matrix on which stem cells grow and divide. The niche may be vascular (lined by endothelium) or endosteal (lined by osteoblasts). There are specific recognition and adhesion sites (see p. 8); extracellular glycoproteins and other compounds are involved in the binding.




Figure 1.5 Growth factors may stimulate proliferation of early bone marrow cells, direct differentiation to one or other cell type, stimulate cell maturation, suppress apoptosis or affect the function of mature non-dividing cells, as illustrated here for granulocyte colony-stimulating factor (G-CSF) for an early myeloid progenitor and a neutrophil.

Table 1.2 General characteristics of myeloid and lymphoid growth factors.
Glycoproteins that act at very low concentrations
Act hierarchically
Usually produced by many cell types
Usually affect more than one lineage
Usually active on stem/progenitor cells and on differentiated cells
Usually show synergistic or additive interactions with other growth factors
Often act on the neoplastic equivalent of a normal cell
Multiple actions: proliferation, differentiation, maturation, functional activation, prevention of apoptosis of progenitor cells

granulocyte–macrophage colony-stimulating factor (GM-CSF) are multipotential growth factors with overlapping activities. G-CSF and thrombopoietin enhance the effects of SCF, FLT-L, IL-3 and GM-CSF on survival and differentiation of the early haemopoietic cells.

These factors maintain a pool of haemopoietic stem and progenitor cells on which later-acting factors, erythropoietin, G-CSF, macrophage colony-stimulating factor (M-CSF), IL-5 and thrombopoietin, act to increase production of one or other cell lineage in response to the body’s need. Granulocyte and monocyte formation, for example, can be stimulated by infection or inflammation through release of IL-1 and tumour necrosis factor (TNF) which then stimulate stromal cells to produce growth factors in an interacting network (see Fig. 8.4). In contrast, cytokines, such as transforming growth factor- β (TGF- β) and γ -interferon (IFN- γ), can exert a negative effect on haemopoiesis and may have a role in the development of aplastic anaemia (see p. 244).

Table 1.3 Haemopoietic growth factors.

Act on stromal cells

IL-1
TNF

Act on pluripotent stem cells

SCF
FLT3-L
VEGF

Act on multipotent progenitor cells

IL-3
GM-CSF
IL-6
G-CSF
Thrombopoietin

Act on committed progenitor cells

G-CSF*
M-CSF
IL-5 (eosinophil-CSF)
Erythropoietin
Thrombopoietin*

CSF, colony-stimulating factor; FLT3-L, FLT3 ligand; G-CSF, granulocyte colony-stimulating factor; GM-CSF, granulocyte-macrophage colony-stimulating factor; IL, interleukin; M-CSF, macrophage colony-stimulating factor; SCF, stem cell factor; TNF, tumour necrosis factor; VEGF, vascular endothelial growth factor.

* These also act synergistically with early acting factors on pluripotent progenitors.

Growth factor receptors and signal transduction

The biological effects of growth factors are mediated through specific receptors on target cells. Many receptors (e.g. erythropoietin (epo) receptor (R), GMCSF-R) are from the *haematopoietin receptor superfamily* which dimerize after binding their ligand.

Dimerization of the receptor leads to activation of a complex series of intracellular signal transduction pathways, of which the three major ones are the JAK/STAT, the mitogen-activated protein (MAP) kinase and the phosphatidylinositol 3 (PI3) kinase pathways (Fig. 1.7; see Fig. 15.2). The Janus-associated kinase (JAK) proteins are a family of four tyrosine-specific protein kinases that associate with the intracellular domains of the growth factor receptors (Fig. 1.7). A growth factor molecule binds simultaneously to the extracellular domains of two or three receptor molecules, resulting in their aggregation. Receptor aggregation induces activation of the JAKs which now phosphorylate members of the signal transducer and activator of transcription (STAT) family of transcription factors. This results in their dimerization and translocation from the cell cytoplasm across the nuclear membrane to the cell nucleus. Within the nucleus STAT dimers activate transcription of specific genes. A model for control of gene expression by a transcription factor is shown in Fig. 1.8. The clinical importance of




Figure 1.6 A diagram of the role of growth factors in normal haemopoiesis. Multiple growth factors act on the earlier marrow stem and progenitor cells. EPO, erythropoietin; PSC, pluripotent stem cell; SCF, stem cell factor; TPO, thrombopoietin; FLT3-L, FLT3 ligand. For other abbreviations see Fig. 1.2.




Figure 1.7 Control of haemopoiesis by growth factors. The factors act on cells expressing the corresponding receptors. Binding of a growth factor to its receptor activates the JAK/STAT, MAPK and phosphatidyl-inositol 3-kinase (PI3K) pathways (see Fig. 15.2) which leads to transcriptional activation of specific genes. E2F is a transcription factor needed for cell transition from G1 to S phase. E2F is inhibited by the tumour suppressor gene Rb (retinoblastoma) which can be indirectly activated by p53. The synthesis and degradation of different cyclins stimulates the cell to pass through the different phases of the cell cycle. The growth factors may also suppress apoptosis by activating AKT (protein kinase B).




Figure 1.8 Model for control of gene expression by a transcription factor. The DNA-binding domain of a transcription factor binds a specific enhancer sequence adjacent to a structural gene. The transactivation domain then binds a molecule of RNA polymerase, thus augmenting its binding to the TATA box. The RNA polymerase now initiates transcription of the structural gene to form mRNA. Translation of the mRNA by the ribosomes generates the protein encoded by the gene.

this pathway is revealed by the finding of an activating mutation of the *JAK2* gene as the cause of polycythaemia rubra vera (see p. 166).

JAK can also activate the MAPK pathway, which is regulated by RAS and controls proliferation. PI3 kinases phosphorylate inositol lipids which have a wide range of downstream effects including activation of AKT leading to block of apoptosis and other actions (Fig. 1.7; see Fig. 15.2). Different domains of the intracellular receptor protein may signal for the different processes (e.g. proliferation or suppression of apoptosis) mediated by growth factors.

A second smaller group of growth factors, including SCF, FLT-3L and M-CSF (Table 1.3), bind to receptors that have an extracellular immunoglobulin-like domain linked via a transmembrane bridge to a cytoplasmic tyrosine kinase domain. Growth factor binding results in dimerization of these receptors and consequent activation of the tyrosine kinase domain. Phosphorylation of tyrosine residues in the receptor itself generates binding sites for signalling proteins which initiate complex cascades of biochemical events resulting in changes in gene expression, cell proliferation and prevention of apoptosis.

Adhesion molecules

A large family of glycoprotein molecules termed adhesion molecules mediate the attachment of marrow precursors, leucocytes and platelets to various components of the extracellular matrix, to endothelium, to other surfaces and to each other. The adhesion molecules on the surface of leucocytes are termed receptors and these interact with proteins termed ligands on the surface of target cells, e.g. endothelium. The adhesion molecules are important in the development and maintenance of inflammatory and immune responses, and in platelet–vessel wall and leucocyte–vessel wall interactions.

The pattern of expression of adhesion molecules on tumour cells may determine their mode of spread and tissue localization (e.g. the pattern of metastasis of carcinoma cells or non-Hodgkin lymphoma cells into a follicular or diffuse pattern). The adhesion molecules may also determine whether or not cells circulate in the bloodstream or remain fixed in tissues. They may also partly determine whether or not tumour cells are susceptible to the body's immune defences.

The cell cycle

The cell division cycle, generally known simply as the **cell cycle**, is a complex process that lies at the heart of haemopoiesis. Dysregulation of cell proliferation is also the key to the development of malignant disease. The duration of the cell cycle is variable between different tissues but the basic principles remain constant. The cycle is divided into the mitotic phase (**M phase**), during which the cell physically divides, and **interphase**, during which the chromosomes are duplicated and cell growth occurs prior to division (Fig. 1.7). The M phase

is further partitioned into classical **mitosis**, in which nuclear division is accomplished, and **cytokinesis**, in which cell fission occurs.

Interphase is divided into three main stages: a **G₁ phase**, in which the cell begins to commit to replication, an **S phase**, during which DNA content doubles and the chromosomes replicate, and the **G₂ phase**, in which the cell organelles are copied and cytoplasmic volume is increased. If cells rest prior to division they enter a G₀ state where they can remain for long periods of time. The number of cells at each stage of the cell cycle can be assessed by exposing cells to a chemical or radiolabel that gets incorporated into newly generated DNA or by flow cytometry.

The cell cycle is controlled by two **checkpoints** which act as brakes to coordinate the division process at the end of the G₁ and G₂ phases. Two major classes of molecules control these checkpoints, **cyclin-dependent protein kinases** (Cdk), which phosphorylate downstream protein targets, and **cyclins**, which bind to Cdks and regulate their activity. An example of the importance of these systems is demonstrated by mantle cell lymphoma which results from the constitutive activation of cyclin D1 as a result of a chromosomal translocation (see p. 223).

Transcription factors

Transcription factors regulate gene expression by controlling the transcription of specific genes or gene families (Fig. 1.8). Typically, they contain at least two domains: a **DNA-binding domain**, such as a leucine zipper or helix–loop–helix motif which binds to a specific DNA sequence, and an **activation domain**, which contributes to assembly of the transcription complex at a gene promoter. Mutation, deletion or translocation of transcription factors underlie many cases of haematological neoplasms (see Chapter 11).

Epigenetics

This refers to changes in DNA and chromatin that affect gene expression other than those that affect DNA sequence.

Cellular DNA is packaged by wrapping it around histones, a group of specialized nuclear proteins. The complex is tightly compacted as chromatin. In order for the DNA code to be read, transcription factors and other proteins need to physically attach to DNA. Histones act as custodians for this access and so for gene expression. Histones may be modified by methylation, acetylation and phosphorylation which can result in increased or decreased gene expression and so changes in cell phenotype. Epigenetics also includes changes to DNA itself, such as methylation which regulates gene expression in normal and tumour tissues. The methylation of cytosine residues to methyl cytosine results in inhibition of gene transcription. The genes *DNMT 3A* and *B* are involved in the methylation, and *TET 1,2,3* and *IDH1* and *2* in the hydroxylation and therefore breakdown of methylcytosine and restoration of gene

expression (see Fig. 16.1). These genes are frequently mutated in the myeloid malignancies (see Chapters 13, 15 and 16).

Apoptosis

Apoptosis (programmed cell death) is a regulated process of physiological cell death in which individual cells are triggered to activate intracellular proteins that lead to the death of the cell. Morphologically it is characterized by cell shrinkage, condensation of the nuclear chromatin, fragmentation of the nucleus and cleavage of DNA at internucleosomal sites. It is an important process for maintaining tissue homeostasis in haemopoiesis and lymphocyte development.

Apoptosis results from the action of intracellular cysteine proteases called **caspases** which are activated following cleavage and lead to endonuclease digestion of DNA and disintegration of the cell skeleton (Fig. 1.9). There are two major pathways by which caspases can be activated. The first is by signalling through membrane proteins such as Fas or TNF receptor via their intracellular death domain. An example of this mechanism is shown by activated cytotoxic T cells expressing Fas ligand which induces apoptosis in target cells. The second pathway is via the release of cytochrome c from mitochondria. Cytochrome c binds to APAF-1 which then activates caspases. DNA damage induced by irradiation or chemotherapy may act through this pathway. The protein p53 has an important

role in sensing DNA damage. It activates apoptosis by raising the cell level of BAX which then increases cytochrome c release (Fig. 1.9). P53 also shuts down the cell cycle to stop the damaged cell from dividing (Fig. 1.7). The cellular level of p53 is rigidly controlled by a second protein, MDM2. Following death, apoptotic cells display molecules that lead to their ingestion by macrophages.

As well as molecules that mediate apoptosis there are several intracellular proteins that protect cells from apoptosis. The best characterized example is BCL-2. BCL-2 is the prototype of a family of related proteins, some of which are anti-apoptotic and some, like BAX, pro-apoptotic. The intracellular ratio of BAX and BCL-2 determines the relative susceptibility of cells to apoptosis (e.g. determines the lifespan of platelets) and may act through regulation of cytochrome c release from mitochondria.

Many of the genetic changes associated with malignant disease lead to a reduced rate of apoptosis and hence prolonged cell survival. The clearest example is the translocation of the *BCL-2* gene to the immunoglobulin heavy chain locus in the t(14;18) translocation in follicular lymphoma (see p. 222). Overexpression of the BCL-2 protein makes the malignant B cells less susceptible to apoptosis. Apoptosis is the normal fate for most B cells undergoing selection in the lymphoid germinal centres.

Several translocations leading to the generation of fusion proteins, such as t(9;22), t(1;14) and t(15;17), also result in inhibition of apoptosis (see Chapter 11). In addition, genes encoding




Figure 1.9 Representation of apoptosis. Apoptosis is initiated via two main stimuli: (i) signalling through cell membrane receptors such as FAS or tumour necrosis factor (TNF) receptor; or (ii) release of cytochrome c from mitochondria. Membrane receptors signal apoptosis through an intracellular death domain leading to activation of caspases which digest DNA. Cytochrome c binds to the cytoplasmic protein Apaf-1 leading to activation of caspases. The intracellular ratio of pro-apoptotic (e.g. BAX) or anti-apoptotic (e.g. BCL-2) members of the BCL-2 family may influence mitochondrial cytochrome c release. Growth factors raise the level of BCL-2 inhibiting cytochrome c release, whereas DNA damage, by activating p53, raises the level of BAX which enhances cytochrome c release.

proteins that are involved in mediating apoptosis following DNA damage, such as p53 and ATM, are also frequently mutated and therefore inactivated in haemopoietic malignancies.

Necrosis is death of cells and adjacent cells due to ischaemia, chemical trauma or hyperthermia. The cells swell, the plasma


membrane loses integrity. There is usually an inflammatory infiltrate in response to spillage of cell contents. Autophagy is the digestion of cell organelles by lysosomes. It may be involved in cell death but in some situations also in maintaining cell survival by recycling nutrients.

SUMMARY

- Haemopoiesis (blood cell formation) arises from pluripotent stem cells in the bone marrow. Stem cells give rise to progenitor cells which, after cell divisions and differentiation, form red cells, granulocytes (neutrophils, eosinophils and basophils), monocytes, platelets and B and T lymphocytes.
- Haemopoietic tissue occupies about 50% of the marrow space in normal adult marrow. Haemopoiesis in adults is confined to the central skeleton but in infants and young children haemopoietic tissue extends down the long bones of the arms and legs.
- Stem cells reside in the bone marrow in niches formed by stromal cells and circulate in the blood.
- Growth factors attach to specific cell receptors and produce a cascade of phosphorylation events to the cell nucleus. Transcription factors carry the message to those genes that are to be 'switched on', to stimulate cell division, differentiation, functional activity or suppress apoptosis.
- Adhesion molecules are a large family of glycoproteins that mediate attachment of marrow precursors and mature leucocytes and platelets to extracellular matrix, endothelium and to each other.
- Epigenetics refers to changes in DNA and chromatin that affect gene expression other than those that affect DNA sequence. Histone modification and DNA methylation are two important examples relevant to haemopoiesis and haematological malignancies.
- Transcription factors are molecules that bind to DNA and control the transcription of specific genes or gene families.
- Apoptosis is a physiological process of cell death resulting from activation of caspases. The intracellular ratio of pro-apoptotic proteins (e.g. BAX) to anti-apoptotic proteins (e.g. BCL-2) determines the cell susceptibility to apoptosis.



Now visit www.wileyessential.com/haematology to test yourself on this chapter.




CHAPTER 2

Erythropoiesis and general aspects of anaemia

Key topics

■ Blood cells	12
■ Erythropoietin	13
■ Haemoglobin	16
■ Red cell metabolism	18
■ Red cell membrane	18
■ Clinical features of anaemia	20
■ Classification and laboratory findings in anaemia	21
■ Assessment of erythropoiesis	25



Blood cells

All the circulating blood cells derive from pluripotent stem cells in the marrow. They divide into three main types. The most numerous are red cells which are specialized for carriage of oxygen from the lungs to the tissues and of carbon dioxide in the reverse direction (Table 2.1). They have a 4-month lifespan, whereas the smallest cells, platelets involved in haemostasis, circulate for only 10 days. The white cells are made up of four types of phagocyte, neutrophils, eosinophils, basophils and monocytes, which protect against bacterial and fungal infections, and of lymphocytes, which include B cells, involved in antibody production, and T cells (CD4 helper and CD8 suppressor), concerned with the immune response and in protection against viruses and other foreign cells. White cells have a wide range of lifespan (Table 2.1).

The red cells and platelets are counted and their diameter and other parameters measured by an automated cell counter (Fig. 2.1). This also enumerates the different types of white cell by flow cytometry and detects abnormal cells.

We each make approximately 10^{12} new erythrocytes (red cells) each day by the complex and finely regulated process of erythropoiesis. Erythropoiesis passes from the stem cell through the progenitor cells, colony-forming




Figure 2.1 Automated blood cell counter. *Source:* Mehta AB & Hoffbrand AV (2014) *Haematology at a Glance*, 4th edn. Reproduced with permission of John Wiley & Sons.

Table 2.1 The blood cells.

Cell	Diameter (µm)	Lifespan in blood	Number	Function
Red cells 	6–8	120 days	Male: $4.5\text{--}6.5 \times 10^{12}/\text{L}$ Female: $3.9\text{--}5.6 \times 10^{12}/\text{L}$	Oxygen and carbon dioxide transport
Platelets 	0.5–3.0	10 days	$140\text{--}400 \times 10^9/\text{L}$	Haemostasis
Phagocytes				
Neutrophils 	12–15	6–10 h	$1.8\text{--}7.5 \times 10^9/\text{L}$	Protection from bacteria, fungi
Monocytes 	12–20	20–40 h	$0.2\text{--}0.8 \times 10^9/\text{L}$	Protection from bacteria, fungi
Eosinophils 	12–15	Days	$0.04\text{--}0.44 \times 10^9/\text{L}$	Protection against parasites
Basophils 	12–15	Days	$0.01\text{--}0.1 \times 10^9/\text{L}$	
Lymphocytes   B T	7–9 (resting) 12–20 (active)	Weeks or years	$1.5\text{--}3.5 \times 10^9/\text{L}$	B-cells: immunoglobulin synthesis T-cells: protection against viruses; immune functions




Figure 2.2 Erythroblasts (normoblasts) at varying stages of development. The earlier cells are larger, with more basophilic cytoplasm and a more open nuclear chromatin pattern. The cytoplasm of the later cells is more eosinophilic as a result of haemoglobin formation.

unit granulocyte, erythroid, monocyte and megakaryocyte (CFU_{GEMM}), burst-forming unit erythroid (BFU_E) and erythroid CFU (CFU_E) (Fig. 2.2), to the first recognizable erythrocyte precursor in the bone marrow, the pronormoblast. This process occurs in an erythroid niche in which about 30 erythroid cells at various stages of development surround a central macrophage.

The pronormoblast is a large cell with dark blue cytoplasm, a central nucleus with nucleoli and slightly clumped chromatin (Fig. 2.2). It gives rise to a series of progressively smaller normoblasts by a number of cell divisions. They also contain progressively more haemoglobin (which stains pink) in the cytoplasm; the cytoplasm stains paler blue as it loses its RNA and protein synthetic apparatus while nuclear chromatin becomes more condensed (Figs 2.2 and 2.3). The nucleus is finally extruded from the late normoblast within the marrow and a reticulocyte results, which still contains some ribosomal RNA and is still able to synthesize haemoglobin (Fig. 2.4). This cell is slightly larger than a mature red cell, and circulates in the peripheral blood for 1–2 days before maturing, when

RNA is completely lost. A completely pink-staining mature erythrocyte results which is a non-nucleated biconcave disc. One pronormoblast usually gives rise to 16 mature red cells (Fig. 2.3). Nucleated red cells (normoblasts) are not present in normal human peripheral blood (Fig. 2.4). They appear in the blood if erythropoiesis is occurring outside the marrow (extramedullary erythropoiesis) and also with some marrow diseases.

Erythropoietin

Erythropoiesis is regulated by the hormone erythropoietin. Erythropoietin is a heavily glycosylated polypeptide. Normally, 90% of the hormone is produced in the peritubular interstitial cells of the kidney and 10% in the liver and elsewhere. There are no preformed stores and the stimulus to erythropoietin production is the oxygen (O_2) tension in the tissues of the kidney (Fig. 2.5). Hypoxia induces synthesis of hypoxia-inducible factors ($\text{HIF-1}\alpha$ and β), which stimulate erythropoietin production and also new vessel formation and transferrin




Figure 2.3 The amplification and maturation sequence in the development of mature red cells from the pronormoblast.

	Normoblast	Reticulocyte	Mature RBC
Nuclear DNA	Yes	No	No
RNA in cytoplasm	Yes	Yes	No
In marrow	Yes	Yes	Yes
In blood	No	Yes	Yes

Figure 2.4 Comparison of the DNA and RNA content, and marrow and peripheral blood distribution, of the erythroblast (normoblast), reticulocyte and mature red blood cell (RBC).

receptor synthesis, and reduces hepcidin synthesis, increasing iron absorption. Von Hippel-Lindau (VHL) protein breaks down HIFs and PHD2 hydroxylates HIF-1 α allowing VHL binding (Fig. 2.5). Abnormalities in these proteins may cause polycythaemia (see Chapter 15).

Erythropoietin production therefore increases in anaemia, and also when haemoglobin for some metabolic or structural

reason is unable to give up O₂ normally, when atmospheric O₂ is low or when defective cardiac or pulmonary function or damage to the renal circulation affects O₂ delivery to the kidney.

Erythropoietin stimulates erythropoiesis by increasing the number of progenitor cells committed to erythropoiesis. The transcription factor GATA-2 is involved in initiating erythroid differentiation from pluripotent stem cells. Subsequently the transcription factors GATA-1 and FOG-1 are activated by erythropoietin receptor stimulation and are important in enhancing expression of erythroid-specific genes (e.g. globin, haem biosynthetic and red cell membrane proteins) and also enhancing expression of anti-apoptotic genes and of the transferrin receptor (CD71). Late BFU_E and CFU_E, which have erythropoietin receptors, are stimulated to proliferate, differentiate and produce haemoglobin. The proportion of erythroid cells in the marrow increases and, in the chronic state, there is anatomical expansion of erythropoiesis into fatty marrow and sometimes into extramedullary sites. In infants, the marrow cavity may expand into cortical bone resulting in bone deformities with frontal bossing and protrusion of the maxilla (see p. 78).

Conversely, increased O₂ supply to the tissues (because of an increased red cell mass or because haemoglobin is able to release its O₂ more readily than normal) reduces the erythropoietin drive. Plasma erythropoietin levels can be valuable in clinical diagnosis. They are high in anaemia unless this is due to renal failure and if a tumour secreting erythropoietin is present, but low in severe renal disease or polycythaemia vera (Fig. 2.6).




Figure 2.5 The production of erythropoietin by the kidney in response to its oxygen (O₂) supply. Erythropoietin stimulates erythropoiesis and so increases O₂ delivery. BFU_E, erythroid burst-forming unit; CFU_E, erythroid colony-forming unit. Hypoxia induces hypoxia inducible factors (HIFs) α and β , which stimulate erythropoietin production. Von-Hippel-Lindau (VHL) protein breaks down HIFs. PHD2 (prolyl hydroxylase) hydroxylates HIF-1 α allowing VHL binding to HIFs. Mutations in VHL, PHD2 or HIF-1 α underlie congenital polycythaemia (see p. 171).




Figure 2.6 The relation between erythropoietin (EPO) in plasma and haemoglobin concentration. Anaemias exclude conditions shown to be associated with impaired production of EPO. Source: Modified from M. Pippard et al. (1992) *B J Haematol* 82: 445. Reproduced with permission of John Wiley & Sons.

Indications for erythropoietin therapy

Recombinant erythropoietin is needed for treating anaemia resulting from renal disease or from various other causes. It is given subcutaneously either three times weekly or once every 1–2 weeks or every 4 weeks, depending on the indication and on the preparation used (erythropoietin alpha or beta, darbepoetin alpha (a heavily glycosylated longer-acting form), or Micera the longest-acting preparation). The main indication is end-stage renal disease (with or without dialysis). The patients often also need oral or intravenous iron. Other uses are listed in Table 2.2. The haemoglobin level and quality of life may be improved. A low serum erythropoietin level prior to treatment is valuable in predicting an effective response. Side-effects include a rise in blood pressure, thrombosis and local injection site reactions. It has been associated with progression of some tumours which express Epo receptors.

The marrow requires many other precursors for effective erythropoiesis. These include metals such as iron and cobalt, vitamins (especially vitamin B₁₂, folate, vitamin C, vitamin E, vitamin B₆, thiamine and riboflavin) and hormones such as androgens and thyroxine. Deficiency in any of these may be associated with anaemia.

Haemoglobin

Haemoglobin synthesis

The main function of red cells is to carry O₂ to the tissues and to return carbon dioxide (CO₂) from the tissues to the lungs. In order to achieve this gaseous exchange they contain the specialized protein haemoglobin. Each molecule of normal adult haemoglobin A (Hb A) (the dominant haemoglobin in blood after the age of 3–6 months) consists of four polypeptide chains, $\alpha_2\beta_2$, each with its own haem group. Normal adult blood also contains small quantities of two other haemoglobins: Hb F and Hb A₂. These also contain α chains, but with γ and δ chains, respectively, instead of β (Table 2.3). The synthesis of the various globin chains in the fetus and adult is discussed in more detail in Chapter 7.

Table 2.2 Clinical uses of erythropoietin.

Anaemia of chronic renal disease
Myelodysplastic syndrome
Anaemia associated with malignancy and chemotherapy
Anaemia of chronic diseases, e.g. rheumatoid arthritis
Anaemia of prematurity
Perioperative uses

Table 2.3 Normal haemoglobins in adult blood.

	Hb A	Hb F	Hb A ₂
Structure	$\alpha_2\beta_2$	$\alpha_2\gamma_2$	$\alpha_2\delta_2$
Normal (%)	96–98	0.5–0.8	1.5–3.2




Figure 2.7 Haemoglobin synthesis in the developing red cell. The mitochondria are the main sites of protoporphyrin synthesis, iron (Fe) is supplied from circulating transferrin; globin chains are synthesized on ribosomes. δ -ALA, δ -aminolaevulinic acid; CoA, coenzyme A.

Haem synthesis occurs largely in the mitochondria by a series of biochemical reactions commencing with the condensation of glycine and succinyl coenzyme A under the action of the key rate-limiting enzyme δ -aminolaevulinic acid (ALA) synthase (Fig. 2.7). Pyridoxal phosphate (vitamin B₆) is a coenzyme for this reaction. Ultimately, protoporphyrin combines with iron in the ferrous (Fe²⁺) state to form haem (Fig. 2.8).




Figure 2.8 The structure of haem.




Figure 2.9 The oxygenated and deoxygenated haemoglobin molecule. α , β , globin chains of normal adult haemoglobin (Hb A). 2,3-DPG, 2,3-diphosphoglycerate.

A tetramer of four globin chains each with its own haem group in a 'pocket' is then formed to make up a haemoglobin molecule (Fig. 2.9).

Haemoglobin function

The red cells in systemic arterial blood carry O_2 from the lungs to the tissues and return in venous blood with CO_2 to the lungs. As the haemoglobin molecule loads and unloads O_2 the individual globin chains move on each other (Fig. 2.9). The $\alpha_1\beta_1$ and $\alpha_2\beta_2$ contacts stabilize the molecule. When O_2 is unloaded the β chains are pulled apart, permitting entry of the metabolite 2,3-diphosphoglycerate (2,3-DPG) resulting in a lower affinity of the molecule for O_2 . This movement is responsible for the sigmoid form of the haemoglobin O_2 dissociation curve (Fig. 2.10). The P_{50} (i.e. the partial pressure of O_2 at which haemoglobin is half saturated with O_2) of normal blood is 26.6 mmHg. With increased affinity for O_2 , the curve shifts to the left (i.e. the P_{50} falls) while with decreased affinity for O_2 , the curve shifts to the right (i.e. the P_{50} rises).

Normally, *in vivo*, O_2 exchange operates between 95% saturation (arterial blood) with a mean arterial O_2 tension of 95 mmHg and 70% saturation (venous blood) with a mean venous O_2 tension of 40 mmHg (Fig. 2.10).

The normal position of the curve depends on the concentration of 2,3-DPG, H^+ ions and CO_2 in the red cell and on the structure of the haemoglobin molecule. High concentrations of 2,3-DPG, H^+ or CO_2 , and the presence of sickle haemoglobin (Hb S), shift the curve to the right (oxygen is given up more easily) whereas fetal haemoglobin (Hb F) – which is unable to bind 2,3-DPG – and certain rare abnormal haemoglobins associated with polycythaemia shift the curve to the left because they give up O_2 less readily than normal.




Figure 2.10 The haemoglobin oxygen (O_2) dissociation curve. 2,3-DPG, 2,3-diphosphoglycerate.

Methaemoglobinaemia

This is a clinical state in which circulating haemoglobin is present with iron in the oxidized (Fe^{3+}) instead of the usual Fe^{2+} state. It may arise because of a hereditary deficiency of methaemoglobin reductase deficiency or inheritance of a structurally abnormal haemoglobin (Hb M). Hb Ms contain an amino acid substitution affecting the haem pocket of the globin chain. Toxic methaemoglobinaemia (and/or sulphaemoglobinaemia) occurs when a drug or other toxic substance oxidizes haemoglobin. In all these states, the patient is likely to show cyanosis.

The red cell

In order to carry haemoglobin into close contact with the tissues and for successful gaseous exchange, the red cell, 8 μ m in diameter, must be able: to pass repeatedly through the microcirculation whose minimum diameter is 3.5 μ m, to maintain haemoglobin in a reduced (ferrous) state and to maintain osmotic equilibrium despite the high concentration of protein (haemoglobin) in the cell. A single journey round the body takes 20 seconds and its total journey throughout its 120-day lifespan has been estimated to be 480 km (300 miles). To fulfil these functions, the cell is a flexible biconcave disc with an ability to generate energy as adenosine triphosphate (ATP) by the anaerobic glycolytic (Embden–Meyerhof) pathway (Fig. 2.11) and to




Figure 2.11 The Embden–Meyerhof glycolytic pathway. The Luebering–Rapoport shunt regulates the concentration of 2,3-diphosphoglycerate (2,3-DPG) in the red cell. ATP, adenosine triphosphate; NAD, NADH, nicotinamide adenine dinucleotide; PG, phosphoglycerate.

generate reducing power as nicotinamide adenine dinucleotide (NADH) by this pathway and as reduced nicotinamide adenine dinucleotide phosphate (NADPH) by the hexose monophosphate shunt (see Fig. 6.6).

Red cell metabolism

Embden–Meyerhof pathway

In this series of biochemical reactions, glucose that enters the red cell from plasma by facilitated transfer is metabolized to lactate (Fig. 2.11). For each molecule of glucose used, two molecules of ATP and thus two high-energy phosphate bonds are generated. This ATP provides energy for maintenance of red cell volume, shape and flexibility.

The Embden–Meyerhof pathway also generates NADH, which is needed by the enzyme methaemoglobin reductase to reduce functionally dead methaemoglobin containing ferric iron (produced by oxidation of approximately 3% of haemoglobin each day) to functionally active, reduced haemoglobin containing ferrous ions. The Luebering–Rapoport shunt, or side arm, of this pathway (Fig. 2.11) generates 2,3-DPG, important in the regulation of haemoglobin's oxygen affinity (Fig. 2.9).

Hexose monophosphate (pentose phosphate) shunt

Approximately 10% of glycolysis occurs by this oxidative pathway in which glucose-6-phosphate is converted to 6-phosphogluconate and so to ribulose-5-phosphate (see Fig. 6.6). NADPH is generated and is linked with glutathione which maintains sulphhydryl (SH) groups intact in the cell, including those in haemoglobin and the red cell membrane. In one of the most common inherited abnormalities of red cells, glucose-6-phosphate dehydrogenase (G6PD) deficiency, the red cells are extremely susceptible to oxidant stress (see p. 66).

Red cell membrane

The red cell membrane comprises a lipid bilayer, integral membrane proteins and a membrane skeleton (Fig. 2.12). Approximately 50% of the membrane is protein, 20% phospholipids, 20% cholesterol molecules and up to 10% is carbohydrate. Carbohydrates occur only on the external surface while proteins are either peripheral or integral, penetrating the lipid bilayer. Several red cell proteins have been numbered according to their mobility on polyacrylamide gel electrophoresis (PAGE), e.g. band 3, proteins 4.1, 4.2 (Fig. 2.12).




Figure 2.12 The structure of the red cell membrane. Some of the penetrating and integral proteins carry carbohydrate antigens; other antigens are attached directly to the lipid layer.

The membrane skeleton is formed by structural proteins that include α and β spectrin, ankyrin, protein 4.1 and actin. These proteins form a horizontal lattice on the internal side of the red cell membrane and are important in maintaining the biconcave shape. Spectrin is the most abundant and consists of two chains, α and β , wound around each other to form heterodimers which then self-associate head-to-head to form tetramers. These tetramers are linked at the tail end to actin and are attached to protein band 4.1. At the head end, the β spectrin chains attach to ankyrin which connects to band 3, the transmembrane protein that acts as an anion channel ('vertical connections') (Fig. 2.12). Protein 4.2 enhances this interaction.

Defects of the membrane proteins explain some of the abnormalities of shape of the red cell membrane (e.g. hereditary spherocytosis and elliptocytosis) (see Chapter 6), while alterations in lipid composition because of congenital or acquired abnormalities in plasma cholesterol or phospholipid may be associated with other membrane abnormalities (see Fig. 2.16).

Anaemia

This is defined as a reduction in the haemoglobin concentration of the blood below normal for age and sex (Table 2.4). Although normal values can vary between laboratories,

typical values would be less than 135 g/L in adult males and less than 115 g/L in adult females (Fig. 2.13). From the age of 2 years to puberty, less than 110 g/L indicates anaemia. As newborn infants have a high haemoglobin level, 140 g/L is taken as the lower limit at birth (Fig. 2.13).

Alterations in total circulating plasma volume as well as of total circulating haemoglobin mass determine the haemoglobin concentration. Reduction in plasma volume (as in dehydration) may mask anaemia or even cause (apparent, pseudo) polycythaemia (see p. 168); conversely, an increase in plasma volume (as with splenomegaly or pregnancy) may cause anaemia even with a normal total circulating red cell and haemoglobin mass.


After acute major blood loss, anaemia is not immediately apparent because the total blood volume is reduced. It takes up to a day for the plasma volume to be replaced and so for the degree of anaemia to become apparent (see p. 345). Regeneration of red cells and haemoglobin mass takes substantially longer. The initial clinical features of major blood loss are therefore a result of reduction in blood volume rather than of anaemia.

Global incidence

The WHO defines anaemia in adults as a haemoglobin less than 130 g/L in males and less than 120 g/L in females. On

Table 2.4 Normal values for blood cells and haematinics.

	Males	Females
Haemoglobin (g/L)	135.0–175.0	115.0–155.0
Red cells (erythrocytes) ($\times 10^{12}/L$)	4.5–6.5	3.9–5.6
PCV (haematocrit) (%)	40–52	36–48
Mean cell volume (MCV) (fL)	80–95	
Mean cell haemoglobin (MCH) (pg)	27–34	
Reticulocyte count ($\times 10^9/L$)	50–150	
White cells (leucocytes)		
Total ($\times 10^9/L$)	4.0–11.0	
Neutrophils ($\times 10^9/L$)	1.8–7.5	
Lymphocytes ($\times 10^9/L$)	1.5–3.5	
Monocytes ($\times 10^9/L$)	0.2–0.8	
Eosinophils ($\times 10^9/L$)	0.04–0.44	
Basophils ($\times 10^9/L$)	0.01–0.1	
Platelets ($\times 10^9/L$)	150–400	
Serum iron ($\mu\text{mol}/L$)	10–30	
Total iron-binding capacity ($\mu\text{mol}/L$)	40–75 (2.0–4.0 g/L as transferrin)	
Serum ferritin** ($\mu\text{g}/L$)	40–340	14–150
Serum vitamin B ₁₂ ** (ng/L)	160–925 (20–680 pmol/L)	
Serum folate** ($\mu\text{g}/L$)	3.0–15.0 (4–30 nmol/L)	
Red cell folate** ($\mu\text{g}/L$)	160–640 (360–1460 nmol/L)	
PCV, packed cell volume.		
* Lower limit $1.5 \times 10^9/L$ in some ethnic groups, e.g. in Middle East and black-skinned people.		
* *Normal ranges differ between different laboratories.		

**Figure 2.13** The lower limit of normal blood haemoglobin concentration in men, women and children of various ages.

this basis, anaemia was estimated in 2010 to occur in about 33% of the global population. Prevalence was greater in females than males at all ages and most frequent in children less than 5 years old. Anaemia was most frequent in South Asia, and Central, West and East Sub-Saharan Africa. The main causes are iron deficiency (hookworm, schistosomiasis), sickle cell diseases, thalassaemia, malaria and the anaemia of chronic disorders (see p. 37).

Clinical features of anaemia

The major adaptations to anaemia are in the cardiovascular system (with increased stroke volume and tachycardia) and in the haemoglobin O_2 dissociation curve. In some patients with quite severe anaemia there may be no symptoms or signs, whereas others with mild anaemia may be severely incapacitated. The presence or absence of clinical features can be considered under four major headings.

- 1. Speed of onset** Rapidly progressive anaemia causes more symptoms than anaemia of slow onset because there is less time for adaptation in the cardiovascular system and in the O_2 dissociation curve of haemoglobin.
- 2. Severity** Mild anaemia often produces no symptoms or signs but these are usually present when the haemoglobin is less than 90 g/L. Even severe anaemia (haemoglobin concentration as low as 60 g/L) may produce remarkably few symptoms, when there is very gradual onset in a young subject who is otherwise healthy.
- 3. Age** The elderly tolerate anaemia less well than the young because normal cardiovascular compensation is impaired.
- 4. Haemoglobin O_2 dissociation curve** Anaemia, in general, is associated with a rise in 2,3-DPG in the red cells and a shift in the O_2 dissociation curve to the right so that oxygen is given up more readily to tissues. This adaptation is particularly marked in some anaemias that either raise 2,3-DPG directly (e.g. pyruvate kinase deficiency [p. 67]) or that are associated with a low-affinity haemoglobin (e.g. Hb S) (see Fig. 2.10).

Symptoms

If the patient does have symptoms these are usually shortness of breath, particularly on exertion, weakness, lethargy, palpitation and headaches. In older subjects, symptoms of cardiac failure, angina pectoris or intermittent claudication or confusion may be present. Visual disturbances because of retinal haemorrhages may complicate very severe anaemia, particularly of rapid onset (Fig. 2.14).

Signs

These may be divided into general and specific. General signs include pallor of mucous membranes or nail beds, which occurs if the haemoglobin level is less than 90 g/L (Fig. 2.15). Conversely, skin colour is not a reliable sign. A hyperdynamic circulation may be present with tachycardia, a bounding pulse, cardiomegaly and a systolic flow murmur especially at the apex. Particularly in the elderly, features of congestive heart failure may be present.




Figure 2.14 Retinal haemorrhages in a patient with severe anaemia (haemoglobin 25 g/L) caused by severe haemorrhage.



(a)



(b)

Figure 2.15 Pallor of the conjunctival mucosa (a) and of the nail bed (b) in two patients with severe anaemia (haemoglobin 60 g/L).

Specific signs are associated with particular types of anaemia, e.g. koilonychia (spoon nails) with iron deficiency, jaundice with haemolytic or megaloblastic anaemias, leg ulcers with sickle cell and other haemolytic anaemias, bone deformities with thalassaemia major.

The association of features of anaemia with excess infections or spontaneous bruising suggest that neutropenia or thrombocytopenia may be present, possibly as a result of bone marrow failure.

Classification and laboratory findings in anaemia

Red cell indices

The most useful classification is that based on red cell indices (Table 2.4) and divides the anaemia into microcytic, normocytic and macrocytic (Table 2.5). As well as suggesting

the nature of the primary defect, this approach may also indicate an underlying abnormality before overt anaemia has developed.

In two common physiological situations, the mean corpuscular volume (MCV) may be outside the normal adult range. In the newborn for a few weeks the MCV is high but in infancy it is low (e.g. 70 fL at 1 year of age) and rises slowly throughout childhood to the normal adult range. In normal pregnancy there is a slight rise in MCV, even in the absence of other causes of macrocytosis (e.g. folate deficiency).

Other laboratory findings

Although the red cell indices will indicate the type of anaemia, further useful information can be obtained from the initial blood sample.

Table 2.5 Classification of anaemia.

Microcytic, hypochromic	Normocytic, normochromic	Macrocytic
MCV <80 fL	MCV 80–95 fL	MCV >95 fL
MCH <27 pg	MCH ≥27 pg	Megaloblastic: vitamin B ₁₂ or folate deficiency Non-megaloblastic: alcohol, liver disease, myelodysplasia, aplastic anaemia, etc. (see Table 5.10)
Iron deficiency	Many haemolytic anaemias	
Thalassaemia Anaemia of chronic disease (some cases) Lead poisoning Sideroblastic anaemia (some cases)	Anaemia of chronic disease (some cases)	
	After acute blood loss	
	Renal disease	
	Mixed deficiencies	
	Bone marrow failure (e.g. post-chemotherapy, infiltration by carcinoma, etc.)	
MCH, mean corpuscular haemoglobin; MCV, mean corpuscular volume.		

Leucocyte and platelet counts

Measurement of these helps to distinguish ‘pure’ anaemia from ‘pancytopenia’ (subnormal levels of red cells, neutrophils and platelets), which suggests a more general marrow defect or destruction of cells (e.g. hypersplenism). In anaemias caused by haemolysis or haemorrhage, the neutrophil and platelet counts are often raised; in infections and leukaemias, the leucocyte count is also often raised and there may be abnormal leucocytes or neutrophil precursors present.

Reticulocyte count

The normal percentage is 0.5–2.5%, and the absolute count $50\text{--}150 \times 10^9/\text{L}$ (Table 2.4). This should rise in anaemia because of erythropoietin increase, and be higher the more severe the anaemia. This is particularly so when there has been time for erythroid hyperplasia to develop in the marrow as in chronic haemolysis. After an acute major haemorrhage there is an erythropoietin response in 6 hours, the reticulocyte count rises within 2–3 days, reaches a maximum in 6–10 days and remains raised until the haemoglobin returns to the normal level. If the reticulocyte count

is not raised in an anaemic patient this suggests impaired marrow function or lack of erythropoietin stimulus (Table 2.6).

Blood film

It is essential to examine the blood film in all cases of anaemia. Abnormal red cell morphology (Fig. 2.16) or red cell inclusions

Table 2.6 Factors impairing the normal reticulocyte response to anaemia.

Marrow diseases, e.g. hypoplasia, infiltration by carcinoma, lymphoma, myeloma, acute leukaemia, tuberculosis
Deficiency of iron, vitamin B ₁₂ or folate
Lack of erythropoietin, e.g. renal disease
Reduced tissue O ₂ consumption, e.g. myxoedema, protein deficiency
Ineffective erythropoiesis, e.g. thalassaemia major, megaloblastic anaemia, myelodysplasia, myelofibrosis
Chronic inflammatory or malignant disease


Red cell abnormality	Causes	Red cell abnormality	Causes
 Normal		 Microspherocyte	Hereditary spherocytosis, autoimmune haemolytic anaemia, septicaemia
 Macrocyte	Liver disease, alcoholism. Oval in megaloblastic anaemia	 Fragments	DIC, microangiopathy, HUS, TTP, burns, cardiac valves
 Target cell	Iron deficiency, liver disease, haemoglobinopathies, post-splenectomy	 Elliptocyte	Hereditary elliptocytosis
 Stomatocyte	Liver disease, alcoholism	 Tear drop poikilocyte	Myelofibrosis, extramedullary haemopoiesis
 Pencil cell	Iron deficiency	 Basket cell	Oxidant damage—e.g. G6PD deficiency, unstable haemoglobin
 Echinocyte	Liver disease, post-splenectomy, storage artefact	 Sickle cell	Sickle cell anaemia
 Acanthocyte	Liver disease, abetalipoproteinaemia, renal failure	 Microcyte	Iron deficiency, haemoglobinopathy

Figure 2.16 Some of the more frequent variations in size (anisocytosis) and shape (poikilocytosis) that may be found in different anaemias. DIC, disseminated intravascular coagulopathy; G6PD, glucose-6-phosphate dehydrogenase; HUS, haemolytic uraemic syndrome; TTP, thrombotic thrombocytopenic purpura.

(Fig. 2.17) may suggest a particular diagnosis. During the blood film examination, white cell abnormalities are sought, platelet number and morphology are assessed and the presence or absence of abnormal cells (e.g. normoblasts, granulocyte precursors or blast cells) is noted.

Bone marrow examination

This is needed when the cause of anaemia or other abnormality of the blood cells cannot be diagnosed from the blood count, film and other blood tests alone. It may be performed by aspiration or trephine biopsy (Fig. 2.18). During bone marrow aspiration a needle is inserted into the marrow and a liquid sample

of marrow is sucked into a syringe. This is then spread on a slide for microscopy and stained by the usual Romanowsky technique. The detail of the developing cells can be examined (e.g. normoblastic or megaloblastic), the proportion of the different cell lines assessed (myeloid:erythroid ratio, the proportion of granulocyte precursors to red cell precursors in the bone marrow, normally 2.5:1 to 12:1), and the presence of cells foreign to the marrow (e.g. secondary carcinoma) observed. The cellularity of the marrow can also be viewed provided fragments are obtained. An iron stain is performed routinely so that the amount of iron in reticuloendothelial stores (macrophages) and as fine granules ('siderotic' granules) in the developing erythroblasts can be assessed (see Fig. 3.10).




Figure 2.17 Red blood cell (RBC) inclusions which may be seen in the peripheral blood film in various conditions. The reticulocyte RNA and Heinz bodies are only demonstrated by supravital staining (e.g. with new methylene blue). Heinz bodies are oxidized denatured haemoglobin. Siderotic granules (Pappenheimer bodies) contain iron. They are purple on conventional staining but blue with Perls' stain. The Howell–Jolly body is a DNA remnant. Basophilic stippling is denatured RNA.




Figure 2.18 (a) The bone marrow aspiration needle and a smear made from a bone marrow aspirate. (b) The bone marrow trephine (biopsy) needle and normal trephine section.

An aspirate sample may also be used for a number of other specialized investigations (Table 2.7).


A trephine biopsy provides a solid core of bone including marrow and is examined as a histological specimen after fixation in formalin, decalcification and sectioning. Usually immunohistology is performed depending on the diagnosis suspected (see Chapter 11). A trephine biopsy specimen is less valuable than aspiration when individual cell detail is to be examined but provides a panoramic view of the marrow from which overall marrow architecture, cellularity and presence of fibrosis or abnormal infiltrates can, with immunohistology, be reliably determined.

Ineffective erythropoiesis

Erythropoiesis is not entirely efficient because approximately 10–15% of developing erythroblasts die within the marrow without producing mature cells. This is termed ineffective erythropoiesis and it is substantially increased in a number of chronic anaemias (Fig. 2.19). The serum unconjugated bilirubin (derived from breaking down haemoglobin) and lactate

Table 2.7 Indications for bone marrow aspiration and trephine biopsy.

	Aspiration	Trephine
Site	Posterior iliac crest (sternum if obese; tibia in infants)	Posterior iliac crest
Stains	Romanowsky; Perls' reaction (for iron)	Haematoxylin and eosin; reticulin (silver stain)
Result available	1–2 hours	1–7 days (according to decalcification method)
Indications	Investigation of unexplained anaemia, neutropenia, thrombocytopenia, suspicion of leukaemia, myeloproliferative disorders, myelodysplasia, aplastic anaemia, lymphoma, myeloma, amyloid, secondary carcinoma, cases of splenomegaly or pyrexia of undetermined cause	
Special tests	Flow cytometry, cytogenetics and molecular tests such as FISH (see p. 131) and DNA or RNA analysis for gene abnormalities. Consider microbiological culture, cytochemical markers and progenitor cell culture	Immunohistological staining

FISH, fluorescence *in situ* hybridization.**Figure 2.19** The relative proportions of marrow erythroblastic activity, circulating red cell mass and red cell lifespan in normal subjects and in three types of anaemia.

dehydrogenase (LDH, derived from breaking down cells) are usually raised when ineffective erythropoiesis is marked. The reticulocyte count is low in relation to the degree of anaemia and to the proportion of erythroblasts in the marrow.

Assessment of erythropoiesis

Total erythropoiesis and the amount of erythropoiesis that is effective in producing circulating red cells can be assessed by examining the bone marrow, haemoglobin level and reticulocyte count.

Total erythropoiesis is assessed from the marrow cellularity and the myeloid:erythroid ratio. This ratio falls and may be reversed when total erythropoiesis is selectively increased.


Effective erythropoiesis is assessed by the reticulocyte count. This is raised in proportion to the degree of anaemia when erythropoiesis is effective, but is low when there is ineffective erythropoiesis or an abnormality preventing normal marrow response (Table 2.6).

SUMMARY

- Erythropoiesis (red cell production) is regulated by erythropoietin, which is secreted by the kidney in response to hypoxia. Erythropoiesis occurs from mixed progenitor cells through a series of nucleated red cell precursors (normoblasts) to a reticulocyte stage, containing RNA but not DNA.
- Various short- or long-acting preparations of erythropoietin are used clinically to treat anaemia in renal failure and other diseases.
- Haemoglobin is the main protein in red cells. It consists of four polypeptide (globin) chains, in adults 2α and 2β , each containing an iron atom bound to protoporphyrin to form haem.
- The red cell has two biochemical pathways for metabolizing glucose, the Embden–Meyerhof, which generates ATP needed for maintenance of red cell shape and flexibility and NADH which prevents oxidation of haemoglobin, and the hexose monophosphate pathway, which generates NADPH, important for maintaining glutathione which keeps cell proteins in the membrane and haemoglobin in the reduced state.
- The red cell membrane consists of a lipid bilayer with a membrane skeleton of penetrating and integral proteins and carbohydrate surface antigens.
- Anaemia is defined as a haemoglobin level in blood below the normal level for age and sex. It is classified according to the size of the red cells into macrocytic, normocytic and microcytic. The reticulocyte count, morphology of the red cells and changes in the white cell and/or platelet count help in the diagnosis of the cause of anaemia.
- The general clinical features of anaemia include shortness of breath on exertion, pallor of mucous membranes and tachycardia.
- Other features relate to particular types of anaemia, e.g. jaundice, leg ulcers.
- Bone marrow examination by aspiration or trephine biopsy may be important in the investigation of anaemia as well as of many other haematological diseases. Special tests, e.g. immunology, cytogenetics, can be performed on the cells obtained.



Now visit www.wileyessential.com/haematology to test yourself on this chapter.




CHAPTER 3

Hypochromic anaemias

Key topics

■ Nutritional and metabolic aspects of iron	28
■ Iron absorption	30
■ Iron deficiency	32
■ Causes of iron deficiency	32
■ Laboratory findings	33
■ Treatment	36
■ Anaemia of chronic disorders	37
■ Sideroblastic anaemia	38



Iron is one of the most common elements in the Earth's crust, yet iron deficiency is the most common cause of anaemia, affecting about 500 million people worldwide. It is particularly frequent in low-income populations, such as in sub-Saharan Africa or South Asia, where the diet can be of poor quality and parasites (e.g. hookworm or schistosomiasis), which cause iron loss due to haemorrhage, may be present. Moreover the body has limited ability to absorb iron. **Iron deficiency is the major cause of a microcytic, hypochromic anaemia, in which the two red cell indices, mean corpuscular volume (MCV) and mean corpuscular haemoglobin (MCH), are reduced and the blood film shows small (microcytic) and pale (hypochromic) red cells.** This appearance is caused by a defect in haemoglobin synthesis (Fig. 3.1). The major differential diagnosis of a microcytic, hypochromic anaemia is thalassaemia, which is considered in Chapter 7, and anaemia of chronic disease, which is dealt with in this chapter.

Nutritional and metabolic aspects of iron

Body iron distribution and transport

The transport and storage of iron is largely mediated by three proteins: transferrin, transferrin receptor 1 (TfR1) and ferritin.

Transferrin molecules can each contain up to two atoms of iron. Transferrin delivers iron to tissues that have transferrin receptors, especially erythroblasts in the bone marrow which incorporate the iron into haemoglobin (Figs 2.7, 3.2). The transferrin is then reutilized. At the end of their life, red cells are broken down in the macrophages of the reticuloendothelial system and the iron is released from haemoglobin, enters the plasma and provides most of the iron on transferrin. Only a small proportion of plasma transferrin iron comes from dietary iron, absorbed through the duodenum and jejunum.

Some iron is stored in the macrophages as ferritin and haemosiderin, the amount varying widely according to overall body iron status. Ferritin is a water-soluble protein-iron complex. It is made up of an outer protein shell, apoferritin, consisting of 22 subunits and an iron-phosphate-hydroxide




Figure 3.1 The causes of a hypochromic microcytic anaemia. These include lack of iron (iron deficiency) or of iron release from macrophages to serum (anaemia of chronic inflammation or malignancy), failure of protoporphyrin synthesis (sideroblastic anaemia) or of globin synthesis (α - or β -thalassaemia). Lead also inhibits haem and globin synthesis.

core. It contains up to 20% of its weight as iron and is not visible by light microscopy.

Haemosiderin is an insoluble protein-iron complex of varying composition containing approximately 37% iron by weight. It is derived from partial lysosomal digestion of ferritin molecules and is visible in macrophages and other cells by light microscopy after staining by Perls' (Prussian blue) reaction (see Fig. 3.10). Iron in ferritin and haemosiderin is in the ferric form. It is mobilized after reduction to the ferrous form. A copper-containing enzyme, caeruloplasmin, catalyses oxidation of the iron to the ferric form for binding to plasma transferrin.

Iron is also present in muscle as myoglobin and in most cells of the body in iron-containing enzymes (e.g. cytochromes or catalase) (Table 3.1). This tissue iron is less likely to become depleted than haemosiderin, ferritin and haemoglobin in states of iron deficiency, but some reduction of these haem-containing enzymes may occur.

Table 3.1 The distribution of body iron.

Amount of iron in average adult	Male (g)	Female (g)	Percentage of total
Haemoglobin	2.4	1.7	65
Ferritin and haemosiderin	1.0 (0.3–1.5)	0.3 (0–1.0)	30
Myoglobin	0.15	0.12	3.5
Haem enzymes (e.g. cytochromes, catalase, peroxidases, flavoproteins)	0.02	0.015	0.5
Transferrin-bound iron	0.004	0.003	0.1




Figure 3.2 Daily iron cycle. Most of the iron in the body is contained in circulating haemoglobin (see Table 3.1) and is reutilized for haemoglobin synthesis after the red cells die. Iron is transferred from macrophages to plasma transferrin and so to bone marrow erythroblasts. Iron absorption is normally just sufficient to make up for iron loss. The dashed line indicates ineffective erythropoiesis.

Regulation of ferritin and transferrin receptor 1 synthesis

The levels of ferritin, TfR1, δ -aminolaevulinic acid synthase (ALA-S) and divalent metal transporter 1 (DMT-1) are linked to iron status so that iron overload causes a rise in tissue ferritin and a fall in TfR1 and DMT-1, whereas in iron deficiency ferritin and ALA-S are low and TfR1 increased. This linkage arises through the binding of an iron regulatory protein (IRP) to iron response elements (IREs) on the ferritin, TfR1, ALA-S and DMT-1 mRNA molecules. Iron deficiency increases the ability of IRP to bind to the IREs whereas iron overload reduces the binding. The site of IRP binding to IREs, whether upstream (5') or downstream (3') from the coding gene, determines whether the amount of mRNA and so protein produced

is increased or decreased (Fig. 3.3). Upstream binding reduces translation, whereas downstream binding stabilizes the mRNA, increasing translation and so protein synthesis.

When plasma iron is raised and transferrin is saturated, the amount of iron transferred to parenchymal cells (e.g. those of the liver, endocrine organs and heart) is increased and this is the basis of the pathological changes associated with iron loading conditions. There may also be free iron in plasma which is toxic to different organs (see Chapter 4).

Hepcidin

Hepcidin is a polypeptide produced by liver cells. It is the major hormonal regulator of iron homeostasis (Fig. 3.4a). It inhibits iron release from macrophages and from intestinal




Figure 3.3 Regulation of transferrin receptor 1 (TfR1), divalent metal transporter 1 (DMT-1) and ferritin expression by iron regulatory protein (IRP) sensing of intracellular iron levels. IRPs are able to bind to stem-loop structures called iron response elements (IREs). IRP binding to the IRE within the 3' untranslated region of TfR1 and DMT-1 leads to stabilization of the mRNA and increased protein synthesis, whereas IRP binding to the IRE within the 5' untranslated region of ferritin and δ -aminolaevulinic acid synthase (ALA-S) mRNA reduces translation. IRPs can exist in two states: at times of high iron levels the IRP binds iron and exhibits a reduced affinity for the IREs, whereas when iron levels are low the binding of IRPs to IREs is increased. In this way synthesis of TfR, ALA-S, DMT-1 and ferritin is coordinated to physiological requirements.

epithelial cells by its interaction with the transmembrane iron exporter, ferroportin. It accelerates degradation of ferroportin mRNA. Raised hepcidin levels therefore profoundly affect iron metabolism by reducing its absorption and release from macrophages.

Control of hepcidin expression

Membrane-bound hemojuvelin (HJV) is a co-receptor with bone morphogenetic protein (BMP) which stimulates hepcidin expression (Fig. 3.4b). A complex between HFE and transferrin receptor 2 (TfR2) promotes HJV binding to BMP. The amount of HFE-TfR2 complex is determined by the degree of iron saturation of transferrin as follows. Diferric transferrin competes with TfR1 for binding to HFE. The more diferric transferrin, the less TfR1 is bound to HFE and more HFE is available to bind to TfR2, with consequently increased hepcidin synthesis. Low concentrations of diferric transferrin, as in iron deficiency, allow HFE binding to TfR1, reducing the amount of HFE able to bind TfR2 and thus reducing hepcidin secretion. HFE also increases BMP expression, directly increasing hepcidin synthesis.

Matriptase 2 digests membrane-bound HJV. In iron deficiency, increased matriptase activity therefore results in decreased hepcidin synthesis. Erythroblasts secrete two proteins, erythroferrone and GDF 15, which suppress hepcidin secretion. In conditions with increased numbers of early erythroblasts in the marrow (e.g. conditions of ineffective erythropoiesis, such as thalassaemia major), iron absorption is increased because of suppression of hepcidin secretion by these proteins. Hypoxia also suppresses hepcidin synthesis, whereas in inflammation interleukin 6 (IL-6) and other cytokines increase hepcidin synthesis (Fig. 3.4a).

Dietary iron

Iron is present in food as ferric hydroxides, ferric-protein and haem-protein complexes. Both the iron content and the proportion of iron absorbed differ from food to food; in general meat, in particular liver, is a better source than vegetables, eggs or dairy foods. The average Western diet contains 10–15 mg iron daily from which only 5–10% is normally absorbed. The proportion can be increased to 20–30% in iron deficiency or pregnancy (Table 3.2) but even in these situations most dietary iron remains unabsorbed.

Iron absorption

Organic dietary iron is partly absorbed as haem and partly broken down in the gut to inorganic iron. Absorption occurs through the duodenum. Haem is absorbed through a receptor, yet to be identified, on the apical membrane of the duodenal enterocyte. Haem is then digested to release iron. Inorganic iron absorption is favoured by factors such as acid and reducing agents that keep iron in the gut lumen in the Fe^{2+} rather than the Fe^{3+} state (Table 3.2). The protein DMT-1 is involved in transfer of iron from the lumen of the gut across the enterocyte

Table 3.2 Iron absorption.

Factors favouring absorption	Factors reducing absorption
Haem iron	Inorganic iron
Ferrous form (Fe^{2+})	Ferric form (Fe^{3+})
Acids (HCl, vitamin C)	Alkalis – antacids, pancreatic secretions
Solubilizing agents (e.g. sugars, amino acids)	Precipitating agents – phytates, phosphates, tea
Reduced serum hepcidin	Increased serum hepcidin
Ineffective erythropoiesis	Decreased erythropoiesis
Pregnancy	Inflammation
Hereditary haemochromatosis	




Figure 3.4 (a) Hepcidin reduces iron absorption and release from macrophages by stimulating degradation of ferroportin. Its synthesis is increased by transferrin saturation and inflammation but reduced by increased erythropoiesis, erythropoietin, hypoxia and matriptase. (b) The proposed mechanism by which the degree of transferrin saturation by iron affects hepcidin synthesis. BMP, bone morphogenetic protein; HJV, hemojuvelin; TFR1 and 2, transferrin receptor 1 and 2. BMP stimulates hepcidin synthesis and this is enhanced by HJV binding to BMP. Diferric transferrin competes with TFR1 for binding of HFE. The more diferric transferrin, the less TFR1 is bound to HFE and more HFE is available to bind to TFR2. The HFE/TFR2 complex promotes HJV binding to BMP and so promotes hepcidin synthesis. Low concentrations of diferric transferrin, as in iron deficiency, allow HFE binding to TFR1, reducing the amount of HFE able to bind TFR2 and thus reducing HJV binding to BMP and so reducing hepcidin secretion. HFE also enhances BMP expression directly but mutated HFE (see Chapter 4) inhibits its expression.

microvilli (Fig. 3.5). Ferroportin at the basolateral surface controls exit of iron from the cell into portal plasma. The amount of iron absorbed is regulated according to the body's needs by changing the levels of DMT-1 and ferroportin. For DMT-1 this occurs by the IRP/IRE binding mechanism (Fig. 3.3), and for ferroportin by hepcidin (Fig. 3.4a).

Ferrireductase present at the apical surface converts iron from the Fe^{3+} to Fe^{2+} state and another enzyme, hephaestin

(ferrioxidase), converts Fe^{2+} to Fe^{3+} at the basal surface prior to binding to transferrin.

Iron requirements

The amount of iron required each day to compensate for losses from the body and for growth varies with age and sex; it is highest in pregnancy, adolescent and menstruating females




Figure 3.5 The regulation of iron absorption. Dietary ferric (Fe^{3+}) iron is reduced to Fe^{2+} and its entry to the enterocyte is through the divalent cation binder DMT-1. Its export into portal plasma is controlled by ferroportin. It is oxidized before binding to transferrin in plasma. Haem is absorbed after binding to its receptor protein.

(Table 3.3). Therefore these groups are particularly likely to develop iron deficiency if there is additional iron loss or prolonged reduced intake.

Iron deficiency

Clinical features

When iron deficiency is developing, the reticuloendothelial stores (haemosiderin and ferritin) become completely depleted before anaemia occurs (Fig. 3.6). As the condition develops, the patient may show the general symptoms and signs of anaemia (see p. 20) and also a painless glossitis, angular stomatitis, brittle, ridged or spoon nails (koilonychia) (Fig. 3.7) and unusual dietary cravings (pica). The cause of the epithelial cell changes is not clear but may be related to reduction of iron-containing enzymes. In children, iron deficiency is particularly significant as it can cause irritability, poor cognitive function and a decline in psychomotor development. There is also evidence that oral or parenteral iron may reduce fatigue in iron-deficient (low serum ferritin) non-anaemic women.

Causes of iron deficiency

In developed countries, chronic blood loss, especially uterine or from the gastrointestinal tract, is the dominant cause of iron deficiency (Table 3.4) and dietary deficiency is rarely a cause on its own. Five hundred millilitres of blood contain approximately 250 mg iron and, despite the increased absorption of food iron at an early stage of iron deficiency, negative iron balance is usual in chronic blood loss.

Increased demands during infancy, adolescence, pregnancy, lactation and in menstruating women account for the high risk of iron deficiency anaemia in these particular clinical groups. Newborn infants have a store of iron derived from delayed clamping of the cord and the breakdown of excess red cells. From 3 to 6 months there is a tendency for negative iron balance because of growth. From 6 months, supplemented formula milk and mixed feeding, particularly with iron-fortified foods, prevents iron deficiency.

In pregnancy increased iron is needed for an increased maternal red cell mass of approximately 35%, transfer of 300 mg of iron to the fetus and because of blood loss at delivery. Although iron absorption is also increased, iron therapy is often needed if the haemoglobin (Hb) falls below 100 g/L

Table 3.3 Estimated daily iron requirements. Units are mg/day.

	Urine, sweat, faeces	Menses	Pregnancy	Growth	Total
Adult male	0.5–1				0.5–1
Postmenopausal female	0.5–1				0.5–1
Menstruating female*	0.5–1	0.5–1			1–2
Pregnant female*	0.5–1		1–2		1.5–3
Children (average)	0.5			0.6	1.1
Female (age 12–15)*	0.5–1	0.5–1		0.6	1.6–2.6

* These groups are more likely to develop iron deficiency.

	Normal	Latent iron deficiency	Iron deficiency anaemia
Red cell iron (peripheral film and indices)	 Normal	 Normal	 Hypochromic, microcytic MCV↓ MCH↓
Iron stores (bone marrow macrophage iron)	 ++	 0	 0

Figure 3.6 The development of iron deficiency anaemia. Reticuloendothelial (macrophage) stores are lost completely before anaemia develops. MCH, mean corpuscular haemoglobin; MCV, mean corpuscular volume.

or the mean cell volume (MCV) is below 82 fL in the third trimester.

Menorrhagia (a loss of 80 mL or more of blood at each cycle) is difficult to assess clinically, although the loss of clots, the use of large numbers of pads or tampons or prolonged periods all suggest excessive loss.

It takes about 8 years for a normal adult male to develop iron deficiency anaemia solely as a result of a poor diet or malabsorption resulting in no iron intake at all. In developed countries inadequate intake or malabsorption are only rarely the sole cause of iron deficiency anaemia. Gluten-induced enteropathy, partial or total gastrectomy and atrophic gastritis (often autoimmune and with *Helicobacter pylori* infection) may, however, predispose to iron deficiency. In developing countries, iron deficiency may occur as a result of a life-long poor diet, consisting mainly of cereals and vegetables. Hookworm

Table 3.4 Causes of iron deficiency.

Chronic blood loss

Uterine

Gastrointestinal, e.g. peptic ulcer, oesophageal varices, aspirin (or other non-steroidal anti-inflammatory drugs) ingestion, gastrectomy, carcinoma of the stomach, caecum, colon or rectum, hookworm, schistosomiasis, angiodysplasia, inflammatory bowel disease, piles, diverticulosis

Rarely, haematuria, haemoglobinuria, pulmonary haemosiderosis, self-inflicted blood loss

Increased demands (see also Table 3.3)

Prematurity

Growth

Pregnancy

Erythropoietin therapy

Malabsorption

Gluten-induced enteropathy, gastrectomy, autoimmune gastritis

Poor diet

A major factor in many developing countries but rarely the sole cause in developed countries

may aggravate iron deficiency, as may repeated pregnancies or growth and menorrhagia in young females.

Laboratory findings

These are summarized and contrasted with those in other hypochromic anaemias in Table 3.7.



(a)



(b)

Figure 3.7 Iron deficiency anaemia. **(a)** Koilonychia: typical 'spoon' nails. **(b)** Angular cheilosis: fissuring and ulceration of the corner of the mouth.

Red cell indices and blood film

Even before anaemia occurs, the red cell indices fall and they fall progressively as the anaemia becomes more severe. The blood film shows hypochromic, microcytic cells with occasional target cells and pencil-shaped poikilocytes (Fig. 3.8). The reticulocyte count is low in relation to the degree of anaemia. When iron deficiency is associated with severe folate or vitamin B₁₂ deficiency, a 'dimorphic' film occurs with a dual population of red cells of which one is macrocytic and the other microcytic and hypochromic; the indices may be normal. A dimorphic blood film is also seen in patients with iron deficiency anaemia who have received recent iron therapy and produced a population of new haemoglobinized normal-sized red cells (Fig. 3.9) and when the patient has been transfused. The platelet count is often moderately raised in iron deficiency, particularly when haemorrhage is continuing.

Bone marrow iron

Bone marrow examination is not essential to assess iron stores except in complicated cases. In iron deficiency anaemia there is a complete absence of iron from stores (macrophages) and from developing erythroblasts (Fig. 3.10). The erythroblasts are small and have a ragged cytoplasm.

Serum iron and total iron-binding capacity

The serum iron falls and total iron-binding capacity (TIBC) rises so that the TIBC is less than 20% saturated (Fig. 3.11). This contrasts both with the anaemia of chronic disorders (see




Figure 3.8 The peripheral blood film in severe iron deficiency anaemia. The cells are microcytic and hypochromic with occasional target cells.




Figure 3.9 Dimorphic blood film in iron deficiency anaemia responding to iron therapy. Two populations of red cells are present: one microcytic and hypochromic, the other normocytic and well haemoglobinized.

below), when the serum iron and the TIBC are both reduced, and with other hypochromic anaemias where the serum iron is normal or even raised.

Serum ferritin

A small fraction of body ferritin circulates in the serum, the concentration being related to tissue, particularly reticulo-endothelial, iron stores. The normal range in men is higher

than in women (Fig. 3.11). In iron deficiency anaemia the serum ferritin is very low while a raised serum ferritin indicates iron overload or excess release of ferritin from damaged tissues or an acute phase response (e.g. in inflammation). The serum ferritin is normal or raised in the anaemia of chronic disorders.

Investigation of the cause of iron deficiency (Fig. 3.12)

In premenopausal women, menorrhagia and/or repeated pregnancies are the usual causes of the deficiency. If these are not present, other causes must be sought. In some patients with menorrhagia a clotting or platelet abnormality (e.g. von Willebrand disease) is present. In men and postmenopausal women, gastrointestinal blood loss is the main cause of iron deficiency and the exact site is sought from the clinical history, physical and rectal examination, by occult blood tests, and by appropriate use of upper and lower gastrointestinal endoscopy and/or radiology (e.g. computed tomography (CT) of the pneumocolon) or virtual colonoscopy using the 3D colon system (Figs 3.12 and 3.13). Tests for parietal cell antibodies, *Helicobacter* infection and serum gastrin level may help to diagnose autoimmune gastritis. In difficult cases a camera in a capsule can be swallowed which relays pictures of the gastrointestinal tract electronically. Tests for transglutaminase antibodies and duodenal biopsy to look for gluten-induced enteropathy can be valuable. Hookworm ova are sought in stools of subjects from areas where this infestation occurs. Rarely, a coeliac axis angiogram is needed to demonstrate angiodysplasia.

If gastrointestinal blood loss is excluded, loss of iron in the urine as haematuria or haemosiderinuria (resulting from




Figure 3.10 Bone marrow iron assessed by Perls' stain. **(a)** Normal iron stores indicated by blue staining in the macrophages. Inset: normal siderotic granule in erythroblast. **(b)** Absence of blue staining (absence of haemosiderin) in iron deficiency. Inset: absence of siderotic granules in erythroblasts.




Figure 3.11 The serum iron, unsaturated serum iron-binding capacity (UIBC) and serum ferritin in normal subjects and in those with iron deficiency, anaemia of chronic disorders and iron overload. The total iron-binding capacity (TIBC) is made up of the serum iron and the UIBC. In some laboratories, the transferrin content of serum is measured directly by immunodiffusion, rather than by its ability to bind iron, and is expressed in g/L. Normal serum contains 2–4 g/L transferrin (1 g/L transferrin = 20 $\mu\text{mol/L}$ binding capacity). Normal ranges for serum iron are 10–30 $\mu\text{mol/L}$; for TIBC, 40–75 $\mu\text{mol/L}$; for serum ferritin, male, 40–340 $\mu\text{g/L}$; female, 14–150 $\mu\text{g/L}$.

chronic intravascular haemolysis) is considered. A normal chest X-ray excludes the rare condition of pulmonary haemosiderosis. Rarely, patients bleed themselves producing iron deficiency.

Treatment

The underlying cause is treated as far as possible. In addition, iron is given to correct the anaemia and replenish iron stores.

Oral iron

The best preparation is ferrous sulphate which is cheap, contains 67 mg iron in each 200-mg tablet and is best given on an empty stomach in doses spaced by at least 6 hours. Ferrous




Figure 3.12 Investigation and management of iron deficiency anaemia. GI, gastrointestinal; TIBC, total iron-binding capacity.




Figure 3.13 Virtual colonoscopy to show carcinoma of colon causing colonic obstruction and iron deficiency.

Table 3.5 Failure of response to oral iron.

Continuing haemorrhage
Failure to take tablets
Wrong diagnosis – especially thalassaemia trait, sideroblastic anaemia
Mixed deficiency – associated folate or vitamin B ₁₂ deficiency
Another cause for anaemia (e.g. malignancy, inflammation)
Malabsorption – coeliac disease, atrophic gastritis, <i>Helicobacter</i> infection
Use of slow-release preparation

fumarate is equally cheap and effective. If side-effects occur (e.g. nausea, abdominal pain, constipation or diarrhoea), these can be reduced by giving iron with food or by using a preparation of lower iron content (e.g. ferrous gluconate which contains less iron (37 mg) per 300-mg tablet). An elixir is available for children. Slow-release preparations should not be used.

Oral iron therapy should be given for long enough both to correct the anaemia and to replenish body iron stores, which usually means for at least 6 months. The haemoglobin should rise at the rate of approximately 20 g/L every 3 weeks. Failure of response to oral iron has several possible causes (Table 3.5) which should all be considered before parenteral iron is used. Iron fortification of the diet in infants in Africa reduces the incidence of anaemia but increases susceptibility to malaria.

Parenteral iron

Many different preparations are available with varying licensing in different countries. The dose is calculated according to body weight and degree of anaemia. Iron dextran (CosmoFer®) can be given as slow intravenous injection or infusion either in small single doses or as a total dose infusion given in 1 day. Ferric carboxymaltose (Ferinject®) and ferric isomaltoside (Monofer) may also be given as a total dose in 1 day by slow intravenous infusion. Ferric hydroxide–sucrose (Venofer®) is administered by slow intravenous injection or infusion, to a maximum of 200 mg iron in each infusion. Ferumoxytol (Feraheme®) is licensed for chronic renal failure. There may be hypersensitivity or anaphylactoid reactions to parenteral iron, especially in those with a previous reaction, multiple drug allergies and severe atopy. If the reaction is severe, it is treated with intravenous hydrocortisone and possibly adrenaline. Parenteral iron is given slowly and only when there are high iron requirements as in gastrointestinal bleeding, severe menorrhagia, chronic haemodialysis, with erythropoietin therapy, and when oral iron is ineffective (e.g. iron malabsorption resulting from gluten-induced enteropathy or atrophic gastritis) or impractical (e.g. active Crohn's disease). The haematological response to parenteral iron is no faster than to adequate

dosage of oral iron but the stores are replenished faster. Intravenous iron has also been found to increase functional capacity and quality of life in some patients with congestive heart failure, even in the absence of anaemia (see p. 326).

Iron refractory iron deficiency anaemia (IRIDA)

Rare autosomal recessive cases of hypochromic microcytic anaemia have been described caused by inherited mutations of matriptase 2, which allow uninhibited hepcidin secretion, or *DMT-1* genes (Figs 3.4 and 3.5). There may be a haematological response to intravenous but usually not to oral iron.

Anaemia of chronic disorders

One of the most common anaemias occurs in patients with a variety of chronic inflammatory and malignant diseases (Table 3.6). The characteristic features are:

- 1 Normochromic, normocytic or mildly hypochromic (MCV rarely <75 fL) indices and red cell morphology.
- 2 Mild and non-progressive anaemia (haemoglobin rarely <90 g/L) – the severity being related to the severity of the disease.
- 3 Both the serum iron and TIBC are reduced.
- 4 The serum ferritin is normal or raised.
- 5 Bone marrow storage (reticuloendothelial) iron is normal but erythroblast iron is reduced (Table 3.7).

The pathogenesis of this anaemia appears to be related to decreased release of iron from macrophages to plasma because of raised serum hepcidin levels, reduced red cell lifespan and an inadequate erythropoietin response to anaemia caused by the effects of cytokines such as IL-1 and tumour necrosis factor (TNF) on erythropoiesis.

The anaemia is corrected by successful treatment of the underlying disease and does not respond to iron therapy. Erythropoietin injections improve the anaemia in some cases. In many conditions this anaemia is complicated by anaemia resulting from other causes (e.g. iron, vitamin B₁₂ or folate deficiency, renal failure, bone marrow failure, hypersplenism, endocrine abnormality, leucoerythroblastic anaemia) and these are discussed in Chapter 29.

Table 3.6 Causes of the anaemia of chronic disorders.

Chronic inflammatory diseases

Infections (e.g. pulmonary abscess, tuberculosis, osteomyelitis, pneumonia, bacterial endocarditis)

Non-infectious (e.g. rheumatoid arthritis, systemic lupus erythematosus and other connective tissue diseases, sarcoidosis, inflammatory bowel disease, liver disease)

Malignant diseases

Carcinoma, lymphoma, sarcoma

Table 3.7 Laboratory diagnosis of a hypochromic anaemia.

	Iron deficiency	Chronic inflammation or malignancy	Thalassaemia trait (α or β)	Sideroblastic anaemia
MCV/ MCH	Reduced in relation to severity of anaemia	Normal or mild reduction	Reduced; very low for degree of anaemia	Usually low in congenital type but MCV usually raised in acquired type
Serum iron	Reduced	Reduced	Normal	Raised
TIBC	Raised	Reduced	Normal	Normal
Serum ferritin	Reduced	Normal or raised	Normal	Raised
Bone marrow iron stores	Absent	Present	Present	Present
Erythroblast iron	Absent	Absent	Present	Ring forms
Haemoglobin electrophoresis	Normal	Normal	Hb A ₂ raised in β form	Normal

MCH, mean corpuscular haemoglobin; MCV, mean corpuscular volume; TIBC, total iron-binding capacity.

Sideroblastic anaemia

This is a refractory anaemia defined by the presence of many pathological ring sideroblasts in the bone marrow (Fig. 3.14). These are abnormal erythroblasts containing




Figure 3.14 Ring sideroblasts with a perinuclear ring of iron granules in sideroblastic anaemia.

numerous iron granules arranged in a ring or collar around the nucleus instead of the few randomly distributed iron granules seen when normal erythroblasts are stained for iron. There is also usually erythroid hyperplasia with ineffective erythropoiesis. Sideroblastic anaemia is diagnosed when 15% or more of marrow erythroblasts are ring sideroblasts. They can be found at lower numbers in a variety of haematological conditions.

Sideroblastic anaemia is classified into different types (Table 3.8) and the common link is a defect in haem synthesis. In the **hereditary forms** the anaemia is usually characterized by a markedly hypochromic and microcytic blood picture. The most common mutations are in the *ALA-S* gene which is on the X chromosome. Pyridoxal-6-phosphate is a coenzyme for ALA-S. Other rare types include an X-linked disease with spinocerebellar degeneration and ataxia, mitochondrial defects (e.g. Pearson's syndrome when there is also pancreatic insufficiency), thiamine-responsive and other autosomal defects. The much more common form is **refractory anaemia with ring sideroblasts**, which is a subtype of myelodysplasia (see Chapter 16). Acquired reversible forms may be due to alcohol, lead and drugs, e.g. isoniazid.

In some patients, particularly with the hereditary type, there is a response to pyridoxine therapy. Folate deficiency may occur and folic acid therapy may also be tried. Other treatments, e.g. erythropoietin, may be tried in the myelodysplasia form (see Chapter 16). In many severe cases, however, repeated blood transfusions are the only method of maintaining a satisfactory haemoglobin concentration, and transfusional

Table 3.8 Classification of sideroblastic anaemia.**Hereditary**

X chromosome linked ALA-S mutation or rarely with spinocerebellar degeneration and ataxia

Usually occurs in males, transmitted by females; also occurs rarely in females

Other rare types (see text)

Acquired*Primary*

Myelodysplasia (refractory anaemia with ring sideroblasts) (see p. 178)

N.B. Ring sideroblast formation (<15% of erythroblasts) may also occur in the bone marrow in:

other malignant diseases of the marrow (e.g. other types of myelodysplasia, myelofibrosis, myeloid leukaemia, myeloma)

drugs, e.g. antituberculous (isoniazid, cycloserine), alcohol, lead

other benign conditions (e.g. haemolytic anaemia, megaloblastic anaemia, malabsorption, rheumatoid arthritis)

ALA-S, δ -aminolaevulinic acid synthase.

iron overload requiring iron chelation therapy becomes a major problem.

Lead poisoning

Lead inhibits both haem and globin synthesis at a number of points. In addition it interferes with the breakdown of RNA by inhibiting the enzyme pyrimidine 5' nucleotidase, causing accumulation of denatured RNA in red cells, the RNA giving an appearance called basophilic stippling on the ordinary (Romanowsky) stain (see Fig. 2.17). The anaemia may be hypochromic or predominantly haemolytic, and the bone marrow may show ring sideroblasts. Free erythrocyte protoporphyrin is raised.

Differential diagnosis of hypochromic anaemia

Table 3.7 lists the laboratory investigations that may be necessary. The clinical history is particularly important as the source of the haemorrhage leading to iron deficiency or the presence of a chronic disease may be revealed. The country of origin and the family history may suggest a possible diagnosis of thalassaemia or other genetic defect of haemoglobin. Physical examination may also be helpful in determining a site of haemorrhage, features of a chronic inflammatory or malignant disease, koilonychia or, in some haemoglobinopathies, an enlarged spleen or bony deformities.

In thalassaemia trait the red cells tend to be small, often with an MCV of 70 fL or less, even when anaemia is mild or absent; the red cell count is usually over $5.5 \times 10^{12}/L$. Conversely, in iron deficiency anaemia the indices fall progressively with the degree of anaemia and when anaemia is mild the indices are normal or only just reduced below normal (e.g. MCV 75–80 fL). In the anaemia of chronic disorders the indices are also not markedly low, an MCV in the range 75–82 fL being usual.

It is usual to perform a serum iron and TIBC measurement, or alternatively serum ferritin estimation, to confirm a diagnosis of iron deficiency. Haemoglobin high-performance liquid chromatography (HPLC) or electrophoresis with an estimation of Hb A₂ and Hb F is carried out in all patients suspected of thalassaemia or other genetic defect of haemoglobin, because of the family history, country of origin, red cell indices and blood film. Iron deficiency or the anaemia of chronic disorders may also occur in these subjects. β -Thalassaemia trait is characterized by a raised Hb A₂ above 3.5%, but in α -thalassaemia trait there is no abnormality on simple haemoglobin studies so the diagnosis is usually made by exclusion of all other causes of hypochromic red cells and by the presence of a red cell count $>5.5 \times 10^{12}/L$. DNA studies can be used to confirm the diagnosis.


Bone marrow examination is essential if a diagnosis of sideroblastic anaemia is suspected but is not usually needed in diagnosis of the other hypochromic anaemias.

SUMMARY

- Iron is present in the body in haemoglobin, myoglobin, haemosiderin and ferritin, and in iron-containing enzymes. Transferrin is the main transport protein in blood.
- Hepcidin is the main regulator of iron absorption and iron release from macrophages.
- Iron metabolism is regulated according to iron status by intracellular iron regulatory proteins and by control of hepcidin synthesis. Hepcidin synthesis is also affected by erythropoiesis and by inflammation.
- Iron deficiency is the most common cause of anaemia throughout the world. The serum ferritin, serum iron and saturation of the iron-binding capacity are reduced.
- In Western countries, it is usually caused by haemorrhage from the gastrointestinal or the female genital tract. Dietary intake is important particularly in underdeveloped countries where hookworm and schistosomiasis are also important causes of blood loss.
- The red cells are hypochromic and microcytic. It is treated by oral or parenteral iron and by treating, as far as possible, the underlying cause.
- Other frequent causes of a hypochromic, microcytic anaemia are the anaemia of chronic disorders, which occurs in patients with chronic inflammatory or malignant diseases, and α - or β -thalassaemia. Less frequent causes include sideroblastic anaemia (some cases) and lead poisoning.
- Sideroblastic anaemias are characterised by ring sideroblasts in the marrow. The most frequent is a sub-type of myelodysplasia.



Now visit www.wileyessential.com/haematology to test yourself on this chapter.


A vertical strip on the left side of the page showing a microscopic view of blood cells. It features several red blood cells (erythrocytes) in various shades of red and pink, some with a biconcave disc shape. There are also some white blood cells (leukocytes) visible, which are larger and have more complex, irregular shapes. The background is a dark blue.

CHAPTER 4

Iron overload

Key topics

■ Assessment of iron status and organ function	42
■ Hereditary (genetic, primary) haemochromatosis	43
■ Transfusional iron overload	44
■ Iron chelation therapy	45



There is no physiological mechanism for eliminating excess iron from the body, so iron absorption is normally regulated to avoid accumulation. Iron overload (haemosiderosis) occurs in disorders associated with excessive absorption or in patients with severe refractory anaemias who receive regular blood transfusion. Excessive iron deposition in tissues may result in serious damage to organs, particularly the heart, liver and endocrine organs. The causes of iron overload are listed in Table 4.1 and of genetic haemochromatosis in Table 4.2.

Assessment of iron status and organ function

The tests that can be performed to assess iron overload and the degree of damage caused by iron are listed in Table 4.3. The serum ferritin is the most widely used assay and is used

Table 4.1 The causes of iron overload.

Increased iron absorption	Hereditary (primary) haemochromatosis Ineffective erythropoiesis, e.g. thalassaemia intermedia Chronic liver disease
Increased iron intake	African siderosis (dietary and genetic)
Repeated red cell transfusions	Transfusion siderosis

Table 4.2 Genetic causes of haemochromatosis and hyperferritinaemia.

Type	Inheritance	Clinical condition	Gene defect
I	AR	Classical hereditary haemochromatosis	HFE
II	AR	Juvenile haemochromatosis	Hemojuvelin Hepcidin
III	AR	Hereditary haemochromatosis	Transferrin receptor 2
IV	AD	Marked increase in RE iron, less hepatic iron	Ferroportin 1
	AD	Hereditary hyperferritinaemia – cataract syndrome (no iron deposition)	Ferritin

AD, autosomal dominant; AR, autosomal recessive; RE, reticuloendothelial.

Table 4.3 Assessment of iron overload.

<i>Assessment of iron stores</i>	
Serum ferritin	
Serum iron and percentage saturation of transferrin (iron-binding capacity)	
Serum non-transferrin bound iron	
Bone marrow biopsy (Perls' stain) for iron within reticuloendothelial stores	
Liver biopsy (parenchymal and reticuloendothelial (RE) stores)	
Liver CT scan or MRI	
Cardiac MRI (T ₂ * or Ferriscan technique)	
Test of deferiprone or deferoxamine urinary iron excretion (assessment of chelatable iron but rarely performed)	
<i>Assessment of tissue damage caused by iron overload</i>	
Cardiac	Clinical; chest X-ray; ECG; 24-h monitor; echocardiography radionuclide (MUGA) scan to check left ventricular ejection fraction at rest and with stress
Liver	Liver function tests, alphafetoprotein; liver biopsy; CT scan or MRI, fibroscan
Endocrine	Clinical examination (growth and sexual development); glucose tolerance test; pituitary gonadotrophin release tests; thyroid, parathyroid, gonadal, adrenal function, growth hormone assays; radiology for bone age; isotopic bone density study

CT, computed tomography; ECG, electrocardiography; MRI, magnetic resonance imaging; MUGA, multiple gated acquisition.

to assess iron overload and for monitoring its treatment. The percentage saturation of transferrin (which is a measure of iron-binding capacity) is also valuable. Serum non-transferrin bound iron is a toxic form of iron that occurs particularly in severe transfusional iron overload. Liver biopsy with staining for iron (Fig. 4.1) and chemical analysis of iron content is useful for assessment of fibrosis and both parenchymal iron (within hepatic cells) and reticuloendothelial iron (within Kupffer cells). Fibroscan is a non-invasive method of assessing liver fibrosis and magnetic resonance imaging (MRI) is the best non-invasive guide to liver and cardiac iron, measured in the T₂* technique. A commercial Ferriscan MRI technique is widely used for measuring liver iron. Serum alphafetoprotein and liver ultrasound are used for screening for hepatocellular carcinoma.




Figure 4.1 Liver biopsy. Iron loading of hepatic parenchymal cells (Perls' stain). Source: Courtesy of Professor A.P. Dhillon.



Figure 4.2 Melanin skin pigmentation. The right hand is of a teenager with iron overload caused by thalassaemia major. The left hand is of her mother who has normal iron status.

Hereditary haemochromatosis

Hereditary haemochromatosis (also called genetic or primary haemochromatosis) is a group of diseases in which there is excessive absorption of iron from the gastrointestinal tract leading to iron overload of the parenchymal cells of the liver (Fig. 4.1), endocrine organs and, in severe cases, the heart.

Most patients are homozygous for a missense mutation in the *HFE* gene which leads to insertion of a tyrosine residue rather than cysteine in the mature protein (C282Y). The allele has a prevalence of approximately 1 in 300 within the white North European population. Only a small proportion of people who are homozygous for the mutation actually present with clinical features of the disease, and these usually show a serum ferritin greater than 1000 µg/L. A second mutation resulting in a histidine to aspartic acid substitution H63D is found with the C282Y mutation in approximately 5% of patients but homozygotes for the H63D mutation do not have the disease.

HFE is involved in hepcidin synthesis and therefore hereditary haemochromatosis caused by *HFE* mutation is due to low serum hepcidin levels (see Fig. 3.4). Low serum hepcidin levels lead to high levels of ferroportin and therefore increased iron absorption and increased release of iron from macrophages. Iron overload therefore develops and damages parenchymal cells such that patients may present in adult life with hepatic disease (fibrosis, cirrhosis, hepatocellular carcinoma), endocrine disturbances such as diabetes mellitus, hypothyroidism or impotence, melanin skin pigmentation (Fig. 4.2) and arthropathy (resulting from pyrophosphate deposition). In some severe cases there is cardiac failure or arrhythmia. Diagnosis is suspected by increased levels of serum iron, serum transferrin saturation and ferritin. It is confirmed by testing for the *HFE* mutation. Liver biopsy may quantify the degree of iron overload and assess liver damage. MRI can also be used to measure liver and cardiac iron.

Treatment is with regular venesection, initially at 1–2-week intervals, with each unit of blood removing 200–250 mg iron. There are differences of opinion as to whether patients without evidence of organ dysfunction due to iron overload should be treated but usually this is done when the ferritin is raised. Venesection is monitored by serum ferritin and the aim is to restore this to normal.

Rarer forms of genetic haemochromatosis are caused by mutations in the genes for hemojuvelin, transferrin receptor 2 and hepcidin (Table 4.2). All three are involved in hepcidin synthesis and the mutations are associated with low levels of hepcidin in serum. They often present as severe iron overload with cardiomyopathy in children, adolescents or young adults. Genetic iron overload in Asian populations is usually due to these mutations rather than mutation of *HFE*. On the other hand, ferroportin gene mutations usually cause reticuloendothelial but not parenchymal cell iron overload but may rarely cause parenchymal overload, depending on the site of the mutation in the ferroportin gene. Mutations of the ferritin light chain gene cause a raised monoclonal serum ferritin with cataracts resulting from ferritin deposition in the eye but no tissue iron overload.

African iron overload

This occurs in sub-Saharan Africa through a combination of increased iron absorption due to a genetic defect, possibly in the ferroportin gene, and consumption of beverages with high iron content due to the use of iron cooking pots.

Thalassaemia intermedia

Moderately severe forms of thalassaemia may lead to increased iron levels even in patients who do not need regular blood transfusions. This is due to increased absorption and may lead to

increased levels of iron in the liver. Iron chelation is indicated if the liver iron concentration is above 5 mg/g dry weight, where serum ferritin reaches 800 µg/L or when the iron leads to organ damage (see also Chapter 7).

Transfusional iron overload

This develops in patients with chronic anaemia who need to have regular blood transfusions. **Each 500 mL of transfused blood contains approximately 250 mg iron and iron overload is inevitable unless iron chelation therapy is given (Table 4.4).** To make matters worse, iron absorption from food is *increased* in β -thalassaemia major and many other anaemias secondary to ineffective erythropoiesis because of inappropriately low serum hepcidin levels. This is thought to be due to release of proteins from early erythroblasts that inhibit hepcidin synthesis (see Fig. 3.4b). Non-transferrin bound iron may appear in plasma, because transferrin is 100% saturated, and cause widespread iron deposition in parenchymal tissues.

Iron damages the **liver** (Fig. 4.3) and the **endocrine organs** with failure of growth, delayed or absent puberty, diabetes mellitus, hypothyroidism and hypoparathyroidism. Skin pigmentation as a result of excess melanin and haemosiderin gives a slate grey appearance even at an early stage of iron overload. Most importantly, iron can damage the **heart**. In the absence of intensive iron chelation, death occurs in the second or third

Table 4.4 Causes of refractory anaemia that may lead to transfusional iron overload.

Congenital	Acquired
β -Thalassaemia major	Myelodysplasia
β -Thalassaemia/Hb E disease	Red cell aplasia
Sickle cell anaemia (some cases)	Aplastic anaemia
Red cell aplasia (Diamond–Blackfan)	Primary myelofibrosis
Sideroblastic anaemia	
Dyserythropoietic anaemia	

decade in thalassaemia major, usually from congestive heart failure or cardiac arrhythmias. T_2^* MRI is a valuable measure of cardiac and liver iron loading (Fig. 4.4). It can detect increased cardiac iron before sensitive tests detect impaired cardiac function. The shorter the relaxation time, the greater the cardiac iron burden and the greater risk of cardiac failure or arrhythmia (Fig. 4.5). Serum ferritin and liver iron correlate poorly with cardiac iron (Figs 4.4 and 4.5). Moreover serum ferritin is raised in viral hepatitis and other inflammatory disorders and should therefore be interpreted in conjunction with more accurate tests of iron status such as T_2^* MRI or liver biopsy.




Figure 4.3 β -Thalassaemia major: needle biopsy of liver. **(a)** Grade IV siderosis with iron deposition in the hepatic parenchymal cells, bile duct epithelium, macrophages and fibroblasts (Perls' stain). **(b)** Reduction of iron excess in liver after intensive chelation therapy.




Figure 4.4 T_2^* magnetic resonance images (MRIs) showing tissue appearance in iron overload: (a) normal volunteer, (b) severe iron overload. Green arrow, normal appearance; red arrow, iron overload. Lack of correlation: liver and cardiac iron in two cases of thalassaemia major (c) and (d).




Figure 4.5 Comparison of T_2^* MRI measurement of cardiac iron and serum ferritin in thalassaemia major patients. There are substantial numbers of patients with very high serum ferritin ($>3000 \mu\text{g/L}$) but normal heart iron ($T_2^* > 20 \text{ msec}$) and conversely, patients with serum ferritin levels $<1000 \mu\text{g/L}$ with severe cardiac iron loading ($T_2^* < 10 \text{ msec}$). Source: Anderson *et al.* (2001) *Eur Heart J* 22: 2171–79.

Iron chelation therapy

Iron chelation therapy is used to treat transfusional iron overload and three effective drugs are available. Thalassaemia major is the most frequent indication worldwide but chelation is also used for heavily transfused patients with the other anaemias listed in Table 4.4.

Deferasirox is given orally once daily and leads to iron loss in the faeces. Skin rashes and transient changes in liver enzymes and rise in serum creatinine are the main side-effects. The ease of administration and its lack of major side-effects have resulted in its widespread use as first choice.

Deferiprone is also an oral chelator and causes predominantly urinary iron excretion. It is usually given in three doses daily. It may be used alone or, if this is inadequate, in combination with deferoxamine infused on one or more days a week, since the drugs have an additive or even synergetic effect on iron excretion. Alone it is more effective than deferoxamine at removing cardiac iron (Fig. 4.6). Side-effects include an arthropathy, agranulocytosis (in about 1%), neutropenia,




Figure 4.6 (a) Reduction of cardiac iron assessed by T₂* MRI is greater in patients treated with deferiprone than deferoxamine. **(b)** Improvement in left ventricular (LV) ejection fraction is greater with deferiprone than with deferoxamine. Source: Pennell D.J *et al.* (2006) *Blood* 107: 3738–44.

gastrointestinal disturbance and, in diabetics, zinc deficiency. Monitoring of the blood count initially weekly is needed in all patients receiving deferiprone.

The third drug, **deferioxamine**, has been in use longest, for over 40 years, but is not active orally. It is usually given by subcutaneous infusion over 8–12 hours for 5–7 days each week; vitamin C can be given to further increase iron excretion. Most iron is lost in the urine but up to one-third is also excreted in the stools. Because of the difficult administration, compliance is a major problem. It may be given on 1 or more days each week in combination with daily deferiprone and it is used intravenously in

patients with severe iron overload at risk of dying from cardiac failure. Side-effects include high tone deafness, retinal damage, bone abnormalities and growth retardation. Patients should have auditory and fundoscopic examinations at regular intervals.

All three chelators can be given in children. Deferasirox is most frequently used and a liquid formulation of deferiprone is available.

Chelation is started in thalassaemia major after 10–15 units have been transfused.

Chelation is given to keep the serum ferritin level between 1000 and 1500 µg/L, when the body iron stores are


approximately 5–10 times normal. MRI assesses cardiac and liver iron more accurately (Fig. 4.5), but serum ferritin measurement is useful in monitoring changes in iron stores. Serial tests of the function of the heart, liver and endocrine organs are also needed if there is organ impairment (Table 4.3).

Life expectancy has improved dramatically for thalassaemia major patients since the introduction of iron chelation. Chelation may even reverse liver, endocrine and cardiac damage in cases where this has developed before chelation is started or is due to inadequate chelation therapy (Table 4.4).

- Iron overload is caused by excessive absorption of iron from food (genetic haemochromatosis) or by repeated blood transfusions in patients with refractory anaemias. Each unit of blood contains 200–250 mg of iron.
- Excess iron absorbed from the gastrointestinal tract accumulates in the parenchymal cells of the liver, endocrine organs and, in severe cases, the heart.
- Transfusional iron overload causes damage to these organs and also iron accumulation in macrophages of the reticuloendothelial system.
- Genetic haemochromatosis is usually caused by homozygous mutation of the *HFE* gene causing C282Y protein change and a low serum hepcidin level. Rarer forms exist caused by mutations of other genes coding for proteins (hemojuvelin, hepcidin, transferrin receptor 2 and ferroportin). Repeated venesections are used to reduce the body iron burden.
- Transfusional iron overload most frequently occurs in thalassaemia major but also in other transfusion dependent refractory anaemias (e.g. some cases of myelodysplasia, sickle cell anaemia, primary myelofibrosis, red cell aplasia and aplastic anaemia).
- Cardiac failure or arrhythmia caused by cardiac siderosis, best detected by MRI, is the most frequent cause of death from transfusional iron overload.
- Treatment is with iron chelating drugs: deferasirox and deferiprone, which are active orally, or deferoxamine, given subcutaneously or intravenously.
- Life expectancy has improved dramatically in thalassaemia major as a result of iron chelation therapy and the use of MRI to accurately measure cardiac and liver iron.



Now visit www.wileyessential.com/haematology to test yourself on this chapter.



CHAPTER 5

Megaloblastic anaemias and other macrocytic anaemias

Key topics

■ Megaloblastic anaemias	49
■ Vitamin B ₁₂	49
■ Folate	51
■ Vitamin B ₁₂ deficiency	52
■ Folate deficiency	53
■ Clinical features of megaloblastic anaemia	53
■ Diagnosis of vitamin B ₁₂ or folate deficiency	56
■ Other megaloblastic anaemias	58
■ Other macrocytic anaemias	58

Introduction to macrocytic anaemia

In macrocytic anaemia the red cells are abnormally large (mean corpuscular volume, MCV >98 fL). There are several causes (see Table 2.5) but they can be broadly subdivided into megaloblastic and non-megaloblastic (Table 5.10), based on the appearance of developing erythroblasts in the bone marrow.

Megaloblastic anaemias

This is a group of anaemias in which the erythroblasts in the bone marrow show a characteristic abnormality – maturation of the nucleus being delayed relative to that of the cytoplasm. The underlying defect accounting for the asynchronous maturation of the nucleus is defective DNA synthesis and this is usually caused by deficiency of vitamin B₁₂ or folate. Less commonly, abnormalities of metabolism of these vitamins or other lesions in DNA synthesis may cause an identical haematological appearance (Table 5.1).

Vitamin B₁₂ (B₁₂, cobalamin)

Vitamin B₁₂ is synthesized in nature by microorganisms; animals acquire it by eating food of animal origin, by internal production from intestinal bacteria (not in humans) or by eating bacterially contaminated foods. The vitamin consists of a small group of compounds, the cobalamins, which have the same basic structure, with a cobalt atom at the centre of a corrin ring which is attached to a nucleotide portion (Fig. 5.1). The vitamin is found in foods of animal origin such as liver, meat, fish and dairy produce but does not occur in fruit, cereals or vegetables (Table 5.2).

Absorption

A normal diet contains a large excess of B₁₂ compared with daily needs (Table 5.2). B₁₂ is released from protein-binding in food and is combined with the glycoprotein **intrinsic factor (IF)**, which is synthesized by the gastric parietal cells. The IF–B₁₂

complex then binds in the ileum to a specific surface receptor for IF, cubilin. This binds to a second protein, amnionless, which directs endocytosis of the cubilin IF–B₁₂ complex in the ileal cell so that B₁₂ is absorbed and IF destroyed (Fig. 5.2).

Transport: the transcobalamins

Vitamin B₁₂ is absorbed into portal blood where it becomes attached to the plasma-binding protein transcobalamin (TC, also called transcobalamin II), which delivers B₁₂ to bone marrow and other tissues. Although TC is the essential plasma protein for transferring B₁₂ into the cells of the body, the amount of B₁₂ on TC is normally very low (<50 ng/L). Congenital TC deficiency causes megaloblastic anaemia because of failure of B₁₂ to enter marrow (and other cells) from plasma but the serum B₁₂ level in TC deficiency is normal. This is because most B₁₂ in plasma is bound to another protein, haptocorrin (also called transcobalamin I). This is a glycoprotein largely synthesized by granulocytes and macrophages. In myeloproliferative diseases where granulocyte production may be greatly increased, the haptocorrin and B₁₂ levels in serum both rise considerably. B₁₂ bound to haptocorrin does not transfer to marrow; it appears to be functionally ‘dead’. Closely related glycoproteins to plasma haptocorrin are present in gastric juice, milk and other body fluids.

Table 5.1 Causes of megaloblastic anaemia.

Vitamin B₁₂ deficiency

Folate deficiency

Abnormalities of vitamin B₁₂ or folate metabolism (e.g. transcobalamin deficiency, nitrous oxide, antifolate drugs)

Other defects of DNA synthesis

Congenital enzyme deficiencies (e.g. orotic aciduria)

Acquired enzyme deficiencies (e.g. alcohol, therapy with hydroxyurea, cytosine arabinoside)





Figure 5.1 The structure of methylcobalamin (methyl B₁₂), the main form of vitamin B₁₂ in human plasma. Other forms include deoxyadenosylcobalamin (ado B₁₂), the main form in human tissues; hydroxocobalamin (hydroxo B₁₂), the main form used in treatment.

Table 5.2 Vitamin B₁₂ and folate: nutritional aspects.

	Vitamin B ₁₂	Folate
Normal daily dietary intake	7–30 µg	200–250 µg
Main foods	Animal produce only	Most, especially liver, greens and yeast
Cooking	Little effect	Easily destroyed
Minimal adult daily requirement	1–2 µg	100–150 µg
Body stores	2–3 mg (sufficient for 2–4 years)	10–12 mg (sufficient for 4 months)
Absorption		
Site	Ileum	Duodenum and jejunum
Mechanism	Intrinsic factor	Conversion to methyltetrahydrofolate
Limit	2–3 µg/day	50–80% of dietary content
Enterohepatic circulation	5–10 µg/day	90 µg/day
Transport in plasma	Most bound to haptocorrin; TC essential for cell uptake	Weakly bound to albumin
Major intracellular physiological forms	Methyl- and deoxyadenosylcobalamin	Reduced polyglutamate derivatives
Usual therapeutic form	Hydroxocobalamin	Folic (pteroylglutamic) acid

TC, transcobalamin; haptocorrin = transcobalamin 1.

**Figure 5.2** The absorption of dietary vitamin B₁₂ after combination with intrinsic factor (IF), through the ileum. Folate absorption occurs through the duodenum and jejunum after conversion of all dietary forms to methyltetrahydrofolate (methyl THF). TC, transcobalamin.




Figure 5.3 The biochemical reactions of vitamin B₁₂ in humans. Ado B₁₂, deoxyadenosylcobalamin; CoA, coenzyme A; THF, tetrahydrofolate.

Biochemical function

Vitamin B₁₂ is a coenzyme for two biochemical reactions: first, as methyl B₁₂ it is a cofactor for methionine synthase, the enzyme responsible for methylation of homocysteine to methionine using methyl tetrahydrofolate (methyl THF) as methyl donor (Fig. 5.3); second, as deoxyadenosyl B₁₂ (ado B₁₂) it assists in conversion of methylmalonyl coenzyme A (CoA) to succinyl CoA (Fig. 5.3).

Folate

Folic (pteroylglutamic) acid is the parent compound of a large group of compounds, the folates, that are derived from it (Fig. 5.4).

Absorption, transport and function

Dietary folates (a complex mixture of variously reduced and polyglutamated folates) are all converted to one compound, methyltetrahydrofolate (methylTHF), a reduced




Figure 5.4 The structure of folic (pteroylglutamic) acid. Dietary folates may contain: (a) additional hydrogen atoms at positions 7 and 8 (dihydrofolate) or 5, 6, 7 and 8 (tetrahydrofolate); (b) a formyl group at N₅ or N₁₀, a methyl group at N₅ or other 1-carbon groups; and (c) additional glutamate moiety attached to the γ -carboxyl group of the glutamate moiety.

monoglutamate form which circulates in plasma. After entering cells this is converted to folate polyglutamate forms by addition of usually four, five or six glutamate moieties (Fig. 5.5).

Folates are needed in a variety of biochemical reactions in the body involving single carbon unit transfer, in amino acid interconversions (e.g. homocysteine conversion to methionine) (Figs 5.3, 5.5) and serine to glycine or in synthesis of purine precursors of DNA.




Figure 5.5 The biochemical basis of megaloblastic anaemia caused by vitamin B₁₂ or folate deficiency. Folate is required in one of its coenzyme forms, 5,10-methylene tetrahydrofolate (THF) polyglutamate, in the synthesis of thymidine monophosphate from its precursor deoxyuridine monophosphate. Vitamin B₁₂ is needed to convert methyl THF, which enters the cells from plasma, to THF, from which polyglutamate forms of folate are synthesized. Dietary folates are all converted to methyl THF (a monoglutamate) by the small intestine. A, adenine; C, cytosine; d, deoxyribose; DHF, dihydrofolate; DP, diphosphate; G, guanine; MP, monophosphate; T, thymine; TP, triphosphate; U, uracil.

Biochemical basis for megaloblastic anaemia (Fig. 5.5)

DNA is formed by polymerization of the four deoxyribonucleoside triphosphates. Folate deficiency is thought to cause megaloblastic anaemia by inhibiting thymidylate synthesis, a rate-limiting step in DNA synthesis in which thymidine monophosphate (dTMP) is synthesized. This reaction needs 5,10-methylene THF polyglutamate as coenzyme.

The role of B₁₂ in DNA synthesis is indirect. B₁₂ is needed in the conversion of methyl THF, which enters marrow and other cells from plasma, to THF. THF (but not methyl THF) is the substrate for folate polyglutamate synthesis. The folate polyglutamates including 5,10-methylene THF polyglutamate are the intracellular folate coenzymes. B₁₂ deficiency therefore indirectly reduces the supply of the critical folate coenzyme involved in thymidylate synthesis (Fig. 5.5). Other congenital or acquired causes of megaloblastic anaemia (e.g. antimetabolite drug therapy) inhibit purine or pyrimidine synthesis at one or other step. The result is a reduced supply of one or other of the four precursors needed for DNA synthesis.

Folate reduction

During the synthesis of dTMP, the folate polyglutamate coenzyme becomes oxidized from the THF to dihydrofolate (DHF) state (Fig. 5.5). Regeneration of active THF coenzymes requires the enzyme DHF reductase. Inhibitors of this enzyme (e.g. methotrexate) therefore inhibit folate reactions, and so DNA synthesis (Fig. 5.5). Methotrexate is a useful drug, mainly in the treatment of malignant (e.g. acute lymphoblastic leukaemia [see p. 192]), or inflammatory disease (e.g. rheumatoid arthritis, psoriasis) with excessive cell turnover. The weaker antagonist, pyrimethamine, is used primarily against malaria. Trimethoprim, active against bacterial DHF reductase but only very weakly against the human enzyme, is used alone or in combination with a sulphonamide, as co-trimoxazole. Toxicity caused by methotrexate or pyrimethamine may be reversed by giving the reduced folate, folinic acid (5-formyl THF).

Vitamin B₁₂ deficiency

In Western countries, severe deficiency is usually caused by (Addisonian) pernicious anaemia (Table 5.3). Less commonly, it may be caused by veganism in which the diet lacks B₁₂ (usually in Hindu Indians), gastrectomy or small intestinal lesions. Causes of mild B₁₂ deficiency are listed in Table 5.3 and discussed below. There is no syndrome of B₁₂ deficiency as a result of increased utilization or loss of the vitamin. The deficiency takes at least 2 years to develop (i.e. the time needed for body stores to deplete at the rate of 1–2 µg/day) when there is severe malabsorption of B₁₂ from

Table 5.3 Causes of severe vitamin B₁₂ deficiency.

Nutritional Especially vegans
Malabsorption <i>Gastric causes</i> Pernicious anaemia Congenital lack or abnormality of intrinsic factor Total or partial gastrectomy
<i>Intestinal causes</i> Intestinal stagnant loop syndrome – jejunal diverticulosis, blind-loop, stricture, etc. Chronic tropical sprue Ileal resection and Crohn's disease Congenital selective malabsorption with proteinuria (autosomal recessive megaloblastic anaemia) Fish tapeworm
Causes of mild vitamin B₁₂ deficiency include poor diet and other causes of malabsorption of vitamin B ₁₂ (malabsorption of food B ₁₂ caused by atrophic gastritis particularly in the elderly, therapy with proton pump inhibitors or metformin), severe pancreatitis, gluten-induced enteropathy and HIV infection. These conditions do not usually lead to vitamin B ₁₂ deficiency sufficient to cause anaemia or neuropathy.

the diet. Nitrous oxide, however, may rapidly inactivate body B₁₂ (p. 58).

Pernicious anaemia (PA)

This is caused by autoimmune attack on the gastric mucosa leading to atrophy of the stomach. The wall of the stomach becomes thin, with a plasma cell and lymphoid infiltrate of the lamina propria. Intestinal metaplasia may occur. There is achlorhydria and secretion of IF is absent or almost absent. Serum gastrin levels are raised. *Helicobacter pylori* infection may initiate an autoimmune gastritis which presents in younger subjects as iron deficiency and in the elderly as PA.

More females than males are affected (1.6:1), with a peak occurrence at 60 years, and there may be associated autoimmune disease, particularly thyroid diseases (Table 5.4). The disease is found in all races but is most common in Northern Europeans and tends to occur in families. There is also an increased incidence of carcinoma of the stomach (approximately 2–3% of all cases of pernicious anaemia).

Antibodies

Ninety per cent of patients show parietal cell antibody in the serum directed against gastric H⁺/K⁺-ATPase. PA is distinguished from simple atrophic or autoimmune gastritis by the presence of antibodies to IF. Fifty per cent of PA patients show in serum an antibody to IF which inhibits IF binding to B₁₂. A

Table 5.4 Pernicious anaemia: associations.

Female	Vitiligo
Blue eyes	Myxoedema
Early greying	Hashimoto's disease
Northern European	Thyrotoxicosis
Familial	Addison's disease
Blood group A	Hypoparathyroidism Hypogammaglobulinaemia Carcinoma of the stomach

second antibody blocking IF attachment to its ileal binding site is less frequent. IF antibodies also occur in PA in gastric juice. IF antibodies are virtually specific for PA but as they occur in the serum of only half of patients, their absence from serum does not exclude the diagnosis. The more common parietal cell antibody is less specific as it occurs quite commonly in older subjects (e.g. 16% of normal women over 60 years) even without PA.

Other causes of vitamin B₁₂ deficiency

Congenital lack or abnormality of IF usually presents at approximately 2 years of age when stores of B₁₂ that were derived from the mother *in utero* have been used up. Specific malabsorption of B₁₂ is due to genetic mutation of the IF-B₁₂ receptor, cubilin, or of amnionless. It usually presents in infancy or childhood and is associated with proteinuria in 90% of cases.

Lesser degrees of B₁₂ deficiency with borderline or low serum B₁₂ levels, but almost always without megaloblastic anaemia, occur resulting from inadequate intake of B₁₂ or from malabsorption of food B₁₂, especially in the elderly with atrophic gastritis, and in association with prolonged therapy with proton pump inhibitors or metformin. Mild B₁₂ deficiency also may occur with the other conditions listed in Table 5.3.

Folate deficiency

This is most often a result of a poor dietary intake of folate alone or in combination with a condition of increased folate utilization or malabsorption (Table 5.5). Excess cell turnover of any sort, including pregnancy, is the main cause of an increased need for folate, because the folate molecule becomes degraded when DNA synthesis, and so thymidylate synthesis, is increased. The mechanism by which anticonvulsants and barbiturates cause the deficiency is still controversial.

Table 5.5 Causes of folate deficiency.

Nutritional

Especially old age, institutions, poverty, famine, special diets, goat's milk anaemia, etc.

Malabsorption

Tropical sprue, gluten-induced enteropathy (adult or child). Possible contributory factor to folate deficiency in some patients with partial gastrectomy, extensive jejunal resection or Crohn's disease

Excess utilization

Physiological

Pregnancy and lactation, prematurity

Pathological

Haematological diseases: haemolytic anaemias, myelofibrosis

Malignant disease: carcinoma, lymphoma, myeloma

Inflammatory diseases: Crohn's disease, tuberculosis, rheumatoid arthritis, psoriasis, exfoliative dermatitis, malaria

Excess urinary folate loss

Active liver disease, congestive heart failure

Drugs

Anticonvulsants, sulfasalazine

Mixed

Liver disease, alcoholism, intensive care

Clinical features of megaloblastic anaemia

The onset is usually insidious with gradually progressive symptoms and signs of anaemia (see Chapter 2). The patient may be mildly jaundiced (lemon yellow) (Fig. 5.6) because of the excess breakdown of haemoglobin resulting from increased ineffective erythropoiesis in the bone marrow. Glossitis (a beefy-red sore tongue) (Fig. 5.7), angular cheilosis (Fig. 5.8) and mild symptoms of malabsorption with loss of weight may be present because of the epithelial abnormality. Purpura as a result of thrombocytopenia and widespread melanin pigmentation (the cause of which is unclear) are less frequent presenting features (Table 5.6). Many asymptomatic patients are diagnosed when a blood count that has been performed for another reason reveals macrocytosis.

Vitamin B₁₂ neuropathy (subacute combined degeneration of the cord)

Severe B₁₂ deficiency can cause a progressive neuropathy affecting the peripheral sensory nerves and posterior and lateral columns (Fig. 5.9). The neuropathy is symmetrical and affects the lower limbs more than the upper limbs. The patient notices tingling in the feet, difficulty in walking and may fall over in the



Figure 5.6 Megaloblastic anaemia: pallor and mild icterus in a patient with a haemoglobin count of 70.0 g/L and a mean corpuscular volume of 132 fL.



Figure 5.7 Megaloblastic anaemia: glossitis – the tongue is beefy-red and painful.



Figure 5.8 Megaloblastic anaemia: angular cheilosis (stomatitis).

Table 5.6 Effects of vitamin B₁₂ or folate deficiency.

Megaloblastic anaemia
Macrocytosis of epithelial cell surfaces
Neuropathy (for vitamin B ₁₂ only)
Sterility
Rarely, reversible melanin skin pigmentation
Decreased osteoblast activity
Neural tube defects in the fetus are related to folate or B ₁₂ deficiency



Figure 5.9 Cross-section of the spinal cord in a patient who died with subacute combined degeneration of the cord (Weigert–Pal stain). There is demyelination of the dorsal and dorsolateral columns.

dark. Rarely, optic atrophy or severe psychiatric symptoms are present. Anaemia may be severe, mild or even absent, but the blood film and bone marrow appearances are always abnormal. The peripheral neuropathy is usually reversible with B₁₂ therapy but spinal cord recovery is incomplete, especially if the neuropathy has been present for more than a few weeks or months.

The cause of the neuropathy is likely to be related to the accumulation of *S*-adenosyl homocysteine and reduced levels of *S*-adenosyl methionine in nervous tissue resulting in defective methylation of myelin and other substrates. Folate and B₁₂ deficiency has been associated with reduced cognitive function and Alzheimer's disease but no benefit has been shown for prophylactic folic acid or B₁₂.

Neural tube defect

Folate or B₁₂ deficiency in the mother predisposes to neural tube defect (NTD) (anencephaly, spina bifida or encephalocoele) in the fetus (Fig. 5.10). The lower the maternal serum or red cell folate or serum B₁₂ levels, the higher the incidence of



Figure 5.10 A baby with neural tube defect (spina bifida).
Source: Courtesy of Professor C.J. Schorah.

NTDs. Moreover, supplementation of the diet with folic acid at the time of conception and in early pregnancy reduces the incidence of NTD by 75%. The exact mechanism is uncertain but is thought to be related to build-up of homocysteine and *S*-adenosyl homocysteine in the fetus which may impair methylation of various proteins and lipids. A common polymorphism (677C → T) in the enzyme 5,10-methylene tetrahydrofolate reductase (5,10-MTHFR), which reduces 5,10-MTHF to methyl-THF, results in higher serum homocysteine and lower serum and red cell folate levels compared with controls. The incidence of the mutation is higher in the parents and fetuses with NTD than in controls.

Other tissue abnormalities

Sterility is frequent in either sex with severe B₁₂ or folate deficiency. Macrocytosis, excess apoptosis and other morphological abnormalities of cervical, buccal, bladder and other epithelia occur. Widespread reversible melanin pigmentation may also occur. B₁₂ deficiency is associated with reduced osteoblastic activity.

Raised serum homocysteine levels and low serum or red cell folate and the polymorphism in the MTHFR enzyme (see above) have been associated with an increased incidence of cardiovascular diseases, including myocardial infarct, peripheral vascular diseases, stroke and venous thrombosis (see p. 306). Folic acid prophylaxis, however, has not reduced the incidence of the arterial diseases except possibly for stroke.

Various associations have been found between folate status and malignant diseases but meta-analysis of subjects randomized in trials to take folic acid or placebo for 2 years or more do not show any difference in cancer incidence between those taking folic acid and the controls.

Laboratory findings

The anaemia is macrocytic (MCV >98 fL and often as high as 120–140 fL in severe cases) and the macrocytes are typically oval (Fig. 5.11). If iron deficiency is also present the MCV may




Figure 5.11 Megaloblastic anaemia: peripheral blood film showing oval macrocytes.




Figure 5.12 Megaloblastic changes in the bone marrow in a patient with severe megaloblastic anaemia. **(a–c)** Erythroblasts showing fine, open stippled (primitive) appearance of the nuclear chromatin even in late cells (pale cytoplasm with some haemoglobin formation). **(d)** Abnormal giant metamyelocytes and band forms.

be normal. The reticulocyte count is low and the total white cell and platelet counts may be moderately reduced, especially in severely anaemic patients. A proportion of the neutrophils show hypersegmented nuclei (with six or more lobes). The bone marrow is usually hypercellular and the erythroblasts are large and show an open, fine, lacy primitive chromatin pattern but normal cytoplasmic haemoglobinization (Fig. 5.12). Giant and abnormally shaped metamyelocytes are characteristic.

The serum unconjugated bilirubin and lactate dehydrogenase are raised as a result of marrow cell breakdown.

Diagnosis of vitamin B₁₂ or folate deficiency

It is usual to assay serum B₁₂ and folate (Table 5.7). The serum B₁₂ is low in megaloblastic anaemia or neuropathy caused by B₁₂ deficiency. The serum and red cell folate are both low in megaloblastic anaemia caused by folate deficiency. In B₁₂ deficiency, the serum folate tends to rise but the red cell folate falls. In the absence of B₁₂ deficiency, however, the red cell folate may be a more accurate guide of tissue folate status than the

Table 5.7 Laboratory tests for vitamin B₁₂ and folate deficiency.

Test	Normal values*	Result in		
			Vitamin B ₁₂ deficiency	Folate deficiency
Serum vitamin B ₁₂	160–925 ng/L	120–680 pmol/L	Low	Normal or borderline
Serum folate	3.0–15.0 µg/L	4–30 nmol/L	Normal or raised	Low
Red cell folate	160–640 µg/L	360–1460 nmol/L	Normal or low	Low

* Normal values differ with different commercial kits.

serum folate. Measurement of serum methylmalonic acid is a test for B₁₂ deficiency and measurement of homocysteine is a test for folate or B₁₂ deficiency. These are not specific, however, and it is difficult to establish normal levels in different age groups. These tests are also not widely available.

Tests for cause of vitamin B₁₂ or folate deficiency

Useful tests are listed in Table 5.8. These are mainly concerned with assessing gastric function and testing for antibodies to gastric antigens. In all cases of pernicious anaemia

endoscopy studies should be performed at diagnosis to confirm the presence of gastric atrophy and exclude carcinoma of the stomach.

For folate deficiency, the dietary history is most important, although it is difficult to estimate folate intake accurately. Unsuspected gluten-induced enteropathy or other underlying conditions should also be considered (Table 5.5).

Treatment

Most cases only need therapy with the appropriate vitamin (Table 5.9). If large doses of folic acid (e.g. 5 mg/day) are given in B₁₂ deficiency they cause a haematological response but may aggravate the neuropathy. Folic acid should therefore not be given alone unless B₁₂ deficiency has been excluded. In severely anaemic patients who need treatment urgently it may be safer to initiate treatment with both vitamins after blood has been taken for B₁₂ and folate assay. Oral B₁₂ therapy may be used, e.g. in haemophiliacs, but it is difficult to initially replenish stores. Large doses are needed daily to get enough absorbed and compliance with long-term therapy may be a problem. Oral therapy may be more appropriate for those with mild degrees of B₁₂ deficiency, e.g. with malabsorption of food B₁₂. In the elderly, the presence of heart failure should be corrected with diuretics. Blood transfusion should be avoided if possible as it may cause circulatory overload.

Table 5.8 Tests for cause of vitamin B₁₂ or folate deficiency.

Vitamin B ₁₂	Folate
Diet history	Diet history
Serum gastrin	Tests for intestinal malabsorption
IF, parietal cell antibodies	Anti-transglutaminase and endomysial antibodies
Endoscopy	Duodenal biopsy Underlying disease
IF, intrinsic factor.	

Table 5.9 Treatment of megaloblastic anaemia.

	Vitamin B ₁₂ deficiency	Folate deficiency
Compound	Hydroxocobalamin	Folic acid
Route	Intramuscular*	Oral
Dose	1000 µg	5 mg
Initial dose	6 × 1000 µg over 2–3 weeks	Daily for 4 months
Maintenance	1000 µg every 3 months	Depends on underlying disease; life-long therapy may be needed in chronic inherited haemolytic anaemias, myelofibrosis, renal dialysis
Prophylactic	Total gastrectomy Ileal resection	Pregnancy, severe haemolytic anaemias, dialysis, prematurity

* Some authors recommend daily oral or sublingual therapy of vitamin B₁₂ deficiency (see text).




Figure 5.13 Typical haematological response to vitamin B₁₂ (hydroxocobalamin) therapy in pernicious anaemia.

Response to therapy

The patient feels better after 24–48 hours of correct vitamin therapy with increased appetite and well-being. The haemoglobin should rise by 20–30 g/L each fortnight. The white cell and platelet counts become normal in 7–10 days (Fig. 5.13) and the marrow is normoblastic in about 48 hours, although giant metamyelocytes persist for up to 12 days.

Prophylactic therapy

Vitamin B₁₂ is given 3-monthly to patients who have total gastrectomy or ileal resection. Folic acid is given in pregnancy at a recommended dosage of 400 µg/day and all women of child-bearing age are recommended to have an intake of at least 400 µg/day (by increased intake of folate-rich or folate-supplemented foods or as folic acid) to prevent a first occurrence of an NTD fetus.

Folic acid is also given to patients undergoing chronic dialysis, with severe chronic haemolytic anaemias, e.g. sickle cell anaemia, and to premature babies. Food fortification with folic acid (e.g. in flour or grain) is practised in over 70 countries (including North America since 1998) to reduce the incidence of NTD but not in the UK or the rest of Europe.

Other megaloblastic anaemias

See Table 5.1.

Abnormalities of vitamin B₁₂ or folate metabolism

These include congenital deficiencies of enzymes concerned in B₁₂ or folate metabolism or of the serum transport protein for B₁₂, TC. Nitrous oxide (N₂O) anaesthesia causes rapid inactivation of body B₁₂ by oxidizing the reduced cobalt atom of methyl B₁₂. Megaloblastic marrow changes occur with several days of N₂O administration and can cause pancytopenia. Chronic exposure (as in dentists and anaesthetists) has been associated with neurological damage resembling B₁₂ deficiency neuropathy. Antifolate drugs, particularly those which inhibit DHF reductase (e.g. methotrexate and pyrimethamine) may also cause megaloblastic change.

Other macrocytic anaemias

There are many non-megaloblastic causes of macrocytic anaemia (Table 5.10). The exact mechanisms creating large red cells in each of these conditions is not clear although increased lipid deposition on the red cell membrane or alterations of erythroblast maturation time in the marrow may be implicated. Alcohol is the most frequent cause of a raised MCV in the absence of anaemia. Reticulocytes are bigger than mature red cells, so haemolytic anaemia is an important cause of macrocytic anaemia. Antimetabolite drugs such as hydroxycarbamide (see Table 12.1) cause macrocytosis and the marrow may show megaloblastic changes. The other underlying conditions listed in Table 5.10 are usually easily diagnosed provided that they are considered and the appropriate investigations to exclude B₁₂ or folate deficiency are carried out.

Table 5.10 Causes of macrocytosis other than megaloblastic anaemia.

Alcohol
Liver disease
Myxoedema
Myelodysplastic syndromes
Antimetabolite drugs, e.g. hydroxycarbamide
Aplastic anaemia
Pregnancy
Smoking
Reticulocytosis
Myeloma and paraproteinaemia
Neonatal

Differential diagnosis of macrocytic anaemias


The clinical history and physical examination may suggest B₁₂ or folate deficiency as the cause. Diet, drugs, alcohol intake, family history, history suggestive of malabsorption, presence of autoimmune diseases or other associations with pernicious anaemia (Table 5.4), previous gastrointestinal disease or operations are all important. The presence of jaundice, glossitis or neuropathy are also important indications of megaloblastic anaemia.

The laboratory features of particular importance are the shape of macrocytes (oval in megaloblastic anaemia), the presence of hypersegmented neutrophils, of leucopenia and thrombocytopenia in megaloblastic anaemia, and the bone marrow appearance. Assay of serum B₁₂ and folate is essential. Exclusion of alcoholism (particularly if the patient is not anaemic), liver and thyroid function tests, and bone marrow examination for myelodysplasia, aplasia or myeloma are important in the investigation of macrocytosis not caused by B₁₂ or folate deficiency.

- Macrocytic anaemias show an increased size of circulating red cells (MCV >98 fL). Causes include vitamin B₁₂ (B₁₂, cobalamin) or folate deficiency, alcohol, liver disease, hypothyroidism, myelodysplasia, paraproteinaemia, cytotoxic drugs, aplastic anaemia, pregnancy and the neonatal period.
- B₁₂ or folate deficiency cause megaloblastic anaemia, in which the bone marrow erythroblasts have a typical abnormal appearance.
- Folate takes part in biochemical reactions in DNA synthesis. B₁₂ has an indirect role by its involvement in folate metabolism.
- B₁₂ deficiency may also cause a neuropathy due to damage to the spinal cord and peripheral nerves.
- B₁₂ deficiency is usually caused by B₁₂ malabsorption brought about by pernicious anaemia in which there is autoimmune gastritis, resulting in severe deficiency of intrinsic factor, a glycoprotein made in the stomach which facilitates B₁₂ absorption by the ileum.
- Other gastrointestinal diseases as well as a vegan diet may cause B₁₂ deficiency.
- Folate deficiency may be caused by a poor diet, malabsorption (e.g. gluten-induced enteropathy) or excess cell turnover (e.g. pregnancy, haemolytic anaemias, malignancy).
- Treatment of B₁₂ deficiency is usually with injections of hydroxocobalamin and of folate deficiency with oral folic (pteroylglutamic) acid.
- Rare causes of megaloblastic anaemia include inborn errors of B₁₂ or folate transport or metabolism, and defects of DNA synthesis not related to B₁₂ or folate.



Now visit www.wileyessential.com/haematology to test yourself on this chapter.


A vertical strip on the left side of the page showing a microscopic view of various red blood cells. Some are normal biconcave discs, while others are fragmented or irregular in shape, suggesting hemolysis. The background is dark blue.

CHAPTER 6

Haemolytic anaemias

Key topics

■ Normal red cell destruction	61
■ Introduction to haemolytic anaemias	62
■ Intravascular and extravascular haemolysis	63
■ Hereditary haemolytic anaemias	64
■ Acquired haemolytic anaemias	68



Normal red cell destruction

Red cell destruction usually occurs after a mean lifespan of 120 days when the cells are removed extravascularly by the macrophages of the reticuloendothelial (RE) system, especially in the marrow but also in the liver and spleen. As the cells have no nucleus, red cell metabolism gradually deteriorates as enzymes are degraded and the cells become non-viable. The breakdown of haem from haemoglobin liberates iron for recirculation via plasma transferrin mainly to marrow erythroblasts, and protoporphyrin, which is broken down to bilirubin. Bilirubin circulates to the liver where it

is conjugated to glucuronides, which are excreted into the gut via bile and converted to stercobilinogen and stercobilin (excreted in faeces) (Fig. 6.1). Stercobilinogen and stercobilin are partly reabsorbed and excreted in urine as urobilinogen and urobilin. Globin chains are broken down to amino acids which are reutilized for general protein synthesis in the body.

Haptoglobins are proteins in normal plasma which bind haemoglobin. The haemoglobin–haptoglobin complex is removed by the RE system. Intravascular haemolysis (breakdown of red cells within blood vessels) plays little or no part in normal red cell destruction.




Figure 6.1 (a) Normal red blood cell (RBC) breakdown. This takes place extravascularly in the macrophages of the reticuloendothelial system. (b) Intravascular haemolysis occurs in some pathological disorders.

Introduction to haemolytic anaemias

Haemolytic anaemias are defined as anaemias that result from an increase in the rate of red cell destruction. Because of erythropoietic hyperplasia and anatomical extension of bone marrow, red cell destruction may be increased several-fold before the patient becomes anaemic – compensated haemolytic disease. The normal adult marrow, after full expansion, is able to produce red cells at 6–8 times the normal rate provided this is ‘effective’. It leads to a marked reticulocytosis. Therefore, anaemia due to haemolysis may not be seen until the red cell lifespan is less than 30 days.

Classification

Table 6.1 is a simplified classification of the haemolytic anaemias. Hereditary haemolytic anaemias are the result of ‘intrinsic’ red cell defects, whereas acquired haemolytic anaemias are usually the result of an ‘extracorpuscular’ or ‘environmental’ change. Paroxysmal nocturnal haemoglobinuria

(PNH) is the exception because, although it is an acquired disorder, the PNH red cells have an intrinsic defect. PNH is associated with marrow hypoplasia and is therefore discussed in Chapter 22.

Clinical features

The patient may show pallor of the mucous membranes, mild fluctuating jaundice and splenomegaly. There is no bilirubin in urine but this may turn dark on standing because of excess urobilinogen. Pigment (bilirubin) gallstones may complicate the condition (Fig. 6.2) and some patients (particularly with sickle cell disease) develop ulcers around the ankle (see Fig. 7.19). Aplastic crises may occur, usually precipitated by infection with parvovirus which ‘switches off’ erythropoiesis, and are characterized by a sudden increase in anaemia and drop in reticulocyte count (see Fig. 22.7).

Rarely, folate deficiency may cause an aplastic crisis in which the bone marrow is megaloblastic.

Table 6.1 Classification of haemolytic anaemias.	
Hereditary	Acquired
Membrane Hereditary spherocytosis, hereditary elliptocytosis	Immune
Metabolism G6PD deficiency, pyruvate kinase deficiency	<i>Autoimmune</i> Warm antibody type (see Table 6.5) Cold antibody type
Haemoglobin Genetic abnormalities (Hb S, Hb C, unstable); see Chapter 7	<i>Alloimmune</i> Haemolytic transfusion reactions Haemolytic disease of the newborn Allografts, especially marrow transplantation <i>Drug associated</i> Red cell fragmentation syndromes See Table 6.6 March haemoglobinuria Infections Malaria, clostridia Chemical and physical agents Especially drugs, industrial/domestic substances, burns Secondary Liver and renal disease Paroxysmal nocturnal haemoglobinuria (see Chapter 22)
G6PD, glucose-6-phosphate dehydrogenase; Hb, haemoglobin.	




Figure 6.2 Ultrasound of multiple small pigment gallstones typical of those associated with hereditary spherocytosis. Source: Courtesy of Dr P. Wylie.

Laboratory findings

The laboratory findings are conveniently divided into three groups.

- 1 Features of increased red cell breakdown:
 - (a) serum bilirubin raised, unconjugated and bound to albumin;
 - (b) urine urobilinogen increased;
 - (c) serum haptoglobins absent because the haptoglobins become saturated with haemoglobin and the complex is removed by RE cells.
- 2 Features of increased red cell production:
 - (a) reticulocytosis;
 - (b) bone marrow erythroid hyperplasia; the normal marrow myeloid:erythroid ratio of 2:1 to 12:1 is reduced to 1:1 or reversed.
- 3 Damaged red cells:
 - (a) morphology (e.g. microspherocytes, elliptocytes, fragments);
 - (b) osmotic fragility;
 - (c) specific enzyme, protein or DNA tests.

Intravascular and extravascular haemolysis

There are two mechanisms whereby red cells are destroyed in haemolytic anaemia. There may be excessive removal of red cells by cells of the RE system (**extravascular haemolysis**) or they may be broken down directly in the circulation (**intravascular haemolysis**) (Fig. 6.1; Table 6.2). Whichever mechanism dominates will depend on the pathology involved. In intravascular

Table 6.2 Causes of intravascular haemolysis.

Mismatched blood transfusion (usually ABO)
G6PD deficiency with oxidant stress
Red cell fragmentation syndromes
Some severe autoimmune haemolytic anaemias
Some drug- and infection-induced haemolytic anaemias
Paroxysmal nocturnal haemoglobinuria
March haemoglobinuria
Unstable haemoglobin
G6PD, glucose-6-phosphate dehydrogenase

haemolysis, free haemoglobin is released which rapidly saturates plasma haptoglobins and the excess free haemoglobin is filtered by the glomerulus. If the rate of haemolysis saturates the renal tubular reabsorptive capacity, free haemoglobin enters urine (Fig. 6.3). Iron released from haemoglobin in the renal tubules is seen as haemosiderin in a urinary deposit. Methaemalbumin is also formed from the process of intravascular haemolysis.

The main laboratory features of intravascular haemolysis therefore are (Fig. 6.3):

- 1 Haemoglobinaemia and haemoglobinuria.
- 2 Haemosiderinuria.
- 3 Methaemalbuminaemia (detected spectrophotometrically).




Figure 6.3 (a) Progressive urine samples in an acute episode of intravascular haemolysis showing haemoglobinuria of decreasing severity. (b) Prussian blue-positive deposits of haemosiderin in a urine spun deposit (Perls' stain).

Hereditary haemolytic anaemias

Membrane defects

Hereditary spherocytosis

Hereditary spherocytosis (HS) is the most common hereditary haemolytic anaemia in Northern Europeans.

Pathogenesis

HS is usually caused by defects in the proteins involved in the vertical interactions between the membrane skeleton and the lipid bilayer of the red cell (Table 6.3; see Fig. 2.12). The marrow produces red cells of normal biconcave shape but these lose membrane and become increasingly spherical (loss of surface area relative to volume) as they circulate through the spleen and the rest of the RE system. The loss of membrane may be caused by the release of parts of the lipid bilayer that are not supported by the skeleton. Ultimately, the spherocytes are unable to pass through the microcirculation, especially in the spleen, and die prematurely.

Clinical features

The inheritance is autosomal dominant with variable expression; rarely it may be autosomal recessive. The anaemia can

present at any age from infancy to old age. Jaundice is typically fluctuating and is particularly marked if the haemolytic anaemia is associated with Gilbert's disease (a defect of hepatic conjugation of bilirubin); splenomegaly occurs in most patients. Pigment gallstones are frequent (Fig. 6.2); aplastic crises, usually precipitated by parvovirus infection, may cause a sudden increase in severity of anaemia (see Fig. 22.7).

Table 6.3 Molecular basis of hereditary spherocytosis and elliptocytosis.

Hereditary spherocytosis

Ankyrin deficiency or abnormalities
 α - or β -spectrin deficiency or abnormalities
 Band 3 abnormalities
 Pallidin (protein 4.2) abnormalities

Hereditary elliptocytosis

α - or β -spectrin mutants leading to defective spectrin dimer formation
 α - or β -spectrin mutants leading to defective spectrin–ankyrin associations
 Protein 4.1 deficiency or abnormality

South-East Asian ovalocytosis band 3 deletion




Figure 6.4 (a) Blood film in hereditary spherocytosis. The spherocytes are deeply staining and of small diameter. Larger polychromatic cells are reticulocytes (confirmed by supravital staining). (b) Blood film in hereditary elliptocytosis.

Haematological findings

Anaemia is usual but not invariable; its severity tends to be similar in members of the same family. Reticulocytes are usually 5–20%. The blood film shows microspherocytes (Fig. 6.4a) which are densely staining with smaller diameters than normal red cells.

Investigation and treatment

A rapid flow analysis of eosin–maleimide (EMA) bound to erythrocytes is used as a test for HS and membrane protein deficiency (Fig. 6.5). Identification of the exact molecular defect is not needed for management but electrophoresis of

membrane proteins is carried out in difficult cases. The EMA test has replaced the osmotic fragility test which showed the red cells to be excessively fragile in dilute saline solutions. The direct antiglobulin (Coombs') test is normal, excluding an autoimmune cause of spherocytosis and haemolysis.

The principal form of treatment is splenectomy, preferably laparoscopic, although this should not be performed unless clinically indicated by symptomatic anaemia or gallstones, leg ulcers or growth retardation, because of the risk of post-splenectomy sepsis, particularly in early childhood (p. 121). Cholecystectomy should be performed with splenectomy if symptomatic gallstones are present. Splenectomy should always produce a rise in the haemoglobin level to normal, even though microspherocytes formed in the rest of the RE system will remain. Folic acid is given in severe cases to prevent folate deficiency.




Figure 6.5 Eosin-5-maleimide staining in hereditary spherocytosis (HS) showing reduced mean channel fluorescence due to membrane band 3 protein deficiency. Source: Courtesy of Mr G. Ellis.

Hereditary elliptocytosis

This has similar clinical and laboratory features to HS except for the appearance of the blood film (Fig. 6.4b), but is usually a clinically milder disorder. It is predominantly discovered by chance on a blood film and there may be no evidence of haemolysis. Occasional patients require splenectomy. The basic defect is a failure of spectrin heterodimers to self-associate into heterotetramers. A number of genetic mutations affecting horizontal interactions have been detected (Table 6.3). Patients with homozygous or doubly heterozygous elliptocytosis present with a severe haemolytic anaemia termed hereditary pyropoikilocytosis.

South-East Asian ovalocytosis

This is common in Melanesia, Malaysia, Indonesia and the Philippines and is caused by a nine amino acid deletion at the junction of the cytoplasmic and transmembrane domains of the band 3 protein. The cells are rigid and resist invasion by malarial parasites. Most cases are not anaemic and are asymptomatic.

Defective red cell metabolism

Glucose-6-phosphate dehydrogenase deficiency

Glucose-6-phosphate dehydrogenase (G6PD) functions to reduce nicotinamide adenine dinucleotide phosphate (NADP). It is the only source of NADPH which is needed for the production of reduced glutathione; a deficiency renders the red cell susceptible to oxidant stress (Fig. 6.6).

Epidemiology

There is a wide variety of normal genetic variants of the enzyme G6PD, the most common being type B (Western) and type A in Africans. In addition, more than 400 variants caused by point mutations or deletions of the enzyme G6PD have been characterized which show less activity than normal, and worldwide over 400 million people are G6PD deficient (Fig. 6.7).

The inheritance is sex-linked, affecting males, and carried by females who show approximately half the normal red cell G6PD values. The female heterozygotes have an advantage of resistance to *Falciparum* malaria. The degree of deficiency varies with ethnic group, often being mild (10–60% of normal activity) in black African people, more severe in Middle-Eastern and South-East Asian people, and most severe in Mediterranean people (<10% of normal activity). Severe deficiency occurs occasionally in white people.




Figure 6.6 Haemoglobin and red blood cell (RBC) membranes are usually protected from oxidant stress by reduced glutathione (GSH). In G6PD deficiency, NADPH and GSH synthesis is impaired. F6P, fructose-6-phosphate; G6P, glucose-6-phosphate; G6PD, glucose-6-phosphate dehydrogenase; 6-PG, 6-phosphogluconate; GSSG, glutathione (oxidized form); NADP, NADPH, nicotinamide adenine dinucleotide phosphate.




Figure 6.7 Global distribution of *G6PD* gene variants causing G6PD deficiency. Shaded areas indicate the prevalence of G6PD deficiency. Source: Adapted from Luzzatto & Notaro (2001) *Science* 293: 442.

Table 6.4 Agents that may cause haemolytic anaemia in glucose-6-phosphate dehydrogenase (G6PD) deficiency.

<i>Infections and other acute illnesses</i> (e.g. diabetic ketoacidosis)
<i>Drugs</i>
Antimalarials (e.g. primaquine, pamaquine, chloroquine, Fansidar, Maloprim)
Sulphonamides and sulphones (e.g. co-trimoxazole, sulfanilamide, dapsone, sulfasalazine)
Other antibacterial agents (e.g. nitrofurans, chloramphenicol)
Analgesics (e.g. aspirin), moderate doses are safe
Antihelminths (e.g. β -naphthol, stibophen)
Miscellaneous (e.g. vitamin K analogues, naphthalene (mothballs), probenecid)
Fava beans (possibly other vegetables)
N.B. Many common drugs have been reported to precipitate haemolysis in G6PD deficiency in some patients (e.g. aspirin, quinine and penicillin) but not at conventional dosage.

Clinical features

G6PD deficiency is usually asymptomatic with a normal blood count between attacks of haemolysis. Although G6PD is present in all cells, the main clinical syndromes that occur are as follows.

- 1 Acute haemolytic anaemia in response to oxidant stress, e.g. drugs, fava beans or infections (Table 6.4). Fava beans (*Vicia fava*) contain an oxidant chemical, divicine. The acute haemolytic anaemia is caused by rapidly developing intravascular haemolysis with haemoglobinuria (Fig. 6.3a). The anaemia may be self-limiting as new young red cells are made with near normal enzyme levels.
- 2 Neonatal jaundice.
- 3 Rarely, a congenital non-spherocytic haemolytic anaemia.

These syndromes may result from different types of severe enzyme deficiency.

Diagnosis

Between crises the blood count is normal. The enzyme deficiency is detected by one of a number of screening tests or by direct enzyme assay on red cells. During a crisis the blood film may show contracted and fragmented cells, 'bite' cells and 'blister' cells (Fig. 6.8), which have had Heinz bodies removed by the spleen. Heinz bodies (oxidized, denatured haemoglobin) may be seen in the reticulocyte preparation, particularly if the spleen is absent. There are also features of intravascular haemolysis. Because of the higher enzyme level in young red cells, red cell enzyme assay may give a 'false' normal level in the phase of acute haemolysis with a reticulocyte response. Subsequent assay after the acute phase reveals the low G6PD level when the red cell population is of normal age distribution.




Figure 6.8 Blood film in G6PD deficiency with acute haemolysis after an oxidant stress. Some of the cells show loss of cytoplasm with separation of remaining haemoglobin from the cell membrane ('blister' cells). There are also numerous contracted and deeply staining cells. Supravital staining (as for reticulocytes) showed the presence of Heinz bodies (see Fig. 2.17).

Treatment

The offending drug is stopped, any underlying infection is treated, a high urine output is maintained and blood transfusion undertaken where necessary for severe anaemia. G6PD-deficient babies are prone to neonatal jaundice and in severe cases phototherapy and exchange transfusion may be needed. The jaundice is usually not caused by excess haemolysis but by deficiency of G6PD affecting neonatal liver function.

Glutathione deficiency and other syndromes

Other defects in the pentose phosphate pathway leading to similar syndromes to G6PD deficiency have been described – particularly glutathione deficiency.

Glycolytic (Embden–Meyerhof) pathway defects

These are all uncommon and lead to a congenital non-spherocytic haemolytic anaemia. In some there are defects of other systems (e.g. a myopathy). The most frequently encountered is pyruvate kinase (PK) deficiency (see Fig. 2.11).

Pyruvate kinase deficiency

This is inherited as an autosomal recessive, the affected patients being homozygous or doubly heterozygous. Over 100 different mutations have been described. The red cells become rigid as a result of reduced adenosine triphosphate (ATP) formation. The severity of the anaemia varies widely (haemoglobin 40–100 g/L) and causes relatively mild symptoms because of a shift to the right in the oxygen (O_2) dissociation curve caused by a rise in intracellular 2,3-diphosphoglycerate (2,3-DPG). Clinically, jaundice is usual and gallstones are frequent. Frontal bossing may be present. The blood film shows poikilocytosis and distorted 'prickle' cells, particularly post-splenectomy.

Direct enzyme assay is needed to make the diagnosis. Splenectomy may alleviate the anaemia but does not cure it and is indicated in those patients who need frequent transfusions. Iron loading is frequent due to low serum hepcidin levels consequent on increased and ineffective erythropoiesis as well as blood transfusions.

Hereditary disorders of haemoglobin synthesis

Several of these cause clinical haemolysis. They are discussed in Chapter 7.

Acquired haemolytic anaemias

Immune haemolytic anaemias

Autoimmune haemolytic anaemias

Autoimmune haemolytic anaemias (AIHAs) are caused by antibody production by the body against its own red cells. They are characterized by a positive direct antiglobulin test (DAT), also known as the Coombs' test (see Fig. 30.5), and divided into 'warm' and 'cold' types (Table 6.5) according to whether the antibody reacts more strongly with red cells at 37°C or 4°C.

Warm autoimmune haemolytic anaemias

The red cells are coated with immunoglobulin (Ig), usually immunoglobulin G (IgG) alone or with complement, and are therefore taken up by RE macrophages which have receptors for the Ig Fc fragment. Part of the coated membrane is lost so the cell becomes progressively more spherical to maintain the same volume and is ultimately prematurely destroyed, predominantly in the spleen. When the cells are coated with IgG and complement (C3d, the degraded fragment of C3) or complement alone, red cell destruction occurs more generally in the RE system.

Clinical features

The disease may occur at any age, in either sex, and presents as a haemolytic anaemia of varying severity. The spleen is often enlarged. The disease tends to remit and relapse. It may occur alone or in association with other diseases (Table 6.5). When associated with idiopathic thrombocytopenic purpura (ITP), a similar condition affecting platelets (p. 282), it is called Evans' syndrome. When secondary to systemic lupus erythematosus, the cells typically are coated with immunoglobulin and complement.

Laboratory findings

The haematological and biochemical findings are typical of an extravascular haemolytic anaemia with spherocytosis prominent in the peripheral blood (Fig. 6.9a). The DAT is positive as a result of IgG, IgG and complement or IgA on the cells and, in some cases, the autoantibody shows specificity within the Rh system. The antibodies both on the cell surface and free in serum are best detected at 37°C.

Treatment

- 1 Remove the underlying cause (e.g. methyl dopa).
- 2 Corticosteroids. Prednisolone is the usual first-line treatment; 1 mg/kg/day is a typical starting dose in adults and should then be tapered down. Those with predominantly IgG on red cells do best, whereas those with complement often respond poorly, both to corticosteroids and splenectomy.
- 3 Monoclonal antibody. Anti-CD20 (rituximab) has produced prolonged remissions in a proportion of cases and is being used with steroids as first-line therapy.
- 4 Splenectomy may be of value in those who fail to respond well or fail to maintain a satisfactory haemoglobin level on an acceptably small steroid dosage or rituximab maintenance.

Table 6.5 Immune haemolytic anaemias: classification.

Warm type	Cold type
Autoimmune <i>Idiopathic</i> <i>Secondary</i> SLE, other 'autoimmune' diseases CLL, lymphomas Drugs (e.g. methyl dopa)	<i>Idiopathic</i> <i>Secondary</i> Infections – <i>Mycoplasma pneumonia</i> , infectious mononucleosis Lymphoma Paroxysmal cold haemoglobinuria (rare, sometimes associated with infections, e.g. syphilis)
Alloimmune <i>Induced by red cell antigens</i> Haemolytic transfusion reactions Haemolytic disease of the newborn post stem cell grafts <i>Drug induced</i> Drug–red cell membrane complex Immune complex	

CLL, chronic lymphocytic leukaemia; SLE, systemic lupus erythematosus.




Figure 6.9 (a) Blood film in warm autoimmune haemolytic anaemia. Numerous microspherocytes are present and larger polychromatic cells (reticulocytes). (b) Blood film in cold autoimmune haemolytic anaemia. Marked red cell agglutination is present in films made at room temperature. The background is caused by the raised plasma protein concentration.

- 5 Immunosuppression may be tried after the above measures have failed or even before splenectomy. Anti-CD52 (Alem-tuzumab), azathioprine, cyclophosphamide, chlorambucil, ciclosporin and mycophenolate mofetil have been tried.
- 6 It may be necessary to treat the underlying disease, e.g. chronic lymphocytic leukaemia or lymphoma.
- 7 Folic acid is given to severe cases.
- 8 Blood transfusion may be needed if anaemia is severe and causing symptoms. The blood should be the least incompatible and, if the specificity of the autoantibody is known, donor blood is chosen that lacks the relevant antigen(s). The patients also readily make alloantibodies against donor red cells.
- 9 High-dose immunoglobulin has been used but with less success than in ITP (p. 283).

Cold autoimmune haemolytic anaemias

In these syndromes the IgM autoantibody attaches to red cells mainly in the peripheral circulation where the blood temperature is cooled (Table 6.5). The autoantibody may be monoclonal, as in primary cold haemagglutinin syndrome or associated with lymphoproliferative disorders, or may be a transient polyclonal response following infections such as infectious mononucleosis or *Mycoplasma pneumoniae*. The IgM antibodies which bind to red cells optimally at 4°C, are highly efficient at fixing complement such that intravascular and extravascular haemolysis can occur. Only complement factors can be detected on red cells in laboratory tests as the IgM antibody is eluted off as cells flow through warmer parts of the circulation. Interestingly, in nearly all these cold AIHA syndromes

the antibody is directed against the 'I' antigen on the red cell surface. In infectious mononucleosis it is anti-i.

Primary cold agglutinin disease

The patient has a chronic haemolytic anaemia aggravated by the cold and often associated with intravascular haemolysis. Mild jaundice and splenomegaly may be present. The patient may develop acrocyanosis (purplish skin discoloration) at the tip of the nose, ears, fingers and toes caused by the agglutination of red cells in small vessels.

Laboratory findings are similar to those of warm AIHA except that spherocytosis is less marked, red cells agglutinate in the cold (Fig. 6.9b) and the DAT reveals complement (C3d) only on the red cell surface. In most patients, nodules of a monoclonal population of B lymphocytes are present in the bone marrow. Their morphology and immunophenotype differs from that in lymphoplasmacytic lymphoma, also associated with an IgM paraprotein; the MYD88 mutation characteristic of that disease is absent. Cold agglutinin disease is indolent but may transform to an aggressive lymphoma.

Treatment consists of keeping the patient warm. Alkylating agents such as chlorambucil or purine nucleotide analogues (e.g. fludarabine) may be used, and anti-CD20 (rituximab) can also be effective. Splenectomy is not indicated unless massive splenomegaly is present. Corticosteroids are of no value. Underlying lymphoma should be excluded in 'idiopathic' cases.

Paroxysmal cold haemoglobinuria is a rare syndrome of acute intravascular haemolysis after exposure to the cold. It is caused by the Donath–Landsteiner antibody, an IgG antibody with specificity for the P blood group antigens, which binds to red

cells in the cold but causes lysis with complement in warm conditions. Viral infections are predisposing causes and the condition is usually self-limiting.

Alloimmune haemolytic anaemias

In these anaemias, antibody produced by one individual reacts with red cells of another. Two important situations are transfusion of ABO-incompatible blood and Rh disease of the newborn which are considered in Chapters 30 and 31. The increased use of allogeneic transplantation for renal, hepatic, cardiac and bone marrow diseases has led to the recognition of alloimmune haemolytic anaemia, resulting from the destruction of red cells of the recipient by antibodies produced by donor lymphocytes before the blood group of the recipient becomes that of the donor.

Drug-induced immune haemolytic anaemias

Drugs may cause immune haemolytic anaemias via three mechanisms (Fig. 6.10):

- 1 Antibody directed against a drug–red cell membrane complex (e.g. penicillin, ampicillin); this only occurs with massive doses of the antibiotic.
- 2 Deposition of complement via a drug–protein (antigen)–antibody complex onto the red cell surface (e.g. quinidine, rifampicin); or
- 3 A true autoimmune haemolytic anaemia in which the role of the drug is unclear (e.g. methyl dopa).

In each case, the haemolytic anaemia gradually disappears when the drug is discontinued.

Red cell fragmentation syndromes

These arise through physical damage to red cells either on abnormal surfaces (e.g. artificial heart valves or arterial grafts), arteriovenous malformations or as a microangiopathic haemolytic anaemia. This is caused by red cells passing through abnormal small vessels. The latter may be caused by deposition of fibrin strands, often associated with disseminated intravascular

Table 6.6 Red cell fragmentation syndromes.

Cardiac haemolysis	Prosthetic heart valves Patches, grafts Perivalvular leaks
Arteriovenous malformations	
Microangiopathic	TTP-HUS Disseminated intravascular coagulation Malignant disease Vasculitis (e.g. polyarteritis nodosa) Malignant hypertension Pre-eclampsia/HELLP Renal vascular disorders/HELLP syndrome Ciclosporin Homograft rejection

HELLP, haemolysis with elevated liver function tests and low platelets; HUS, haemolytic uraemic syndrome; TTP, thrombotic thrombocytopenic purpura.

coagulation (DIC), or platelet adherence as in thrombotic thrombocytopenic purpura (TTP) (p. 285) or vasculitis (e.g. polyarteritis nodosa) (Table 6.6). The peripheral blood contains many deeply staining red cell fragments (Fig. 6.11). When DIC underlies the haemolysis, clotting abnormalities (p. 297) and a low platelet count are also present. TTP is discussed in detail on p. 285.

March haemoglobinuria

This is caused by damage to red cells between the small bones of the feet, usually during prolonged marching or running. The blood film does not show fragments.




Figure 6.10 Three different mechanisms of drug-induced immune haemolytic anaemia. In each case the coated (opsonized) cells are destroyed in the reticuloendothelial system.




Figure 6.11 Blood film in microangiopathic haemolytic anaemia (in this patient Gram-negative septicaemia). Numerous contracted and deeply staining cells and cell fragments are present.

Infections

Infections can cause haemolysis in a variety of ways. They may precipitate an acute haemolytic crisis in G6PD deficiency or cause microangiopathic haemolytic anaemia (e.g. with meningococcal or pneumococcal septicaemia). Malaria causes haemolysis by extravascular destruction of parasitized red cells as well as by direct intravascular lysis. Blackwater fever is an acute intravascular haemolysis accompanied by acute renal failure caused by *Falciparum* malaria. *Clostridium perfringens* septicaemia can cause intravascular haemolysis with marked microspherocytosis.

Chemical and physical agents

Certain drugs (e.g. dapsone and sulfasalazine) in high doses cause oxidative intravascular haemolysis with Heinz body formation in normal subjects. In Wilson's disease an acute haemolytic anaemia can occur as a result of high levels of copper in the blood. Chemical poisoning (e.g. with lead, chlorate or arsine) can cause severe haemolysis. Severe burns damage red cells causing acanthocytosis or spherocytosis.


Secondary haemolytic anaemias

In many systemic disorders red cell survival is shortened. This may contribute to anaemia (see Chapter 29).

- Haemolytic anaemia is caused by shortening of the red cell life. The red cells may break down in the reticuloendothelial system (extravascular) or in the circulation (intravascular).
- Haemolytic anaemia may be caused by inherited red cell defects, which are usually intrinsic to the red cell, or to acquired causes, which are usually caused by an abnormality of the red cell environment.
- Features of extravascular haemolysis include jaundice, gallstones and splenomegaly with raised reticulocytes, unconjugated bilirubin and absent haptoglobins. In intravascular haemolysis (e.g. caused by ABO mismatched blood transfusion), there is haemoglobinaemia, methaemalbuminaemia, haemoglobinuria and haemosiderinuria.
- Genetic defects include those of the red cell membrane (e.g. hereditary spherocytosis), enzyme deficiencies (e.g. glucose-6-phosphate dehydrogenase or pyruvate kinase deficiency) or haemoglobin defects (e.g. sickle cell anaemia, see Chapter 7).
- Acquired causes of haemolytic anaemia include warm or cold, auto- or allo-antibodies to red cells, red cell fragmentation syndromes, infections, toxins and paroxysmal nocturnal haemoglobinuria (see Chapter 22).



Now visit www.wileyessential.com/haematology to test yourself on this chapter.




CHAPTER 7

Genetic disorders of haemoglobin

Key topics

■ Haemoglobin synthesis	73
■ Haemoglobin abnormalities	74
■ Thalassaemias	75
■ α -Thalassaemias	76
■ β -Thalassaemias	76
■ Thalassaemia intermedia	79
■ Sickle cell anaemia	81
■ Prenatal diagnosis of genetic haemoglobin disorders	85



This chapter deals with inherited diseases caused by reduced or abnormal synthesis of globin. Mutations in the globin genes are the most prevalent monogenic disorders worldwide and affect approximately 7% of the world's population. The synthesis of normal haemoglobin in the fetus and adult is described first.

Haemoglobin synthesis

Normal adult blood contains three types of haemoglobin (see Table 2.3). The major component is haemoglobin A with the molecular structure $\alpha_2\beta_2$. The minor haemoglobins contain γ (fetal Hb or Hb F) or δ (Hb A₂) globin chains instead of β chains. In the embryo and fetus, Gower 1, Portland, Gower 2 and fetal Hb dominate at different stages (Fig. 7.1). The genes for the globin chains occur in two clusters: ϵ , γ , δ and β on chromosome 11 and ζ and α on chromosome 16. Two types of γ chain occur, G_γ and A_γ , which differ by a glycine or alanine amino acid at position 136 in the polypeptide chain. The α -chain gene is duplicated and both α_1 and α_2 on each chromosome are active (Fig. 7.1).

Molecular aspects

All the globin genes have three exons (coding regions) and two introns (non-coding regions whose DNA is not represented in the finished protein). The initial RNA is transcribed from both introns and exons, and from this transcript the RNA derived from introns is removed by splicing (Fig. 7.2). The introns always begin with a G-T dinucleotide and end with an A-G dinucleotide. The splicing machinery recognizes these sequences as well as neighbouring conserved sequences. The newly formed mRNA is polyadenylated at the 3' end (Fig. 7.2). This stabilizes it. Thalassaemia may arise from mutations or deletions of any of these sequences.

A number of other conserved sequences are important in globin synthesis and mutations at these sites may also give rise to thalassaemia. These sequences influence gene transcription, ensure its fidelity, specify sites for the initiation and termination of translation, and ensure the stability of newly synthesized mRNA. Promoters are found 5' of the gene (see Fig. 1.8). Enhancers occur either 5' or 3' to the gene and are important in the tissue-specific regulation of globin gene expression and in regulation of the




Figure 7.1 (a) The globin gene clusters on chromosomes 16 and 11. In embryonic, fetal and adult life different genes are activated or suppressed. The different globin chains are synthesized independently and then combine with each other to produce the different haemoglobins. The γ gene may have two sequences, which code for either a glutamic acid or alanine residue at position 136 (G_γ or A_γ , respectively). LCR, locus control region, HS-40, see text. **(b)** Synthesis of individual globin chains in prenatal and postnatal life.




Figure 7.2 The expression of a human globin gene from transcription, excision of introns, splicing of exons and translation to ribosomes. The primary transcript is 'capped' at the 5' end and a poly A tail is then added.

synthesis of the various globin chains during fetal and postnatal life. The locus control region (LCR) is a genetic regulatory element, situated upstream of the β -globin cluster, that controls genetic activity by opening up the chromatin to allow transcription factors to bind. A similar region applies to α -globin synthesis.

Switch from fetal to adult haemoglobin

The globin genes are arranged on chromosomes 11 and 16 in the order in which they are expressed (Fig. 7.1). Certain embryonic haemoglobins are usually only expressed in yolk sac erythroblasts. The β -globin gene is expressed at a low level in early fetal life, but the main switch to adult haemoglobin occurs 3–6 months after birth when synthesis of the γ chain is replaced by β chains. BCL11A is a major transcriptional regulator of the switch. Other nuclear transcriptional factors are involved. The methylation state of the gene (expressed genes are hypomethylated, non-expressed hypermethylated), the state of the chromosome packaging and enhancer sequences all play a part in determining whether a particular gene will be transcribed.

Haemoglobin abnormalities

These result from the following:

- 1 Synthesis of an abnormal haemoglobin.
- 2 Reduced rate of synthesis of normal α - or β -globin chains (the α - and β -thalassaemias, Fig. 7.3).




Figure 7.3 The ratio of α : β globin chain synthesis (y axis) depending on the number of functioning α and β chain genes (x axis). Source: Mehta A.B. & Hoffbrand A.V. (2014) *Haematology at a Glance*, 4th edn. Reproduced with permission of John Wiley & Sons.

Table 7.1 The clinical syndromes produced by haemoglobin abnormalities.

Syndrome	Abnormality
Haemolysis	Crystalline haemoglobins (Hb S, C, D, E, etc.)
	Unstable haemoglobin
Thalassaemia	α or β resulting from reduced globin chain synthesis
Familial polycythaemia	Altered oxygen affinity
Methaemoglobinaemia	Failure of reduction (Hb Ms)

Table 7.1 shows some of the abnormal haemoglobins that arise from synthesis of an α or β chain with an amino acid substitution. Clinically most important is sickle cell anaemia. Haemoglobin (Hb) C, D and E are also common and, like Hb S, are substitutions in the β chain. Unstable haemoglobins are rare and cause a chronic haemolytic anaemia of varying severity with intravascular haemolysis (see Table 6.2). Abnormal haemoglobins may also cause (familial) polycythaemia (Chapter 15) or congenital methaemoglobinaemia (Chapter 2).

The genetic defects of haemoglobin are the most common genetic disorders worldwide. They occur in tropical and subtropical areas (Fig. 7.4) and most appear to have been selected because the carrier state affords some protection against malaria.

Thalassaemias

These are a heterogeneous group of genetic disorders that result from a reduced rate of synthesis of α or β chains (Fig. 7.3; Table 7.2). β -Thalassaemia is more common in the Mediterranean region while α -thalassaemia is more common in the Far East (Fig. 7.4).




Figure 7.4 The geographical distribution of the thalassaemias and the more common, inherited, structural haemoglobin abnormalities.

Table 7.2 Classification of thalassaemia*.

Clinical

Hydrops fetalis

Four gene deletion α -thalassaemia

Thalassaemia major

Transfusion dependent, homozygous

β^0 -thalassaemia or other combinations of β -thalassaemia trait

Thalassaemia intermedia (non-transfusion dependent thalassaemia)

See Table 7.3

Genetic

Thalassaemia minor

β^0 -Thalassaemia trait

β^+ -Thalassaemia trait

α^0 -Thalassaemia trait

α^+ -Thalassaemia trait

Type	Haplotype	Heterozygous thalassaemia trait (minor)*	Homozygous
α-Thalassaemias[†]			
α^0	— —/	MCV, MCH low	Hydrops fetalis
α^+	— α /	MCV, MCH minimally reduced	As heterozygous α^0 -thalassaemia [†] Compound heterozygote $\alpha^0\alpha^+$ (— —/ α) is haemoglobin H disease
β-Thalassaemias			
β^0		MCV, MCH low (Hb A ₂ >3.5%)	Thalassaemia major (Hb F 98%, Hb A ₂ 2%)
β^+		MCV, MCH low (Hb A ₂ >3.5%)	Thalassaemia major or intermedia (Hb F 70–80%, Hb A 10–20%, Hb A ₂ variable)

α^0 = 2 α genes deleted or mutated, α^+ = one α gene deleted or mutated

*See text for the less common diseases: $\delta\beta$ -thalassaemia, haemoglobin Lepore and dominant β -thalassaemia trait

Clinically the main syndromes are **transfusion dependent thalassaemia major**, non-transfusion dependent thalassaemia (**thalassaemia intermedia**) with a moderate degree of anaemia due to a variety of genetic defects (see Table 7.3) and **thalassaemia minor**, usually due to a carrier state for α - or β -thalassaemia.

α -Thalassaemia syndromes

These are caused by α -globin gene deletions or less frequently mutations (Table 7.2). The clinical severity is related to the number of the four α -globin genes missing or inactive. Loss of all four genes completely suppresses α -chain synthesis (Fig. 7.5) and because the α chain is essential in fetal as well as in adult haemoglobin this is incompatible with life and leads to death *in utero* (**hydrops fetalis**; Fig. 7.6). Three α gene deletion leads to a moderately severe (haemoglobin 70–110 g/L) microcytic, hypochromic anaemia (Fig. 7.7) with splenomegaly. This is known as **Hb H disease** because haemoglobin H (β_4) can be detected in red cells of these patients by electrophoresis or in reticulocyte preparations (Fig. 7.7). In fetal life, Hb Barts (γ_4) occurs.

The **α -thalassaemia** traits are caused by loss of one or two genes and are usually not associated with anaemia. The mean corpuscular volume (MCV) and mean corpuscular haemoglobin (MCH) are low and the red cell count is over $5.5 \times 10^{12}/L$. Haemoglobin electrophoresis is normal and DNA analysis is needed to be certain of the diagnosis. Uncommon

non-deletional forms of α -thalassaemia are caused by point mutations producing dysfunction of the genes or rarely by mutations affecting termination of translation which give rise to an elongated but unstable chain (e.g. Hb Constant Spring). Two rare forms of α -thalassaemia are associated with mental retardation. They are caused by mutation in a gene on chromosome 16 (ATR-16) or on chromosome X (ATR-X) which control the transcription of the α globin and other genes.

β -Thalassaemia syndromes

β -Thalassaemia major

This condition occurs on average in one in four offspring if both parents are carriers of the β -thalassaemia trait. Either no β chain (β^0) or small amounts (β^+) are synthesized (Fig. 7.3). Excess α chains precipitate in erythroblasts and in mature red cells causing severe ineffective erythropoiesis and haemolysis that are typical of this disease. The greater the α -chain excess, the more severe the anaemia. Production of γ chains helps to 'mop up' excess α chains and to ameliorate the condition. Over 400 different genetic defects have been detected (Fig. 7.8).

Unlike α -thalassaemia, the majority of genetic lesions are point mutations rather than gene deletions. These mutations may be within the gene complex itself or in promoter or enhancer regions. Certain mutations are particularly frequent in some communities and this may simplify antenatal diagnosis aimed at detecting the mutations in fetal DNA. Thalassaemia




Figure 7.5 The genetics of α -thalassaemia. Each α gene may be deleted or (less frequently) dysfunctional. The orange boxes represent normal genes, and the blue boxes represent gene deletions or dysfunctional genes.



Figure 7.6 α -Thalassaemia: hydrops fetalis, the result of deletion of all four α -globin genes (homozygous α^0 -thalassaemia). The main haemoglobin present is Hb Barts (γ_4). The condition is incompatible with life beyond the fetal stage. Source: Courtesy of Professor D. Todd.




Figure 7.7 (a) α -Thalassaemia: haemoglobin H disease (three α -globin gene deletion). The blood film shows marked hypochromic, microcytic cells with target cells and poikilocytosis. (b) α -Thalassaemia: haemoglobin H disease. Supravital staining with brilliant cresyl blue reveals multiple fine, deeply stained deposits ('golf ball' cells) caused by precipitation of aggregates of β -globin chains. Hb H can also be detected as a fast-moving band on haemoglobin electrophoresis (see Fig. 7.12).

major is often a result of inheritance of two different mutations, each affecting β -globin synthesis (compound heterozygotes). In some cases, deletion of the β gene, δ and β genes or even δ , β and γ genes occurs. In others, unequal crossing-over has produced $\delta\beta$ fusion genes (so called Lepore syndrome named after the first family in which this was diagnosed) (p. 80).

Clinical features

1 Severe anaemia becomes apparent at 3–6 months after birth when the switch from γ - to β -chain production should take place. Typically the infant presents in the first year with failure to thrive, pallor and a swollen abdomen.




Figure 7.8 Examples of mutations that produce β -thalassaemia. These include single base changes, small deletions and insertions of one or two bases affecting introns, exons or the flanking regions of the β -globin gene. FS, 'frameshifts': deletion of nucleotide(s) that places the reading frame out of phase downstream of the lesion; NS, 'non-sense': premature chain termination as a result of a new translational stop codon (e.g. UAA); SPL, 'splicing': inactivation of splicing or new splice sites generated (aberrant splicing) in exons or introns; promoter, CAP, initiation: reduction of transcription or translation as a result of lesion in promoter, CAP or initiation regions; Poly A, mutations on the poly A addition signal resulting in failure of poly A addition and an unstable mRNA.



Figure 7.9 The facial appearance of a child with β -thalassaemia major. The skull is bossed with prominent frontal and parietal bones; the maxilla is enlarged.



Figure 7.10 Skull X-ray in β -thalassaemia major. There is a 'hair-on-end' appearance as a result of expansion of the bone marrow into cortical bone.

- 2 Enlargement of the liver and spleen** occurs as a result of excessive red cell destruction, extramedullary haemopoiesis and later because of iron overload. The large spleen increases blood requirements by increasing red cell destruction and pooling, and by causing expansion of the plasma volume.
- 3 Expansion of bones** caused by intense marrow hyperplasia leads to a thalassaemic facies (Fig. 7.9) and to thinning of the cortex of many bones with a tendency to fractures and bossing of the skull with a 'hair-on-end' appearance on X-ray (Fig. 7.10).
- 4 Thalassaemia major is the disease that most frequently underlies transfusional iron overload.** Regular transfusions are usually commenced in the first year of life and unless the disease is cured by stem cell transplantation, are continued for life. Also iron absorption is increased because of low serum hepcidin levels because of release of proteins, e.g. GDF 15 from increased numbers of early red cell precursors in the marrow. In the children, failure of growth and delayed puberty are frequent, and without iron chelation, death from cardiac damage usually occurs in teenagers. The clinical features (due to hepatic, endocrine and cardiac iron overload) are further discussed in Chapter 4.

- 5 Infections** occur frequently. In infancy, without adequate transfusion, anaemia predisposes to bacterial infections. Pneumococcal, *Haemophilus* and meningococcal infections are likely if splenectomy has been carried out. *Yersinia enterocolitica* occurs, particularly in iron-loaded patients being treated with deferoxamine; it may cause severe gastroenteritis. Iron overload itself also predisposes to bacterial infection, e.g. *Klebsiella*, and to fungal infection. Transfusion of viruses by blood transfusion may occur. As a result of reduction of deaths from cardiac iron overload by improved chelation therapy, infections now account for an increasing proportion of deaths in thalassaemia major.
- 6 Liver disease** in thalassaemia is most frequently a result of hepatitis C but hepatitis B is also common where the virus is endemic. Human immunodeficiency virus (HIV) has been transmitted to some patients by blood transfusion. Iron overload may also cause liver damage.
- 7 Osteoporosis** may occur in well-transfused patients. It is more common in diabetic patients with endocrine abnormalities.
- 8 Hepatocellular carcinoma** incidence is increased in those with iron overload and chronic hepatitis B or C. Ultrasound and measurement of serum alpha-fetoprotein every 6 months is advisable in such patients.

Laboratory diagnosis

- 1 There is a severe hypochromic, microcytic anaemia with normoblasts, target cells and basophilic stippling in the blood film (Fig. 7.11).
- 2 High performance liquid chromatography (HPLC) is now usually used as the first-line method to diagnose haemoglobin disorders (Fig. 7.12a). HPLC or haemoglobin electrophoresis (Fig. 7.12b) reveals absence or almost complete absence of Hb A, with almost all the circulating haemoglobin being Hb F. The Hb A₂ percentage is normal, low or slightly raised. DNA analysis is used to identify the defect on each allele important in antenatal diagnosis.

Treatment

- 1 Regular blood transfusions are needed to maintain the haemoglobin over 100 g/L at all times. This usually requires 2–3 units every 4–6 weeks. Fresh blood, filtered to remove white cells, gives the best red cell survival with the fewest reactions. The patients should be genotyped at the start of the transfusion programme in case red cell antibodies against transfused red cells develop.
- 2 Iron chelation therapy is essential and available drugs have considerably improved life expectancy (see Chapter 4).
- 3 Regular folic acid (e.g. 5 mg/day) is given if the diet is poor.
- 4 Splenectomy may be needed to reduce blood requirements. This should be delayed until the patient is over 6 years old because of the high risk of dangerous infections post-splenectomy. The vaccinations and antibiotics to be given are described in Chapter 10.
- 5 Endocrine therapy is given either as replacement because of end-organ failure or to stimulate the pituitary if puberty is delayed. Diabetes will require insulin therapy. Patients with osteoporosis may need additional therapy with increased

calcium and vitamin D together with a bisphosphonate and appropriate endocrine therapy.

- 6 Immunization against hepatitis B should be carried out in all non-immune patients. Treatment for transfusion-transmitted hepatitis C is given if viral genomes are detected in plasma.
- 7 Allogeneic stem cell transplantation offers the prospect of permanent cure. The success rate (long-term thalassaemia major-free survival) is over 80% in well-chelated younger patients without liver fibrosis or hepatomegaly. A human leucocyte antigen matching sibling (or rarely other family member or matching unrelated volunteer) acts as donor. Failure is mainly a result of recurrence of thalassaemia, death (e.g. from infection) or severe chronic graft-versus-host disease.

β -Thalassaemia trait (minor)

This is a common, usually symptomless, abnormality characterized like α -thalassaemia trait by a hypochromic, microcytic blood picture (MCV and MCH very low) but high red cell count ($>5.5 \times 10^{12}/L$) and mild anaemia (haemoglobin 100–120 g/L). It is usually more severe than α thalassaemia trait. A raised Hb A₂ ($>3.5\%$) confirms the diagnosis. The diagnosis allows the possibility of prenatal counselling. If the partner also has β -thalassaemia trait there is a 25% risk of a thalassaemia major child.

Non-transfusion dependent thalassaemia (thalassaemia intermedia)

This is thalassaemia of moderate severity (haemoglobin 70–100 g/L) without the need for regular transfusions (Table 7.3). It is a *clinical* syndrome caused by a variety of genetic defects: homozygous β -thalassaemia with production of




Figure 7.11 Blood film in β -thalassaemia major post-splenectomy. There are hypochromic cells, target cells and many nucleated red cells (normoblasts). Howell–Jolly bodies are seen in some red cells.

Table 7.3 Non-transfusion dependent thalassaemia (thalassaemia intermedia).

Homozygous β -thalassaemia

Homozygous or compound heterozygotes with mild β^+ -thalassaemia
Coinheritance of α -thalassaemia
Enhanced ability to make fetal haemoglobin (γ -chain production)

Heterozygous β -thalassaemia

Coinheritance of additional α -globin genes ($\alpha\alpha\alpha/\alpha\alpha$ or $\alpha\alpha\alpha/\alpha\alpha\alpha$)
Dominant β -thalassaemia trait

$\delta\beta$ -Thalassaemia and hereditary persistence of fetal haemoglobin

Homozygous $\delta\beta$ -thalassaemia
Heterozygous $\delta\beta$ -thalassaemia/ β -thalassaemia
Homozygous Hb Lepore (some cases)

Haemoglobin H disease




Figure 7.12 (a) High performance liquid chromatography. The different haemoglobins elute at different times from the column and their concentrations are read automatically. In this example, the patient is a carrier of sickle cell disease. **(b)** Haemoglobin electrophoretic patterns in normal adult human blood and in subjects with sickle cell (Hb S) trait or disease, β -thalassaemia trait, β -thalassaemia major, Hb S/ β -thalassaemia or Hb S/Hb C disease and Hb H disease.

more Hb F than usual, e.g. due to mutations of the *BCL11A* gene or with mild defects in β -chain synthesis, by β -thalassaemia trait alone of unusual severity ('dominant' β -thalassaemia), or by β -thalassaemia trait in association with mild globin abnormalities such as Hb Lepore. The coexistence of α -thalassaemia trait improves the haemoglobin level in homozygous β -thalassaemia by reducing the degree of α : β chain imbalance and thus of

α -chain precipitation and ineffective erythropoiesis. Conversely, patients with β -thalassaemia trait who also have excess (five or six) α genes tend to be more anaemic than usual. The patient with thalassaemia intermedia may show bone deformity, enlarged liver and spleen, extramedullary erythropoiesis (Fig. 7.13), leg ulcers, gallstones, osteoporosis and venous thrombosis. Iron overload is caused by increased iron




Figure 7.13 β -Thalassaemia intermedia: MRI scan showing masses of extramedullary haemopoietic tissue arising from the ribs and in the paravertebral region without encroachment of the spinal cord.

absorption and occasional transfusions, e.g. during pregnancy or infections or given to reduce bone deformity. Iron chelation, usually with oral drugs, may be needed to treat this, e.g. if the serum ferritin is greater than $800 \mu\text{g/L}$ or liver iron is greater than 5 mg/g . Splenectomy may be needed to avoid the need for transfusions. Hb H disease (three-gene deletion α -thalassaemia) is a type of thalassaemia intermedia without iron overload or extramedullary haemopoiesis.

$\delta\beta$ -Thalassaemia

This involves failure of production of both β and δ chains. Fetal haemoglobin production is increased to 5–20% in the heterozygous state, which resembles thalassaemia minor haematologically. In the homozygous state only Hb F is present and haematologically the picture is of thalassaemia intermedia.

Haemoglobin Lepore

This is an abnormal haemoglobin caused by unequal cross-over of the β and δ genes to produce a polypeptide chain consisting of the δ chain at its amino end and β chain at its carboxyl end. The $\delta\beta$ -fusion chain is synthesized inefficiently and normal δ - and β -chain production is abolished. The homozygotes show thalassaemia intermedia and the heterozygotes thalassaemia trait.

Hereditary persistence of fetal haemoglobin

These are a heterogeneous group of genetic conditions caused by deletions or cross-overs affecting the production of β and γ chains or, in non-deletion forms, by point mutations upstream from the γ -globin genes or in the *BCL11A* gene (p. 74).

Association of β -thalassaemia trait with other genetic disorders of haemoglobin

The combination of β -thalassaemia trait with Hb E trait usually causes a transfusion-dependent thalassaemia major syndrome, but some cases are intermediate. β -Thalassaemia trait with Hb S trait produces the clinical picture of sickle cell anaemia rather than of thalassaemia (p. 85). β -Thalassaemia trait with Hb D trait causes a hypochromic, microcytic anaemia of varying severity.

Sickle cell anaemia

Sickle cell disease is a group of haemoglobin disorders resulting from the inheritance of the sickle β -globin gene. The sickle β -globin abnormality is caused by substitution of valine for glutamic acid in position 6 in the β chain (Fig. 7.14). Homozygous sickle cell anaemia (Hb SS) is the most common severe syndrome while the doubly heterozygote conditions of Hb S/C and Hb S/ β thal also cause sickling disease. Hb S (Hb $\alpha_2\beta_2^S$) is insoluble and forms crystals when exposed to low oxygen tension (Fig. 7.15). Deoxygenated sickle haemoglobin polymerizes into long fibres, each consisting of seven intertwined double strands with cross-linking (Fig. 7.15). The red cells sickle and may block different areas of the microcirculation or large vessels causing infarcts of various organs. The carrier state is widespread (Fig. 7.4) and is found in up to 30% of West African people, maintained at this level because of the protection against malaria that is afforded by the carrier state.

Homozygous disease

Clinical features

Clinical features are of a severe haemolytic anaemia punctuated by crises. The symptoms of anaemia are often mild in relation to the severity of the anaemia because Hb S gives up oxygen (O_2) to tissues relatively easily compared with Hb A (see Fig. 2.10). The clinical expression of Hb SS is very variable, some patients having

Normal β -chain	Amino acid	pro	glu	glu
	Base composition	CCT	GAG	GAG
Sickle β -chain	Base composition	CCT	GTT	GAG
	Amino acid	pro	val	glu

Figure 7.14 Molecular pathology of sickle cell anaemia. There is a single base change in the DNA coding for the amino acid in the sixth position in the β -globin chain (adenine is replaced by thymine). This leads to an amino acid change from glutamic acid to valine. A, adenine; C, cytosine; G, guanine; glu, glutamic acid; pro, proline; T, thymine; val, valine.




Figure 7.15 The formation of the sickle polymer. Source: Adapted from Bunn H.F. & Aster J.C. (2011). *Hematologic Pathophysiology*. McGraw Hill.

an almost normal life, free of crises, but others develop severe crises even as infants and may die in early childhood or as young adults. Crises may be vaso-occlusive (painful or visceral), aplastic or haemolytic. There may be serious damage to many organs.


Vaso-occlusive crises

Painful


These are the most frequent. They may be sporadic and unpredictable or precipitated by infection, acidosis, dehydration or deoxygenation (e.g. altitude, operations, obstetric delivery, stasis of the circulation, exposure to cold, violent exercise). Infarcts causing severe pain occur in the bones (hips, shoulders and vertebrae are commonly affected) (Fig. 7.16). The 'hand-foot' syndrome (painful dactylitis caused by infarcts of the small bones) is frequently the first presentation of the disease and may lead to digits of varying lengths (Fig. 7.17).

Visceral

These are caused by sickling within organs causing infarction and pooling of blood, often with a severe exacerbation of anaemia. The **acute sickle chest syndrome** is the most common cause of death both in children and adults. It presents with dyspnoea, falling arterial PO_2 , chest pain and pulmonary infiltrates on chest X-ray. Treatment is with analgesia, oxygen, exchange transfusion and ventilatory support if necessary. Hepatic and girdle sequestration



(a)



(b)

Figure 7.16 Sickle cell anaemia. (a) Radiograph of the pelvis of a young man of West Indian origin which shows avascular necrosis with flattening of the femoral heads, more marked on the right, coarsening of the bone architecture and cystic areas in the right femoral neck caused by previous infarcts. (b) Coronal hip MRI image revealing established osteonecrosis of femoral heads bilaterally (yellow arrow) with crescentic sclerotic margin (blue dot) as a consequence of sickle cell. (Courtesy of Dr A. Malhotra.)

crises may lead to severe illness requiring exchange transfusions. Splenic sequestration is typically seen in infants and presents with an enlarging spleen, falling haemoglobin and abdominal pain. Treatment is with transfusion. Attacks tend to be recurrent and splenectomy is often needed. Priapism and liver and kidney damage due to repeated small infarcts are other complications.




Figure 7.17 Sickle cell anaemia: (a) painful swollen fingers (dactylitis) in a child and (b) the hand of an 18-year-old Nigerian boy with the 'hand-foot' syndrome. There is marked shortening of the right middle finger because of dactylitis in childhood affecting the growth of the epiphysis.

Aplastic crises

These occur as a result of infection with parvovirus or from folate deficiency and are characterized by a sudden fall in haemoglobin and reticulocytes, usually requiring transfusion (see Fig. 22.7).

Haemolytic crises

These are characterized by an increased rate of haemolysis and fall in haemoglobin but rise in reticulocytes and usually accompany a painful crisis.

Other organ damage

The most serious is of the brain (a stroke occurs in 7% of all patients) or spinal cord. Up to a third of children have had a silent cerebral infarct by the age of 6 years (Fig. 7.18). Transcranial Doppler ultrasonography detects abnormal blood flow indicative of arterial stenosis. This may be associated with cognitive impairment and predicts for strokes in children. This can be largely prevented by regular blood transfusions in these cases and trials of hydroxycarbamide therapy are in progress.

Ulcers of the lower legs are common, as a result of vascular stasis and local ischaemia (Fig. 7.19). The spleen is enlarged in infancy and early childhood but later is often reduced in size as a result of infarcts (**autosplenectomy**). Pulmonary hypertension detected by Doppler echocardiography and an increased tricuspid regurgitant velocity is common and increases the risk




Figure 7.18 T2-weighted MRI of a 9-year-old girl with sickle cell anaemia, showing five hyperintensities in the deep white matter (arrows), which are silent cerebral infarcts. Source: Courtesy Dr David Rees, Kings College Hospital.



Figure 7.19 Sick cell anaemia: medial aspect of the ankle of a 15-year-old Nigerian boy showing necrosis and ulceration.

of death. A proliferative retinopathy and priapism are other clinical complications. Chronic damage to the liver may occur through microinfarcts. Pigment (bilirubin) gallstones are frequent. The kidneys are vulnerable to infarctions of the medulla with papillary necrosis. Failure to concentrate urine aggravates the tendency to dehydration and crisis, and nocturnal enuresis


may develop. Infections are frequent partly due to hyposplenism. Pneumonia, urinary tract infections and Gram-negative septicaemia are common. Osteomyelitis may also occur, usually from *Salmonella* spp.

Laboratory findings


- 1 The haemoglobin is usually 60–90 g/L – low in comparison to mild or no symptoms of anaemia.
- 2 Sickle cells and target cells occur in the blood (Fig. 7.20). Features of splenic atrophy (e.g. Howell–Jolly bodies) may also be present.
- 3 Screening tests for sickling are positive when the blood is deoxygenated (e.g. with dithionate and Na_2HPO_4).
- 4 HPLC or haemoglobin electrophoresis (Fig. 7.12): in Hb SS, no Hb A is detected. The amount of Hb F is variable and is usually 5–15%, larger amounts are usually associated with a milder disorder.

Treatment

- 1 Prophylactic – avoid those factors known to precipitate crises, especially dehydration, anoxia, infections, stasis of the circulation and cooling of the skin surface.
- 2 Folic acid (e.g. 5 mg once weekly).
- 3 Good general nutrition and hygiene.
- 4 Pneumococcal, *Haemophilus* and meningococcal vaccination and regular oral penicillin are effective at reducing the infection rate with these organisms. Oral penicillin should start at diagnosis and continue at least until puberty. Hepatitis B vaccination is also given as transfusions may be needed.
- 5 Crises – treat by rest, warmth, rehydration by oral fluids and/or intravenous normal saline (3 L in 24 h) and antibiotics if infection is present. Analgesia at the appropriate level should be given. Suitable drugs are paracetamol, a non-steroidal anti-inflammatory agent and opiates. Blood transfusion is given only if there is very severe anaemia with symptoms. Exchange transfusion may be needed, particularly if there is neurological damage, a visceral sequestration crisis or repeated painful crises. This is aimed at achieving an Hb S percentage of less than 30% in severe cases and, after a stroke, is continued for at least 2 years.



(a)



(b)

Figure 7.20 (a) Sick cell anaemia: peripheral blood film showing deeply staining sickle cells, target cells and polychromasia. (b) Homozygous Hb C disease: peripheral blood film showing many target cells, deeply staining rhomboidal and spherocytic cells.

- 6 Particular care is needed in pregnancy and anaesthesia. There is debate as to whether or not patients need transfusions with normal blood to reduce Hb S levels during pregnancy or before delivery or for minor operations. Routine transfusions throughout pregnancy are given to those with a poor obstetric history or a history of frequent crises. Careful anaesthetic and recovery techniques must be used to avoid hypoxaemia or acidosis.
- 7 Transfusions – these are also sometimes given repeatedly as prophylaxis to patients having frequent crises or who have had major organ damage (e.g. of the brain) or show abnormal transcranial Doppler studies. The aim is to suppress Hb S production over a period of several months or even years. Iron overload, best assessed by the total number of units transfused and liver iron (and may need iron chelation therapy), and alloimmunization against donated blood are common problems.
- 8 Hydroxycarbamide can increase Hb F levels and improves the clinical course of children or adults. It is given to those with severe or moderately severe disease, e.g. who are having three or more painful crises each year. It should not be used during pregnancy.
- 9 Stem cell transplantation can cure the disease, with 80% disease free. The mortality rate is less than 10%. Transplantation is only indicated in the severest of cases whose quality of life or life expectancy are substantially impaired.
- 10 Research into other drugs (e.g. butyrate) to enhance Hb F synthesis or to increase the solubility of Hb S is taking place.

Sickle cell trait

This is a benign condition with no anaemia and normal appearance of red cells in a blood film. Haematuria is the most common symptom and is thought to be caused by minor infarcts of the renal papillae. Hb S varies from 25–45% of the total haemoglobin (Fig. 7.12). Care must be taken with anaesthesia, pregnancy and at high altitudes.

Combination of haemoglobin S with other genetic defects of haemoglobin

The most common of these are Hb S/ β -thalassaemia, and sickle cell/C disease. In Hb S/ β -thalassaemia, the MCV and MCH are lower than in homozygous Hb SS. The clinical picture is of sickle cell anaemia; splenomegaly is usual. Patients with Hb SC disease have a particular tendency to thrombosis and pulmonary embolism, especially in pregnancy. In general, when compared with Hb SS disease, they have a higher incidence of retinal abnormalities, milder anaemia, splenomegaly and generally a longer life expectancy. Diagnosis is made by haemoglobin electrophoresis or HPLC, particularly with family studies.

Haemoglobin C disease

This genetic defect of haemoglobin is frequent in West Africa and is caused by substitution of lysine for glutamic acid in the

β -globin chain at the same point as the substitution in Hb S. Hb C tends to form rhomboidal crystals and in the homozygous state there is a mild haemolytic anaemia with marked target cell formation, cells with rhomboidal shape and microspherocytes (Fig. 7.20b). The spleen is enlarged. The carriers show a few target cells only.

Haemoglobin D disease

This is a group of variants all with the same electrophoretic mobility. Heterozygotes show no haematological abnormality while homozygotes have a mild haemolytic anaemia.

Haemoglobin E disease

This is the most common haemoglobin variant in South-East Asia. In the homozygous state, there is a mild microcytic, hypochromic anaemia. Haemoglobin E/ β^0 -thalassaemia, however, resembles homozygous β^0 -thalassaemia both clinically and haematologically.

Prenatal diagnosis of genetic haemoglobin disorders

It is important to give genetic counselling to couples at risk of having a child with a major haemoglobin defect. If a pregnant woman is found to have a haemoglobin abnormality, her partner should be tested to determine whether he also carries a defect. When both partners show an abnormality and there is a risk of a serious defect in the offspring, particularly β -thalassaemia major, it is important to offer antenatal diagnosis. Several techniques are available, the choice depending on the stage of pregnancy and the potential nature of the defect.

DNA diagnosis

The majority of samples are obtained by chorionic villus biopsy, although amniotic fluid cells are sometimes used. Techniques to sample maternal blood for fetal cells or fetal DNA are being developed. Fetal blood may be sampled directly in mid trimester. The DNA is then analysed after amplification by the polymerase chain reaction (PCR). It may be performed by using primer pairs that only amplify individual alleles ('allele-specific priming') or by using consensus primers that amplify all the alleles followed by restriction digestion to detect a particular allele. This is best illustrated by Hb S in which the enzyme DdeI detects the A-T change (Fig. 7.21).

Pre-implantation genetic diagnosis which avoids the need for pregnancy termination involves performing conventional *in vitro* fertilization, followed by removing one or two cells from the blastomeres on day 3. PCR is used to detect thalassaemia mutations so that unaffected blastomeres can be selected for implantation. HLA typing can also be used to select a blastomere HLA matching a previous thalassaemia major child so that the new baby could potentially act as donor for stem cell transplantation into its older sibling. Ethical considerations are important in deciding to use these applications.




Figure 7.21 Sickle cell anaemia: antenatal diagnosis by DdeI-PCR analysis. The DNA is amplified by two primers that span the sickle cell gene mutation site and produces a product of 473 base pairs (bp) in size. The product is digested with the restriction enzyme DdeI and the resulting fragments analysed by agarose gel electrophoresis. The replacement of an adenine base in the normal β -globin gene by thymine results in Hb S and removes a normal restriction site for DdeI, producing a larger 376 bp fragment than the normal 175 and 201 bp fragments in the digested amplified product. In this case, the CVS DNA shows both the normal fragments and the larger sickle cell product and so is AS. The gel shows DNA from the mother (M), father (F), fetal DNA from a chorionic villus sample (CVS), a normal DNA control (AA) and a homozygous sickle cell DNA control (SS). Source: Courtesy of Dr. J. Old.

SUMMARY


- Genetic disorders of haemoglobin fall into two main groups:
 - 1 The thalassaemias in which synthesis of the α or β globin chain is reduced.
 - 2 Structural disorders such as sickle cell anaemia in which an abnormal haemoglobin is produced.
- The α - or β -thalassaemias occur clinically as minor forms with microcytic hypochromic red cells and a raised red cell count with or without anaemia.
- Total absence of function of all four α globin genes causes hydrops fetalis.
- Total absence of function of both β globin genes causes β -thalassaemia major, a transfusion-dependent anaemia associated with iron overload. Thalassaemia

intermedia is a clinical term for a group of disorders showing mild to moderate anaemia and is usually caused by variants of β -thalassaemia.

- The most frequent structural defect of haemoglobin is the sickle mutation in the β -globin chain causing, in the homozygous form, a severe haemolytic anaemia, associated with vaso-occlusive crises. These may be painful, affecting bone, or affect soft tissues (e.g. chest, spleen or central nervous system). Crises may also be haemolytic or aplastic.
- Antenatal diagnosis using PCR technology to amplify chorionic villous DNA is used to detect severe genetic defects of haemoglobin production, with termination of the pregnancy if appropriate.



Now visit www.wileyessential.com/haematology to test yourself on this chapter.


A vertical strip on the left side of the page showing a microscopic view of blood cells. It includes several red blood cells (erythrocytes) which are biconcave and pinkish-red, and a few white blood cells (leukocytes) which are larger and have more complex, granular or lobed nuclei. The background is a dark blue.

CHAPTER 8

The white cells 1: granulocytes, monocytes and their benign disorders

Key topics

■ Granulocytes	89
■ Granulopoiesis	90
■ Clinical applications of myeloid growth factors	91
■ Monocytes	92
■ Disorders of neutrophil and monocyte function	92
■ Causes of leucocytosis and monocytosis	94
■ Neutropenia	95
■ Eosinophilia	96
■ Histiocytic disorders	99
■ Lysosomal storage diseases	99



The white blood cells (leucocytes) may be divided into two broad groups: the **phagocytes** and the **lymphocytes**. Phagocytes comprise the cells of the **innate** immune system, which can act very quickly after an infection, whereas lymphocytes mediate the **adaptive** immune response, which can develop immunological memory, for example after vaccination. Phagocytes can themselves be subdivided into granulocytes (which include neutrophils, eosinophils and basophils) and monocytes. Their normal development and function, and

benign disorders of white blood cells, are dealt with in this chapter (Table 8.1; Fig. 8.1). Lymphocytes are considered in Chapter 9.

The function of phagocytes and lymphocytes in protecting the body against infection is closely connected with two soluble protein systems of the body: **immunoglobulins** and **complement**. These proteins, which may also be involved in blood cell destruction in a number of diseases, are discussed together with the lymphocytes in Chapter 9.

Table 8.1 White cells: normal blood counts.

Adults	Blood count	Children	Blood count
<i>Total leucocytes</i>	4.00–11.0 × 10 ⁹ /L*	<i>Total leucocytes</i>	
Neutrophils	1.8–7.5 × 10 ⁹ /L*	Neonates	10.0–25.0 × 10 ⁹ /L
Eosinophils	0.04–0.4 × 10 ⁹ /L	1 year	6.0–18.0 × 10 ⁹ /L
Monocytes	0.2–0.8 × 10 ⁹ /L	4–7 years	6.0–15.0 × 10 ⁹ /L
Basophils	0.01–0.1 × 10 ⁹ /L	8–12 years	4.5–13.5 × 10 ⁹ /L
Lymphocytes	1.5–3.5 × 10 ⁹ /L		

* Normal black and Middle Eastern subjects may have lower counts. In normal pregnancy the upper limits are: total leucocytes 14.5 × 10⁹/L, neutrophils 11 × 10⁹/L.




Figure 8.1 White blood cells (leucocytes): (a) neutrophil (polymorph); (b) eosinophil; (c) basophil; (d) monocyte; (e) lymphocyte.

Granulocytes

Neutrophil (polymorph)

This cell has a characteristic dense nucleus consisting of between two and five lobes, and a pale cytoplasm with an irregular outline containing many fine pink–blue (azurophilic) or grey–blue granules (Fig. 8.1a). The granules are divided into primary, which appear at the promyelocyte stage, and secondary (specific), which appear at the myelocyte stage and predominate in the mature neutrophil (Fig. 8.7). Both types of granule are lysosomal in origin: the primary contains myeloperoxidase and other acid hydrolases; the secondary contains lactoferrin, lysozyme and other enzymes. The lifespan of neutrophils in the blood is only 6–10 hours.

Neutrophil precursors

These do not normally appear in normal peripheral blood but are present in the marrow (Fig. 8.2). The earliest recognizable precursor is the **myeloblast**, a cell of variable size which has a large nucleus with fine chromatin and usually two to five nucleoli (Fig. 8.2). The cytoplasm is basophilic and no granules are present. The normal bone marrow contains up to 5% of myeloblasts. Myeloblasts give rise to **promyelocytes** which are

slightly larger cells and have developed primary granules in the cytoplasm. These cells then give rise to **myelocytes** which have specific or secondary granules. The nuclear chromatin is now more condensed and nucleoli are not visible. Separate myelocytes of the neutrophil, eosinophil and basophil series can be identified. The myelocytes give rise to **metamyelocytes**, non-dividing cells, which have an indented or horseshoe-shaped nucleus and a cytoplasm filled with primary and secondary granules. Neutrophil forms between the metamyelocyte and fully mature neutrophil are termed 'band', 'stab' or 'juvenile'. These cells may occur in normal peripheral blood. They do not contain the clear, fine filamentous distinction between nuclear lobes that is seen in mature neutrophils.

Monocytes

These are usually larger than other peripheral blood leucocytes and possess a large central oval or indented nucleus with clumped chromatin (Fig. 8.1d). The abundant cytoplasm stains blue and contains many fine vacuoles, giving a ground-glass appearance. Cytoplasmic granules are also often present. The monocyte precursors in the marrow (monoblasts and promonocytes) are difficult to distinguish from myeloblasts and monocytes.




Figure 8.2 The formation of the neutrophil and monocyte phagocytes. Eosinophils and basophils are also formed in the marrow in a process similar to that for neutrophils.

Eosinophils

These cells are similar to neutrophils, except that the cytoplasmic granules are coarser and more deeply red staining and there are rarely more than three nuclear lobes (Fig. 8.1b). Eosinophil myelocytes can be recognized but earlier stages are indistinguishable from neutrophil precursors. The blood transit time for eosinophils is longer than for neutrophils. They enter inflammatory exudates and have a special role in allergic responses, defence against parasites and removal of fibrin formed during inflammation. Thus they play a role in local immunity and tissue repair.

Basophils

These are only occasionally seen in normal peripheral blood. They have many dark cytoplasmic granules which overlie the nucleus and contain heparin and histamine (Fig. 8.1c). In the tissues they become mast cells. They have immunoglobulin E (IgE) attachment sites and their degranulation is associated with histamine release.

Granulopoiesis

Granulocytes and monocytes are formed in the bone marrow from a common precursor cell (see Fig. 1.2). In the granulopoietic series progenitor cells, myeloblasts, promyelocytes and myelocytes form a proliferative or mitotic pool of cells while the metamyelocytes, band and segmented granulocytes make up a post-mitotic maturation compartment (Fig. 8.3). Large numbers of band and segmented neutrophils (10–15 times more than in the blood) are held in

the normal marrow as a ‘reserve pool’. The bone marrow normally contains more myeloid cells than erythroid cells in the ratio of 2:1 to 12:1, the largest proportion being neutrophils and metamyelocytes. Following their release from the marrow, granulocytes spend only 6–10 hours in the circulation before entering tissues where they perform their phagocytic function. They spend on average 4–5 days in the tissues before they are destroyed during defensive action or as the result of senescence. In the bloodstream there are two pools usually of about equal size: the circulating pool (included in the blood count) and a marginating pool (not included in the blood count).

Control of granulopoiesis: myeloid growth factors

The granulocyte series arises from bone marrow progenitor cells which are increasingly specialized. Many growth factors are involved in this maturation process including interleukin-1 (IL-1), IL-3, IL-5 (for eosinophils), IL-6, IL-11, granulocyte–macrophage colony-stimulating factor (GM-CSF), granulocyte CSF (G-CSF) and monocyte CSF (M-CSF) (see Fig. 1.6). The growth factors stimulate proliferation and differentiation and also affect the function of the mature cells on which they act (e.g. phagocytosis, superoxide generation and cytotoxicity in the case of neutrophils) (see Fig. 1.5). They also inhibit apoptosis.

Increased granulocyte and monocyte production in response to an infection is induced by increased production of growth factors from stromal cells and T lymphocytes, stimulated by endotoxin, and cytokines such as IL-1 or tumour necrosis factor (TNF) (Fig. 8.4).




Figure 8.3 Neutrophil kinetics. CSF, colony-stimulating factor; G, granulocyte; IL, interleukin; M, monocyte; SCF, stem cell factor.




Figure 8.4 Regulation of haemopoiesis; pathways of stimulation of leucopoiesis by endotoxin, for example from infection. It is likely that endothelial and fibroblast cells release basal quantities of granulocyte-macrophage colony-stimulating factor (GM-CSF) and granulocyte colony-stimulating factor (G-CSF) in the normal resting state and that this is enhanced substantially by tumour necrosis factor (TNF) and interleukin-1 (IL-1).

Clinical applications of G-CSF

Clinical administration of G-CSF intravenously or subcutaneously produces a rise in neutrophils. Short-acting G-CSF is given daily. A longer-acting PEGylated G-CSF can be given once in 7–14 days). Indications are:

- **Post-chemotherapy, radiotherapy or stem cell transplantation (SCT)** In these situations, G-CSF accelerates granulocytic recovery and shortens the period of neutropenia (Fig. 8.5). This may translate into a reduction of length of time in hospital, antibiotic usage and frequency of infection but periods of extreme neutropenia after intensive chemotherapy cannot be prevented. The injections may also allow repeated courses of chemotherapy, e.g. for lymphoma, to be given on schedule rather than being delayed because of prolonged neutropenia, particularly a problem in older patients.
- **Myelodysplasia and aplastic anaemia** G-CSF has been given alone or in conjunction with erythropoietin in an attempt to improve bone marrow function and the neutrophil count.
- **Severe benign neutropenia** Both congenital and acquired neutropenia, including cyclical and drug-induced neutropenia, often respond well to G-CSF.
- **Peripheral blood stem cell mobilisation** G-CSF is used to increase the number of circulating multipotent progenitors from donors or the patient, improving the harvest of sufficient peripheral blood stem cells for allogeneic or autologous transplantation.




Figure 8.5 Typical effect of granulocyte colony-stimulating factor (G-CSF) on recovery of neutrophils following autologous bone marrow transplantation.

Monocytes

Monocytes spend only a short time in the marrow and, after circulating for 20–40 hours, leave the blood to enter the tissues where they mature and carry out their principal functions. Their extravascular lifespan after their transformation to macrophages (histiocytes) may be as long as several months or even years. In tissues the macrophages become self-replicating without replenishment from the blood. They assume specific functions in different tissues (e.g. skin, gut, liver) (Fig. 8.6). One particularly important lineage is that of **dendritic cells**, which are involved in antigen presentation to T cells (see Chapter 9). GM-CSF and M-CSF are involved in their production and activation.

Disorders of neutrophil and monocyte function

The normal function of neutrophils and monocytes may be divided into three phases.

Chemotaxis (cell mobilization and migration)

The phagocyte is attracted to bacteria or the site of inflammation by chemotactic substances released from damaged tissues,




Figure 8.6 Reticuloendothelial system: distribution of macrophages.

by complement components and by the interaction of leucocyte adhesion molecules with ligands on the damaged tissues. The leucocyte adhesion molecules also mediate recruitment and interaction with other immune cells. They are also variously expressed on endothelial cells and platelets (see Chapter 1).

Phagocytosis

The foreign material (e.g. bacteria, fungi) or dead or damaged cells of the host are phagocytosed (Fig. 8.7). Recognition of a foreign particle is aided by opsonization with immunoglobulin or complement because both neutrophils and monocytes have Fc and C3b receptors (see Chapter 9).

Macrophages have a central role in antigen presentation: processing and presenting foreign antigens on human leucocyte antigen (HLA) molecules to the immune system. They also secrete a large number of growth factors and chemokines which regulate inflammation and immune responses.

Chemokines are chemotactic cytokines which may be produced constitutively and control lymphocyte traffic under




Figure 8.7 Phagocytosis and bacterial destruction. On entering the neutrophil, the bacterium is surrounded by an invaginated surface membrane and fuses with a primary lysosome to form a phagosome. Enzymes from the lysosome attack the bacterium. Secondary granules also fuse with the phagosomes, and new enzymes from these granules including lactoferrin attack the organism. Various types of activated oxygen, generated by glucose metabolism, also help to kill bacteria. Undigested residual bacterial products are excreted by exocytosis.

physiological conditions; inflammatory chemokines are induced or up-regulated by inflammatory stimuli. They bind to and activate cells via chemokine receptors and play an important part in recruiting appropriate cells to the sites of inflammation.

Killing and digestion

This occurs by **oxygen-dependent** and **oxygen-independent** pathways. In the oxygen-dependent reactions, superoxide (O_2^-), hydrogen peroxide (H_2O_2) and other activated oxygen (O_2) species, are generated from O_2 and reduced nicotinamide adenine dinucleotide phosphate (NADPH). In neutrophils, H_2O_2 reacts with myeloperoxidase and intracellular halide to kill bacteria; activated oxygen may also be involved. Nitric oxide (NO), generated through NO synthase from L-arginine, is an oxygen-independent mechanism by which phagocytes also kill microbes. The other non-oxidative microbicidal mechanisms involve microbicidal proteins. These may act alone (e.g. cathepsin G) or in conjunction with H_2O_2 (e.g. lysozyme, elastase). They may also act with a fall in pH within phagocytic vacuoles into which lysosomal enzymes are released. Lactoferrin, an iron-binding protein, is bacteriostatic by depriving bacteria of iron and generating free radicals (Fig. 8.7).

Defects of phagocytic cell function

Chemotaxis

These defects occur in rare congenital abnormalities (e.g. 'lazy leucocyte' syndrome) and in more common acquired

abnormalities, either of the environment e.g. corticosteroid therapy or of the leucocytes themselves e.g. in acute or chronic myeloid leukaemia, myelodysplasia and the myeloproliferative syndromes.

Phagocytosis

These defects usually arise because of a lack of opsonization which may be caused by congenital or acquired causes of hypogammaglobulinaemia or lack of complement components.

Killing

This abnormality is clearly illustrated by the rare X-linked or autosomal recessive chronic granulomatous disease that results from abnormal leucocyte oxidative metabolism. There is an abnormality affecting different elements of the respiratory burst oxidase or its activating mechanism. The patients have recurring infections, usually bacterial but sometimes fungal, which present in infancy or early childhood.

Other rare congenital abnormalities may also result in defects of bacterial killing (e.g. myeloperoxidase deficiency and the Chédiak–Higashi syndrome; see below). Acute or chronic myeloid leukaemia and myelodysplastic syndromes may also be associated with defective killing of ingested microorganisms.

Benign disorders

A number of the hereditary conditions may give rise to changes in granulocyte morphology (Fig. 8.8).




Figure 8.8 Abnormal white blood cells. **(a)** Neutrophil leucocytosis: toxic changes shown by the presence of red–purple granules in the band form neutrophils. **(b)** Neutrophil leucocytosis: a Döhle body can be seen in the cytoplasm of the neutrophil. **(c)** Megaloblastic anaemia: hypersegmented oversized neutrophil in peripheral blood. **(d)** May–Hegglin anomaly: the neutrophils contain basophilic inclusions 2–5 µm in diameter; there is an associated mild thrombocytopenia with giant platelets. **(e)** Pelger–Huët anomaly: coarse clumping of the chromatin in pince nez configuration. **(f)** Chédiak–Higashi syndrome: bizarre giant granules in the cytoplasm of a monocyte. **(g)** Alder's anomaly: coarse violet granules in the cytoplasm of a neutrophil.

Pelger–Huët anomaly

In this uncommon symptomless condition bilobed neutrophils are found in the peripheral blood. Occasional unsegmented neutrophils are also seen. Inheritance is autosomal dominant.

May–Hegglin anomaly

In this rare condition the neutrophils contain basophilic inclusions of RNA (resembling Döhle bodies) in the cytoplasm. There is an associated mild thrombocytopenia with giant platelets. Inheritance is autosomal dominant.

Other rare disorders

In contrast to these two benign anomalies, other rare congenital leucocyte disorders may be associated with severe disease. The Chédiak–Higashi syndrome is inherited in an autosomal recessive manner, and there are giant granules in the neutrophils, eosinophils, monocytes and lymphocytes, accompanied by neutropenia, thrombocytopenia and marked hepatosplenomegaly. Abnormal leucocyte granulation or vacuolation is also seen in patients with rare mucopolysaccharide disorders (e.g. Hurler's syndrome).

Common morphological abnormalities

Figure 8.8 also shows some of the more common abnormalities of neutrophil morphology that can be seen in peripheral blood. Hypersegmented forms occur in megaloblastic anaemia, Döhle bodies and toxic changes in infection. A 'drumstick' (Barr body) appears on the nucleus of a proportion of the neutrophils in normal females and is caused by the presence of two X chromosomes. Pelger cells, seen in the benign congenital abnormality, also occur in patients with acute myeloid leukaemia or myelodysplasia.

Causes of neutrophil leucocytosis

An increase in circulating neutrophils to levels greater than $7.5 \times 10^9/L$ is one of the most frequently observed blood count changes. The causes of neutrophil leucocytosis are given in Table 8.2. Neutrophil leucocytosis is sometimes accompanied by fever as a result of the release of leucocyte pyrogens. Other characteristic features of reactive neutrophilia may include: (a) a 'shift to the left' in the peripheral blood differential white cell count, an increase in the number of band forms and the occasional presence of more primitive cells such as metamyelocytes and myelocytes; (b) the presence of cytoplasmic toxic granulation and Döhle bodies (Fig. 8.8a,b).

The leukaemoid reaction

The leukaemoid reaction is a reactive and excessive leucocytosis usually characterized by the presence of immature cells (e.g. myeloblasts, promyelocytes and myelocytes) in the peripheral

Table 8.2 Causes of neutrophil leucocytosis.

Bacterial infections (especially pyogenic bacterial, localized or generalized)
Inflammation and tissue necrosis (e.g. myositis, vasculitis, cardiac infarct, trauma)
Metabolic disorders (e.g. uraemia, eclampsia, acidosis, gout)
Pregnancy
Neoplasms of all types (e.g. carcinoma, lymphoma, melanoma)
Acute haemorrhage or haemolysis
Drugs (e.g. corticosteroid therapy (inhibits margination): lithium, tetracycline)
Chronic myeloid leukaemia, myeloproliferative disease, polycythaemia vera, myelofibrosis, essential thrombocythaemia
Treatment with G-CSF
Rare inherited disorders
Asplenia

blood. Associated disorders include severe or chronic infections, severe haemolysis or metastatic cancer. Leukaemoid reactions are often particularly marked in children.

Leucoerythroblastic reaction

This is characterised by the presence of erythroblast and granulocyte precursors in the blood (Fig. 8.9). It is due to metastatic




Figure 8.9 Leucoerythroblastic blood film. This shows an erythroblast, promyelocyte, myelocyte and metamyelocytes in a patient with metastatic breast carcinoma in the bone marrow.

Table 8.3 Causes of leucoerythroblastic blood film.

Metastatic neoplasm in the marrow
Primary myelofibrosis
Acute and chronic myeloid leukaemia
Myeloma, lymphoma
Miliary tuberculosis
Severe megaloblastic anaemia
Severe haemolysis
Osteopetrosis

infiltration of the marrow or certain benign or neoplastic blood disorders (Table 8.3).

Neutropenia

The lower limit of the normal neutrophil count is $1.8 \times 10^9/L$ except in black people and in the Middle East where $1.5 \times 10^9/L$ is normal. When the absolute neutrophil level falls below $0.5 \times 10^9/L$ the patient is likely to have recurrent infections and when the count falls to less than $0.2 \times 10^9/L$ the risks are very serious, particularly if there is also a functional defect. Neutropenia may be selective or part of a general pancytopenia (Table 8.4).

Benign ethnic neutropenia

Black people often have a low neutrophil count which is termed benign ethnic neutropenia. Up to 98% of people of West African origin carry a polymorphism in the Duffy antigen chemokine receptor (*DARC*) gene which leads to loss of DARC expression on red cells. This has been selected during evolution because the malaria parasite *Plasmodium vivax* uses DARC as a receptor to enter the red cell. DARC is a chemokine receptor and the loss of its expression on white cells is associated with a lowering of the median neutrophil count by around $0.5 \times 10^9/L$. The reduction in the neutrophil count may result from increased neutrophil margination, but there are no significant clinical consequences. A similar effect is seen in some populations in the Middle East.

Congenital neutropenia

Severe congenital neutropenia (previously called Kostmann's syndrome) usually presents in the first year of life with life-threatening infections. Most cases are dominantly inherited, caused by mutations of the gene *ELA2* coding for neutrophil elastase. Other types are autosomally recessive, or the neutropenia occurs as part of other syndromes, e.g. Wiskott–Aldrich,

Table 8.4 Causes of neutropenia.

Selective neutropenia

Congenital

Acquired

Drug-induced

Anti-inflammatory drugs (phenylbutazone)
 Antibacterial drugs (chloramphenicol, co-trimoxazole, sulfasalazine, imipenem)
 Anticonvulsants (phenytoin, carbamazepine)
 Antithyroids (carbimazole)
 Hypoglycaemics (tolbutamide)
 Phenothiazines (chlorpromazine, thioridazine)
 Psychotropics and antidepressants (clozapine, mianserin, imipramine)
 Miscellaneous (rituximab, gold, penicillamine, mepacrine, furosemide, deferiprone)

Benign (racial or familial)

Cyclical

Immune

Autoimmune

Systemic lupus erythematosus

Felty's syndrome

Hypersensitivity and anaphylaxis

Large granular lymphocytic leukaemia (see p. 203)

Infections

Viral (e.g. hepatitis, influenza, HIV)

Fulminant bacterial infection (e.g. typhoid, miliary tuberculosis)

Part of general pancytopenia (see Table 22.1)

HIV, human immunodeficiency virus.

Schwachman–Diamond (p. 249) or Chédiak–Higashi (p. 94). G-CSF produces a clinical response. Some of the forms predispose to myelodysplasia or acute myeloid leukaemia.

Drug-induced neutropenia

A large number of drugs have been implicated (Table 8.4) which may induce neutropenia either by direct toxicity or immune-mediated damage.

Cyclical neutropenia

This is a rare syndrome with 3–4-week periodicity. Severe but temporary neutropenia occurs. Monocytes tend to rise as the neutrophils fall. Mutation of the gene for neutrophil elastase underlies some cases.

Autoimmune neutropenia

In some cases of chronic neutropenia an autoimmune mechanism can be demonstrated. The antibody may be directed against one of the neutrophil-specific antigens (e.g. NA, NB).

Idiopathic benign neutropenia

An increase in the marginating fraction of blood neutrophils and a corresponding reduction in the circulating fraction is one cause of benign neutropenia. It may be ethnic in origin (see above).

The term chronic idiopathic neutropenia is used for unexplained acquired neutropenia (neutrophil count below normal for the ethnic group), without phasic variations or underlying disease. It is more common in females and thought to be brought about by immune cells causing inhibition of myelopoiesis in the bone marrow.

Clinical features

Severe neutropenia is particularly associated with infections of the mouth and throat. Painful and often intractable ulceration may occur at these sites (Fig. 8.10), on the skin or at the anus.



Figure 8.10 Ulceration of the tongue in severe neutropenia.

Septicaemia rapidly supervenes. Organisms carried as commensals by normal individuals, such as *Staphylococcus epidermidis* or Gram-negative organisms in the bowel, may become pathogens. Other features of infections associated with severe neutropenia are described on p. 138.

Diagnosis

Bone marrow examination is useful in determining the severity of damage in granulopoiesis (i.e. whether there is reduction in early precursors or whether there is reduction only of circulating and marrow neutrophils with late precursors remaining in the marrow). Marrow aspiration and trephine biopsy may also provide evidence of leukaemia, myelodysplasia or other infiltration.

Management

The treatment of patients with acute severe neutropenia is described on p. 138. In many patients with drug-induced neutropenia, spontaneous recovery occurs within 1–2 weeks after stopping the drug. Patients with chronic neutropenia have recurrent infections which are mainly bacterial in origin, although fungal and viral infections (especially herpes) also occur. Early recognition and vigorous treatment with antibiotics, antifungal or antiviral agents, as appropriate, is essential. Prophylactic antifungal agents, e.g. fluconazole, are often given and antibacterial agents, e.g. ciprofloxacin, may reduce the risk but resistance is of concern (see Chapter 12). G-CSF is effective at raising the neutrophil count in a variety of benign chronic neutropenic states. Corticosteroid therapy or splenectomy has been associated with good results in some patients with autoimmune neutropenia. Corticosteroids impair neutrophil function and should not be used indiscriminately in patients with neutropenia. Rituximab (anti-CD20) may also be effective, although it may itself be a cause of neutropenia.

Causes of monocytosis, eosinophil and basophil leucocytosis

Monocytosis

A rise in blood monocyte count above $0.8 \times 10^9/L$ is infrequent. The conditions causing monocytosis are listed in Table 8.5.

Eosinophilic leucocytosis (eosinophilia)

The causes of an increase in blood eosinophils (Fig. 8.11) above $0.4 \times 10^9/L$ are listed in Table 8.6. It is most frequently due to allergic diseases, parasites, skin diseases or drugs. Sometimes no underlying cause is found, no clonal marker can be detected and if the eosinophil count is elevated (above $1.5 \times 10^9/L$) for over 6 months and associated with tissue damage, then the


Table 8.5 Causes of monocytosis.

Chronic bacterial infections: tuberculosis, brucellosis, bacterial endocarditis, typhoid
Connective tissue diseases: SLE, temporal arteritis, rheumatoid arthritis
Protozoan infections
Chronic neutropenia
Hodgkin lymphoma, AML and other malignancies
Chronic myelomonocytic leukaemia
AML, acute myeloid leukemia; SLE, systemic lupus erythematosus.

hypereosinophilic syndrome is diagnosed. In this the heart valves, central nervous system, skin and lungs may be affected; treatment is usually with steroids or cytotoxic drugs. In 25% of cases a clonal T-cell population is present. Loeffler's syndrome is a transient reactive form affecting the lungs and the Churg–Strauss syndrome consists of a vasculitis with eosinophilic granulomas affecting the respiratory tract. In other cases of chronic eosinophilia, often with similar clinical features, a clonal cytogenetic or molecular abnormality is present and the term chronic eosinophilic leukaemia is diagnosed (see p. 164).

Basophil leucocytosis (basophilia)

An increase in blood basophils above $0.1 \times 10^9/L$ is uncommon. The usual cause is a myeloproliferative disorder such as chronic myeloid leukaemia or polycythaemia vera. Reactive basophil increases are sometimes seen in myxoedema, during smallpox or chickenpox infection and in ulcerative colitis.

**Figure 8.11** Eosinophilia.**Table 8.6** Causes of eosinophilia.

Allergic diseases, especially hypersensitivity of the atopic type (e.g. bronchial asthma, hay fever, urticaria and food sensitivity)
Parasitic diseases (e.g. amoebiasis, hookworm, ascariasis, tapeworm infestation, filariasis, schistosomiasis and trichinosis)
Recovery from acute infection
Certain skin diseases (e.g. psoriasis, pemphigus and dermatitis herpetiformis, urticaria and angioedema, atopic dermatitis)
Drug sensitivity
Polyarteritis nodosa, vasculitis, serum sickness
Graft-versus-host disease
Hodgkin disease and some other tumours, especially clonal T-cell disorders
Metastatic malignancy with tumour necrosis
Hypereosinophilic syndrome
Pulmonary syndromes Eosinophilic pneumonia, transient pulmonary infiltrates (Loeffler's syndrome), allergic granulomatosis (Churg–Strauss syndrome), tropical pulmonary eosinophilia
Chronic eosinophilic leukaemia
Myeloproliferative including systemic mastocytosis
GM-CSF, granulocyte–macrophage colony-stimulating factor.

Histiocytic and dendritic cell disorders

Histiocytes are myeloid-derived tissue macrophages. Disorders are listed in Table 8.7.

Dendritic cells

These are specialized antigen-presenting cells found mainly in the skin, lymph nodes, spleen and thymus. They comprise:

- 1 Myeloid-derived cells, including Langerhans' cells, which are present in skin and mucosae and are characterised by the presence of tennis racquet-shaped Birbeck granules seen in electron-microscopy sections in neutrophils, eosinophils, macrophages and lymphocytes; and
- 2 A lymphocyte-derived subset.

The primary role of dendritic cells is antigen presentation to T and B lymphocytes (p. 109).

Table 8.7 Classification of the histiocytic and dendritic cell disorders.

Benign

Dendritic cell-related

Langerhans' cell histiocytosis

Solitary dendritic cell histiocytoma

Histiocyte related

Haemophagocytic lymphohistiocytosis

primary (familial)

secondary (infection, drug, tumour)

Sinus histiocytosis with massive lymphadenopathy (Rosai–Dorfman syndrome)

Malignant

Dendritic and histiocytic sarcomas (localized or disseminated)

AML monocytic and myelomonocytic (p. 146)

Chronic myelomonocytic leukaemia (p. 184)

AML, acute myeloid leukaemia.

Langerhans' cell histiocytosis

Langerhans' cell histiocytosis (LCH) includes diseases previously called histiocytosis X, and falls into three clinical groups: **Letterer–Siwe (multisystem) disease**, **Hand–Schüller–Christian disease** (a triad of bone lesions, polyuria due to posterior pituitary involvement and exophthalmos) and **eosinophilic granuloma** (one or more bone lesions). Thus the disease may be single

organ or multisystem. There is a clonal proliferation of myeloid-derived cells resembling antigen presenting cells of the skin. Mutations of the *BRAF* or *MAP2K1* genes may be present.

The multisystem disease affects children in the first 3 years of life with hepatosplenomegaly, lymphadenopathy and eczematous skin symptoms. Localized lesions may occur, especially in the skull, ribs and long bones, the posterior pituitary causing diabetes insipidus, the central nervous system, gastrointestinal tract and lungs. The lesions include CD1a-positive Langerhans' cells (characterized by the presence of Birbeck granules), eosinophils, lymphocytes, neutrophils and macrophages.

Haemophagocytic lymphohistiocytosis (haemophagocytic syndrome)

This is a rare, recessively inherited or more frequently acquired disease, usually precipitated by a viral (especially Epstein–Barr), bacterial or fungal infection or occurring in association with immunosuppression or tumours. In the familial form, various genes, such as perforin, have been shown to be mutated. Patients present with fever and pancytopenia, often with splenomegaly and liver dysfunction. There are increased numbers of histiocytes in the bone marrow which ingest red cells, white cells and platelets (Fig. 8.12). Clinical features are fever, pancytopenia and multiorgan dysfunction, often with lymphadenopathy, hepatic and splenic enlargement, coagulopathy and CNS signs. Treatment is of the underlying infection, if known, with support care. T-cell activation is implicated in the aetiology. Chemotherapy with etoposide, corticosteroids, ciclosporin or Rituximab (anti-CD20) may be tried. The condition is often fatal.




Figure 8.12 Haemophagocytic lymphohistiocytosis: bone marrow aspirates showing histiocytes that have ingested red cells, erythroblasts and neutrophils.

Sinus histiocytosis with massive lymphadenopathy

This is also known as the Rosai–Dorfman syndrome. There is painless, chronic cervical lymphadenopathy. There may be fever and weight loss. The histology is typical and the condition subsides over months or years.

Malignant diseases of histiocytic or dendritic cells include sarcomas, chronic myelomonocytic leukaemia (see Chapter 16) and a rare type of acute myeloid leukaemia (see Chapter 13).

Lysosomal storage diseases

Gaucher's, Tay–Sachs and Niemann–Pick diseases all result from hereditary deficiency of the enzymes required for glycolipid breakdown.

Gaucher's disease

Gaucher's disease is an uncommon autosomal recessive disorder characterized by an accumulation of glucosylceramide in the lysosomes of reticuloendothelial cells as a result of deficiency of glucocerebrosidase (Fig. 8.13). **Three types** occur: a chronic adult type (type I); an acute infantile neuronopathic type (type II); and a subacute neuronopathic type with onset in childhood or adolescence (type III). **Type I is caused by a variety of mutations in the glucocerebrosidase gene**, one type of which (a single base pair substitution in codon 444) is particularly common in Ashkenazi Jews and explains the high incidence of the disease in this group. In type I the outstanding physical sign is splenomegaly. Moderate liver enlargement and pingueculae (conjunctival deposits) are other characteristics. Often the presenting symptom is easy bruising. This is due to thrombocytopenia with abnormal platelet function and coagulation defects.




Figure 8.13 Gaucher disease results from a deficiency of glucocerebrosidase. Gal, Galactose; Glc, glucose.

In many cases, bone deposits cause bone pain and pathological fractures. Osteoporosis is also frequent. Expansion of the lower end of the femur may produce the 'Erlenmeyer flask deformity' (Fig. 8.14c).

The clinical manifestations are caused by the accumulation of glucocerebroside-laden macrophages in the spleen, liver and bone marrow (Fig. 8.14). Gaucher's disease at all ages is commonly associated with marked anaemia, leucopenia and thrombocytopenia occurring singly or in combination. Gaucher cells are not inert lipid storage containers but are metabolically active, secreting proteins that cause secondary pathology, e.g. pulmonary hypertension, alveolar fibrosis and cholesterol gallstones. Polyclonal hypergammaglobulinaemia or monoclonal gammopathy are frequent with a risk of myeloma. Carriers of a Gaucher mutation also have an increased incidence and earlier onset of Parkinson's disease.

Diagnosis is made by assay of white cell glucocerebrosidase and DNA analysis. Lysosomal enzymes, chitotriosidase and acid phosphatase are raised and useful in monitoring therapy. Serum levels of pulmonary activation regulated cytokine (Parc), angiotensin-converting enzyme (ACE) and ferritin are also elevated.

Enzyme replacement therapy with glucocerebrosidase as imiglucerase (Cerezyme), velaglucerase or taliglucerase, made by recombinant technology and given intravenously once every 2 weeks, is effective in treating the disease with shrinkage of spleen, rise in blood count and improved bone structure (Fig. 8.14). Oral drugs, miglustat or eliglustat, are useful alone in mild forms or in combination with the intravenous enzyme. They inhibit glucosylceramide synthase (Fig. 8.13) and so reduce the amount of substrate being produced in lysosomes. The use of enzyme replacement has virtually eliminated the need for splenectomy but it cannot reverse established osteonecrosis, bone deformation, hepatic, splenic or marrow fibrosis. Stem cell transplantation has been carried out successfully in severely affected patients, usually with type II or III disease.

Niemann–Pick disease

Niemann–Pick disease shows certain clinical and pathological similarities to Gaucher's disease. It is caused by a sphingomyelinase deficiency. The majority of patients are infants who die in the first few years of life, although occasional patients survive to adult life. Massive hepatosplenomegaly occurs and there is usually lung and nervous system involvement with retarded physical and mental development. A 'cherry-red' spot is commonly seen in the retina of affected infants. Pancytopenia is a regular feature and in marrow aspirates 'foam cells' of similar size to Gaucher cells are seen. Chemical analysis of the tissues reveals that the disorder is caused by an accumulation of sphingomyelin and cholesterol.





Figure 8.14 Gaucher's disease: **(a)** bone marrow aspirate – a Gaucher cell with 'fibrillar' cytoplasmic pattern; **(b)** spleen histology – pale clusters of Gaucher cells in the reticuloendothelial cord; **(c)** magnetic resonance imaging (MRI) scan of the left knee of a patient before treatment showing Erlenmeyer flask deformity with expansion of the marrow and thinning of the cortical bone; **(d)** following a year of glucocerebrosidase therapy with subsequent remodelling of bone; and bone marrow trephine biopsy before **(e)** and after **(f)** 2 years of glucocerebrosidase therapy. **(g)** Improvement in blood counts and chitotriosidase levels with glucocerebrosidase therapy. Source: (g) Mehta A.B. & Hughes D.A. In Hoffbrand A.V. *et al.* (eds) (2016) *Postgraduate Haematology*. 7th edn. Reproduced with permission of John Wiley & Sons, Ltd.

- Granulocytes include neutrophils (polymorphs), eosinophils and basophils. They are made in the bone marrow under the control of a variety of growth factors and have a short lifespan in the blood stream before entering tissues.
- Phagocytes (neutrophils and monocytes) are the body's main defence against bacterial infection. Neutrophil leucocytosis occurs in bacterial infection and in other types of inflammation.
- Neutropenia, if severe, predisposes to infections. It may be caused by bone marrow failure, chemotherapy or radiotherapy drugs, immune mechanisms or occur congenitally.
- Eosinophilia is most frequently caused by allergic diseases, including skin diseases, parasitic infections or drugs. It can be caused by a clonal increase in eosinophils, termed chronic eosinophilic leukaemia, or an idiopathic condition, often associated with tissue damage.
- Defects of function of neutrophils and monocytes may affect their chemotaxis, phagocytosis or killing.
- Histiocytes are tissue macrophages derived from circulation monocytes. They may form clonal diseases called Langerhans' cell histiocytosis which affect single or multiple organs.
- The haemophagocytic syndrome involves destruction of red cells, granulocytes and platelets by tissue macrophages.
- Lysosomal storage diseases are caused by inherited defects in the enzymes responsible for breakdown of glycolipids. Gaucher's disease is caused by glucocerebrosidase deficiency and is associated with accumulation of glycolipids in the reticuloendothelial system with splenomegaly, pancytopenia and bone lesions causing the main clinical manifestations. Treatment is with enzyme replacement or substrate reduction therapy



Now visit www.wileyessential.com/haematology to test yourself on this chapter.


A vertical strip on the left side of the page showing a microscopic view of blood cells. It features numerous red blood cells (erythrocytes) as bright red, biconcave discs, and several white blood cells (leukocytes) with distinct nuclei and granules, appearing in shades of pink, purple, and yellow.

CHAPTER 9

The white cells 2: lymphocytes and their benign disorders

Key topics

■ Lymphocytes	103
■ Natural killer cells	104
■ Immunoglobulins	106
■ Antigen–receptor gene rearrangements	107
■ Complement	109
■ The immune response	109
■ Lymphocytosis	111
■ Immunodeficiency	113
■ Differential diagnosis of lymphadenopathy	114






Figure 9.1 Lymphocytes: (a) small lymphocyte; (b) activated lymphocyte; (c) large granular lymphocyte; (d) plasma cell.

Lymphocytes are the immunologically competent cells that assist phagocytes in defence of the body against infection and other foreign invasion (Fig. 9.1). Two unique features characteristic of the immune system are the ability to generate *antigenic specificity* and the phenomenon of *immunological memory*. A complete description of the functions of lymphocytes is beyond the scope of this book, but information essential to an understanding of the diseases of the lymphoid system, and of the role of lymphocytes in other haematological diseases, is included here.

Lymphocytes

In postnatal life, the bone marrow and thymus are the *primary lymphoid organs* in which lymphocytes develop (Fig. 9.2). The *secondary lymphoid organs* in which specific immune responses are generated are the lymph nodes, spleen and lymphoid tissues of the alimentary and respiratory tracts.

B and T lymphocytes

The immune response depends upon two types of lymphocytes, B and T cells (Table 9.1), which derive from the haemopoietic stem cell. B cells mature in the bone marrow and circulate in

the peripheral blood until they undergo recognition of antigen. The **B-cell receptor (BCR)** is membrane-bound immunoglobulin (Fig. 9.3) and binds to a specific antigen. This leads to activation of phosphoinositide 3-kinase (PI3K), which produces a second messenger PIP3 (Fig. 9.4), and also Bruton tyrosine kinase, which phosphorylates further downstream enzymes. The overall effect is to induce expression of AKT, which is an anti-apoptotic pro-survival kinase. Two effective new drugs for treating the B cell neoplasms, chronic lymphocytic leukaemia and non-Hodgkin lymphoma, inhibit Bruton kinase (ibrutinib) and PI3K (idelalisib) (see Chapter 18). The receptor itself is secreted as free soluble immunoglobulin (Fig. 9.4). The B cell then matures into a memory B cell or **plasma cell**. Plasma cells home to the bone marrow and have a characteristic morphology with an eccentric round nucleus, with a 'clock-face' chromatin pattern, and strongly basophilic cytoplasm (Fig. 9.1d). They express intracellular but not surface immunoglobulin.

T cells develop from cells that have migrated to the thymus where they differentiate into mature T cells during passage from the cortex to the medulla. During this process, self-reactive T cells are deleted (negative selection) whereas T cells with some specificity for host human leucocyte antigen (HLA) molecules are selected (positive selection). The mature helper cells




Figure 9.2 Primary and secondary lymphoid organs and blood.

express CD4 and cytotoxic cells express CD8 (Table 9.1). The cells also express one of two T-cell antigen receptor heterodimers, $\alpha\beta$ (>90%) or $\gamma\delta$ (<10%). They recognize antigen only when it is presented at a cell surface (see below).

Natural killer cells

Natural killer (NK) cells are cytotoxic CD8⁺ cells that lack the T-cell receptor (TCR). They are large cells with cytoplasmic

granules and typically express surface molecules CD16 (Fc receptor), CD56 and CD57. NK cells are designed to kill target cells that have a low level of expression of HLA class I molecules, such as may occur during viral infection or on a malignant cell. NK cells do this by displaying a number of receptors for HLA molecules on their surface. When HLA is expressed on the target cell these deliver an inhibitory signal into the NK cell. When HLA molecules are absent on the target

Table 9.1 Functional aspects of T and B cells.

	T cells	B cells
Origin	Thymus	Bone marrow
Tissue distribution	Parafollicular areas of cortex in nodes, peri-arteriolar in spleen	Germinal centres of lymph nodes, spleen, gut, respiratory tract; also subcapsular and medullary cords of lymph nodes
Blood	80% of lymphocytes; CD4 > CD8	20% of lymphocytes
Membrane receptors	TCR for antigen	BCR (= immunoglobulin) for antigen
Function	CD8 ⁺ : CMI against intracellular organisms CD4 ⁺ : T-cell help for antibody production and generation of CMI	Humoral immunity by generation of antibodies
Characteristic surface markers	CD1 CD2 CD3 CD4 or 8 CD5 CD6 CD7 HLA class I HLA class II when activated	CD19 CD20 CD22 CD9 (pre B cells) CD10 (precursor B cells) CD79 a and b HLA class I and II
Genes rearranged	TCR α , β , γ , δ	IgH, Igk, Ig λ

BCR, B-cell receptor; C, complement; CMI, cell-mediated immunity; IFN, interferon; Ig, immunoglobulin; TCR, T-cell receptor; TNF, tumour necrosis factor.




Figure 9.3 Antigen receptors on lymphocytes and their interaction with antigen. **(a)** The B-cell antigen receptor is membrane-bound immunoglobulin (see Fig. 9.4). The antigen-binding immunoglobulin molecule is associated with the CD79 heterodimer which acts as a signal transduction unit. **(b)** The T-cell receptor consists of a number of components that together constitute the CD3 complex. Two antigen-binding chains (α , β) are associated with several proteins (γ , δ , ϵ , ζ) that mediate signal transduction. Antigen is recognized in the form of short peptides held on the surface of HLA molecules. CD8⁺ T cells interact with peptide on a class I HLA molecule and the CD8 heterodimer interacts with the α_3 domain of the class I protein.




Figure 9.4 Signalling from the B-cell receptor after antigen binding occurs through phosphoinositide-3-kinase (PI3K) which produces a second messenger phosphatidylinositol (3)-phosphate (PIP3) which activates Bruton tyrosine kinase (BTK) and AKT. Idelalisib inhibits PI3K and ibrutinib inhibits BTK.

cell this inhibitory signal is lost and the NK cell can then kill its target. In addition, NK cells display antibody-dependent cell-mediated cytotoxicity (ADCC). In this, antibody binds to antigen on the surface of the target cell and then NK cells bind to the Fc portion of the bound antibody and kill the target cell.

Lymphocyte circulation

Lymphocytes in the peripheral blood migrate through *post-capillary venules* into the substance of the lymph nodes or into the spleen or bone marrow. T cells home to the perifollicular zones of the cortical areas of lymph nodes (paracortical areas) (Fig. 9.2) and to the periarteriolar sheaths surrounding the central arterioles of the spleen. B cells selectively accumulate in follicles of the lymph nodes and spleen. Lymphocytes return to the peripheral blood via the efferent lymphatic stream and the thoracic duct.

Immunoglobulins

These are a group of proteins produced by plasma cells and B lymphocytes that bind to antigen. They are divided into five subclasses or **isotypes**: immunoglobulin G (IgG), IgA, IgM, IgD and IgE. IgG, the most common, contributes approximately 80% of normal serum immunoglobulin and is further subdivided into four **subclasses**: IgG₁, IgG₂, IgG₃ and IgG₄. IgA is subdivided into two types. IgM is usually produced first in response to antigen, IgG subsequently and for a more prolonged period. The same cell can switch from IgM to IgG, or to IgA or IgE synthesis. IgA is the main immunoglobulin in secretions, particularly of the gastrointestinal tract. IgD and IgE (involved in delayed hypersensitivity reactions) are minor fractions. Some important biochemical and biological properties of the three main immunoglobulin subclasses are summarized in Table 9.2.

The immunoglobulins are all made up of the same basic structure (Fig. 9.5) consisting of **two heavy chains** which are called gamma (γ) in IgG, alpha (α) in IgA, mu (μ) in IgM, delta (δ) in IgD and epsilon (ϵ) in IgE, and **two light chains** – kappa (κ) or lambda (λ) – which are common to all five




Figure 9.5 Basic structure of an immunoglobulin molecule. Each molecule is made up of two light (κ or λ) (blue areas) and two heavy (purple) chains, and each chain is made up of variable (V) and constant (C) portions, the V portions including the antigen-binding site. The heavy chain (μ , δ , γ , ϵ or α) varies according to the immunoglobulin class. IgA molecules form dimers, while IgM forms a ring of five molecules. Papain cleaves the molecules into an Fc fragment and two Fab fragments.

immunoglobulins. The heavy and light chains each have highly variable regions, which give the immunoglobulin specificity, and constant regions, in which there is virtual complete correspondence in amino acid sequence in all antibodies of a given isotype (e.g. IgA, IgG) or isotype subclass (e.g. IgG₁, IgG₂). IgG antibody can be broken into a constant Fc fragment and two highly variable Fab fragments. IgM molecules are much larger because they consist of five subunits.

Table 9.2 Some properties of the three main classes of immunoglobulin (Ig).

	IgG	IgA	IgM
Molecular weight	140 000	140 000	900 000
Normal serum level (g/L)	6.0–16.0	1.5–4.5	0.5–1.5
Present in	Serum and extracellular fluid	Serum and other body fluids (e.g. of bronchi and gut)	Serum only
Complement fixation	Usual	Yes (alternative pathway)	Usual and very efficient
Placental transfer	Yes	No	No
Heavy chain	(γ_{1-4})	α (α_1 or α_2)	μ

The main role of immunoglobulins is defence of the body against foreign organisms. However, they also have a vital role in the pathogenesis of a number of haematological disorders. Secretion of a specific immunoglobulin from a monoclonal population of lymphocytes or plasma cells causes *paraproteinaemia* (see p. 229). Bence–Jones protein found in the urine in some cases of myeloma consists of a monoclonal secretion of light chains or light-chain fragments (either κ or λ). Immunoglobulins may bind to blood cells in a variety of immune disorders and cause their agglutination (e.g. in cold agglutinin disease; p. 69) or destruction following direct complement lysis or after elimination by the reticuloendothelial system.

Antigen–receptor gene rearrangements

Immunoglobulin gene rearrangements

The immunoglobulin heavy-chain and κ and λ light-chain genes occur on chromosomes 14, 2 and 22, respectively. In the germline state, the heavy-chain gene consists of separate segments for variable (V), diversity (D), joining (J) and constant (C) regions. Each of the V, D and J regions contain a number (n) of different **gene segments** (Fig. 9.6). In cells not committed to immunoglobulin synthesis these gene segments remain in their separate germline state. During early differentiation of B cells there is rearrangement of heavy-chain genes so that one of the V heavy-chain segments combines with one of the D segments, which has itself already combined with one of the J segments. Thus, they form a transcriptionally active gene for the heavy chain. The protein coding segments of the C region mRNA are joined to the V region after splicing out intervening RNA. The class of immunoglobulin that is secreted depends

on which of the nine (4γ , 2α , 1μ , 1δ and 1ϵ) constant regions is used. Diversity is introduced by the variability of which V segment joins with which D and with which J segment. In the arbitrary example shown in Fig. 9.6, V_2 joins with D_1 and J_2 . Additional diversity is generated by the enzyme terminal deoxynucleotidyl transferase (TdT), which inserts a variable number of new bases into the DNA of the D region at the time of gene rearrangement. Further mutation of the V region genes occurs in the germinal centres of secondary lymphoid tissues (called somatic mutation) (see below).

Similar rearrangements occur during generation of the light-chain gene (Fig. 9.7). Enzymes called **recombinases** are needed both in B and T cells to join up the adjacent pieces of DNA after excision of intervening sequences. These recognize certain heptamer- and nonamer-conserved sequences flanking the various gene segments. Mistakes in recombinase activity play an important part in the chromosome translocations of B- or T-cell malignancy.

T-cell receptor gene rearrangements

The vast majority of T cells contain a TCR composed of a heterodimer of α and β chain. In a minority of T cells, the TCR is composed of γ and δ chains. The α , β , γ and δ genes of the TCRs each include V, D, J and C regions. During T-cell ontogeny, rearrangements of these gene segments occur in a similar fashion to those for immunoglobulin genes, thus creating T cells expressing a wide variety (10^8 or more) of TCR structures (Fig. 9.8). TdT is involved in creating additional diversity and the same recombinase enzymes used in B cells are involved in joining up TCR gene segments.




Figure 9.6 Rearrangement of a heavy-chain immunoglobulin gene. One of the V segments is brought into contact with a D, a J and a C (in this case C_μ) segment, forming an active transcriptional gene from which the corresponding mRNA is produced. The DJ rearrangement precedes VDJ joining. The class of immunoglobulin depends on which of the nine constant regions (1μ , 1δ , 4γ , 2α , 1ϵ) is used.




Figure 9.7 The sequence of immunoglobulin gene rearrangement, antigen and immunoglobulin expression during early B-cell development. Intracytoplasmic CD22 is a feature of very early B cells. HLA, human leucocyte antigen; TdT, terminal deoxynucleotidyl transferase.




Figure 9.8 The sequence of events during early T-cell development. The earliest events appear to be the expression of surface CD7, intranuclear terminal deoxynucleotidyl transferase (TdT) and intracytoplasmic CD3 followed by T-cell receptor (TCR) gene rearrangement. Early medullary thymocytes express both CD4 and CD8, but they then lose one or other of these structures.

Complement

This consists of a series of plasma proteins constituting an **amplification enzyme system** which is capable of lysis of bacteria (or of blood cells) or can 'opsonize' (coat) bacteria or cells so that they are phagocytosed. The complement sequence consists of nine major components – C1, C2, etc. – which are activated in turn (denoted thus <HLO>C1<EHLO>) and form a cascade, resembling the coagulation sequence (Fig. 9.9). The most abundant and pivotal protein is C3, which is present in plasma at a level of approximately 1.2 g/L. The early (opsonizing) stages leading to coating of the cells with C3b can occur by two different pathways:

1. The **classical pathway** usually activated by IgG or IgM coating of cells; or
2. The **alternate pathway**, which is more rapid and activated by IgA, endotoxin (from Gram-negative bacteria) and other factors (Fig. 9.9).

Macrophages and neutrophils have C3b receptors and they phagocytose C3b-coated cells. C3b is degraded to C3d detected in the direct antiglobulin test using an anti-complement agent (p. 340). If the complement sequence goes to completion (C9) there is generation of an active phospholipase that punches holes in the cell membrane (e.g. of the red cell or bacterium), causing direct lysis. The complement pathway also generates the biologically active fragments C3a and C5a which act directly on phagocytes to stimulate the respiratory burst (p. 93). Both may trigger anaphylaxis by release of mediators from tissue mast cells and basophils, which cause vasodilatation and increased permeability.

The immune response

One of the most striking features of the immune system is its capacity to produce a highly *specific* response. For both T and B cells this specificity is achieved by the presence of a particular receptor on the lymphocyte surface (Fig. 9.3). Naïve (or virgin) B and T lymphocytes which leave the bone marrow and thymus are resting cells that are not in cell division. They recirculate in the lymphatic system. Specialized macrophages called dendritic cells (DCs; p. 92) process antigens before presenting them to B and T lymphocytes – they are therefore known as **antigen-presenting cells** (APCs). The immune system contains many different lymphocytes. Each of these lymphocytes has a receptor that shows differences in structure from that of any other lymphocyte. Consequently each lymphocyte will bind to only a restricted number of antigens. T and B cells undergo clonal expansion if they meet an APC that is presenting an antigen that can trigger their antigen receptor molecules. At this stage, lymphocytes may develop into effector cells (such as plasma cells or cytotoxic T cells) or memory cells.

DC precursors constitutively migrate at low levels from blood into tissues but their rate of migration is increased at the site of inflammation. Immature DCs are efficient at macropinocytosis which allows them to capture antigens from the environment.

T cells are unable to bind antigen free in solution and require it to be presented on APCs in the form of peptides held on the surface of HLA molecules (Fig. 9.3b). T cells recognize the antigen only when it is presented with 'self' HLA molecules and so are known as **HLA-restricted**. The CD4 molecule on helper cells recognizes class II (HLA-DR, -DQ and






Figure 9.9 The complement (C) sequence. The activated factors are denoted by a bar over the number. Both pathways generate a C3 convertase. In the classic pathway, the convertase is the major (b) component of C4 and C2 (C4b2b). In the alternate pathway, it is the combination of C3b and the major fragment (b) of factor B (C3bBb).

-DR) molecules, whereas the CD8 molecule recognizes class I (HLA-A, -B and -C) molecules (see Fig. 23.5). The antigen recognition site of the TCR is joined to several other subunits in the CD3 complex which together mediate signal transduction. Depending on their cytokine production, CD4⁺ T cells can be broadly subdivided into T helper type 1 (Th1) and Th2 cells. Th1 cells produce mainly IL-2, TNF- β and γ -interferon (IFN- γ), and are important in boosting cell-mediated immunity (and granuloma formation), whereas Th2 cells produce IL-4 and IL-10 and are mainly responsible for providing help for antibody production.

Antigen-specific immune responses are generated in **secondary lymphoid organs** and commence when antigen is carried into a lymph node (Fig. 9.10) on dendritic cells. B cells recognize antigen through their surface immunoglobulin and although most antibody responses require help from antigen-specific T cells, some antigens such as polysaccharides can lead to T-cell independent B cell antibody production. In the follicle, germinal centres arise as a result of continuing response to antigenic stimulation (Fig. 9.11). These consist of follicular dendritic cells (FDCs), which are loaded with antigen, B cells and activated T cells which have migrated up from the T zone.



(a)



(b)

Figure 9.10 (a) Structure of a lymph node. **(b)** Lymph node showing germinal follicles surrounded by a darker mantle zone rim and lighter, more diffuse marginal and T-zone areas.




Figure 9.11 Generation of a germinal centre. B cells activated by antigen migrate from the T zone to the follicle where they undergo massive proliferation. Cells enter the dark zone as centroblasts and accumulate mutations in their immunoglobulin V genes. Cells then pass back into the light zone (Fig. 9.10) as centrocytes. Only those cells that can interact with antigen on follicular dendritic cells and receive signals from antigen-specific T cells (Fig. 9.9) are selected and migrate out as plasma cells and memory cells. Cells not selected die by apoptosis.

Proliferating B cells move to the dark zone of the germinal centre as **centroblasts** where they undergo somatic mutation of their immunoglobulin variable-region genes (Fig. 9.11). Their progeny are known as **centrocytes** and these must be selected for survival by antigen on FDCs, otherwise they undergo apoptosis. If selected they become memory B cells or plasma cells (Fig. 9.11). Plasma cells migrate to the bone marrow and other sites in the RE system and produce high-affinity antibody.

Lymphocytosis

Lymphocytosis often occurs in infants and young children in response to infections that produce a neutrophil reaction in adults. Conditions particularly associated with lymphocytosis are listed in Table 9.3.

Glandular fever is a general term for a disease characterized by fever, sore throat, lymphadenopathy and atypical lymphocytes in the blood. It may be caused by primary infection with Epstein–Barr virus (EBV), cytomegalovirus, human immunodeficiency virus (HIV) or *Toxoplasma*. EBV infection, otherwise known as infectious mononucleosis, is the most common cause.

Table 9.3 Causes of lymphocytosis.

Infections <i>acute:</i> infectious mononucleosis, rubella, pertussis, mumps, acute infectious lymphocytosis, infectious hepatitis, cytomegalovirus, HIV, herpes simplex or zoster <i>chronic:</i> tuberculosis, toxoplasmosis, brucellosis, syphilis
Chronic lymphoid leukaemias (see Chapter 18)
Acute lymphoblastic leukaemia (Chapter 17)
Non-Hodgkin lymphoma (some) (Chapter 20)
Thyrotoxicosis
HIV, human immunodeficiency virus.

Infectious mononucleosis

This is caused by primary infection with EBV and occurs only in a minority of infected individuals – in most cases infection is subclinical. The disease is characterized by a lymphocytosis caused by clonal expansions of T cells reacting against B lymphocytes infected with EBV. The disease is associated with a

high titre of heterophile ('reacting with cells of another species') antibody which reacts with sheep, horse or ox red cells.

Clinical features

The majority of patients are between the ages of 15 and 40 years. A prodromal period of a few days occurs with lethargy, malaise, headaches, stiff neck and a dry cough. In established disease the following features may be found:

- 1 Bilateral cervical lymphadenopathy is present in 75% of cases. Symmetrical generalized lymphadenopathy occurs in 50% of cases. The nodes are discrete and may be tender.
- 2 Over half of patients have a sore throat with inflamed oral and pharyngeal surfaces. Follicular tonsillitis is frequently seen.
- 3 Fever may be mild or severe.
- 4 A morbilliform rash, severe headache and eye signs (e.g. photophobia, conjunctivitis and periorbital oedema) are not uncommon. The rash may follow therapy with amoxicillin or ampicillin.
- 5 Palpable splenomegaly occurs in over half of patients and hepatomegaly in approximately 15%. Approximately 5% of patients are jaundiced.

- 6 Peripheral neuropathy, severe anaemia (caused by autoimmune haemolysis) or purpura (caused by thrombocytopenia) are less frequent complications.

Diagnosis

Pleomorphic atypical lymphocytosis

A moderate rise in white cell count (e.g. $10\text{--}20 \times 10^9/\text{L}$) with an absolute lymphocytosis is usual, and some patients have even higher counts. Large numbers of atypical lymphocytes are seen in the peripheral blood film (Fig. 9.12). These T cells are variable in appearance but most have nuclear and cytoplasmic features similar to those seen during reactive lymphocyte transformation. The greatest number of atypical lymphocytes are usually found between the seventh and tenth day of the illness.

Heterophile antibodies

Heterophile antibodies against sheep or horse red cells may be found in the serum at high titres. Modern slide screening tests, such as the *monospot test*, use formalinized horse red cells to test for the IgM antibodies which agglutinate the cells. Highest




Figure 9.12a–d Infectious mononucleosis: representative 'reactive' T lymphocytes in the peripheral blood film of a 21-year-old man (see also Fig. 9.1b).

titres occur during the second and third week and the antibody persists in most patients for 6 weeks.

EBV antibody

If viral diagnostic facilities are available, a rise in the titre of IgM antibody against the EBV capsid antigen (VCA) may be demonstrated during the first 2–3 weeks. Specific IgG antibody to the EBV nuclear antigen (EBNA) and IgG VCA antibodies develop later and persist for life.

Haematological abnormalities

Haematological abnormalities other than the atypical lymphocytosis are frequent. Occasional patients develop an autoimmune haemolytic anaemia. The IgM autoantibody is typically of the ‘cold’ type and usually shows ‘i’ blood group specificity. Thrombocytopenia is frequent and an autoimmune thrombocytopenic purpura occurs in a smaller number of patients.

Differential diagnosis

The differential diagnosis of infectious mononucleosis includes cytomegalovirus, HIV or toxoplasmosis infection; acute leukaemia; influenza; rubella; bacterial tonsillitis; and infectious hepatitis.

Treatment

In the great majority of patients only symptomatic treatment is required. Corticosteroids are sometimes given to those with severe systemic symptoms. Patients characteristically develop an erythematous rash if given ampicillin therapy. Most patients recover fully 4–6 weeks after initial symptoms. However, convalescence may be slow and associated with severe malaise and lethargy.

Lymphopenia

Lymphopenia may occur in severe bone marrow failure, with corticosteroid and other immunosuppressive therapy, in Hodgkin lymphoma and with widespread irradiation. It also occurs during treatment with the monoclonal antibody alemtuzumab (anti-CD52) and in a variety of immunodeficiency syndromes, the most important of which is HIV infection (see p. 328).

Immunodeficiency

A large number of inherited or acquired deficits in any of the components of the immune system can cause an impaired immune response with increased susceptibility to infection (Table 9.4). A primary lack of T cells (as in AIDS) leads not only to bacterial infections, but also to viral, protozoal, fungal and mycobacterial infections. In some cases, however, lack of specific subsets of T cells which control B-cell maturation may lead to a secondary lack of B-cell function, as in many cases of common variable immunodeficiency, which may develop in children or adults of either sex. In others, a primary defect of B cells or of APCs is present. X-linked agammaglobulinaemia is caused by failure of B-cell development; pyogenic bacterial infections dominate the clinical course. Immunoglobulin replacement therapy can be given by monthly courses of intravenous immunoglobulin. Rare syndromes include aplasia of the thymus, severe combined (T and B) immunodeficiency as a result of adenosine deaminase deficiency and selective deficiencies of IgA or IgM. Acquired immune deficiency occurs after cytotoxic chemotherapy or radiotherapy and is particularly pronounced after stem cell transplantation where dysregulation of the immune system persists for 1 year or

Table 9.4 Classification of immunodeficiencies.

<i>Primary</i>	X-linked agammaglobulinaemia, acquired common variable hypogammaglobulinaemia, selective IgA or IgG subclass deficiencies
B cell (antibody deficiency)	Thymic aplasia (DiGeorge's syndrome), PNP deficiency
T cell	Severe combined immune deficiency (as a result of ADA deficiency or other causes); Bloom's syndrome; ataxia-telangiectasia; Wiskott–Aldrich syndrome
Mixed B and T cell	
<i>Secondary</i>	Myeloma; nephrotic syndrome, protein-losing enteropathy
B cell (antibody deficiency)	AIDS; Hodgkin lymphoma, non-Hodgkin lymphoma; drugs: steroids, ciclosporin, azathioprine, fludarabine, etc.
T cell	Chronic lymphocytic leukaemia, post-stem cell transplantation, chemotherapy/radiotherapy, antiCD52 (alemtuzumab)
T and B cell	

ADA, adenosine deaminase; AIDS, acquired immune deficiency syndrome; Ig, immunoglobulin; PNP, purine nucleoside phosphorylase.

more. Immunodeficiency is also frequently associated with tumours of the lymphoid system including chronic lymphocytic leukaemia and myeloma.

Differential diagnosis of lymphadenopathy

The principal causes of lymphadenopathy are listed in Figure 9.13. The clinical history and examination give essential information. The age of the patient, length of history, associated symptoms of possible infectious or malignant disease, whether the nodes are painful or tender, consistency of the nodes and whether there is **generalized or local lymphadenopathy** are all important. The size of the liver and spleen are assessed. In the case of local node enlargement, it is important to look for inflammatory or malignant disease in the associated lymphatic drainage area.

Further investigations will depend on the initial clinical diagnosis but it is usual to include a full blood count, blood

film and erythrocyte sedimentation rate (ESR). Chest X-ray, monospot test, cytomegalovirus and *Toxoplasma* titres, and anti-HIV and Mantoux tests are frequently needed. In many cases, it will be essential to make a histological diagnosis by node biopsy, usually trucut, in which a core of node is removed under radiological control. Fine needle aspirates give less material, destroy the architecture and so are less reliable in diagnosis. Computed tomography (CT) scanning is valuable in determining the presence and extent of deep node enlargement. Subsequent investigations will depend on the diagnosis made and the patient's particular features. In some cases of deep node enlargement, where enlarged superficial nodes are not available for biopsy, bone marrow or liver biopsy, CT- or ultrasound-guided trucut deep node biopsy may be needed in an attempt to reach a histological diagnosis. Biopsy of the spleen is not performed as it may cause splenic rupture requiring splenectomy.




Figure 9.13 Causes of lymphadenopathy. ALL, acute lymphoblastic leukaemia; CLL, chronic lymphocytic leukaemia; SLE, systemic lupus erythematosus. Malignancies are listed in red.

- Lymphocytes are immunologically competent white cells that are involved in antibody production (B cells) and with the body's defence against viral infection or other foreign invasion (T cells).
- They arise from haemopoietic stem cells in the marrow, T cells being subsequently processed in the thymus.
- B cells secrete antibodies specific for individual antigens.
- T lymphocytes are further subdivided into helper (CD4⁺) and cytotoxic (CD8⁺) cells. They recognise peptides on HLA antigens.
- Natural killer cells are cytotoxic CD8⁺ cells that kill target cells with low expression of HLA molecules.
- The immune response occurs in the germinal centre of lymph nodes and involves B-cell and T-cell proliferation, somatic mutation, selection of cells by recognition of antigen on antigen-presenting cells and formation of plasma cells (which secrete immunoglobulin) or memory B cells.
- Immunoglobulins include five subclasses or isotypes, IgG, IgA, IgM, IgD and IgG, all made up of two heavy chains and two light chains (κ or λ).
- Complement is a cascade of plasma proteins that can either lyse cells or coat (opsonise) them so they are phagocytosed.
- Lymphocytosis is usually caused by acute or chronic infections or by lymphoid leukaemias or lymphomas.
- Lymphadenopathy may be localized (because of local infection or malignancy) or generalized because of infection, non-infectious inflammatory diseases, malignancy or drugs.




Now visit www.wileyessential.com/haematology to test yourself on this chapter.

CHAPTER 10

The spleen

Key topics

■ The anatomy and circulation of the spleen	117
■ The functions of the spleen	117
■ Extramedullary haemopoiesis	118
■ Imaging the spleen	118
■ Splenomegaly	118
■ Hypersplenism	119
■ Hyposplenism	120
■ Splenectomy	120
■ Prevention of infection in hyposplenic patients	121



The spleen has an important and unique role in the function of the haemopoietic and immune systems. As well as being directly involved in many diseases of these systems, a number of important clinical features are associated with hypersplenic and hyposplenic states.

The anatomy and circulation of the spleen

The spleen lies under the left costal margin, has a normal weight of 150–250 g and a length of between 5 and 13 cm. It is normally not palpable but becomes palpable when the size is increased to over 14 cm.

Blood enters the spleen through the splenic artery, which then divides into **trabecular arteries** which permeate the organ and give rise to **central arterioles** (Fig. 10.1). The majority of the arterioles end in **cords** which lack an endothelial lining and form an open blood system unique to the spleen, with a loose reticular connective tissue network lined by fibroblasts and many macrophages. The blood re-enters the circulation by passing across the endothelium of venous **sinuses**. Blood then passes into the splenic vein and back into the general circulation. The cords and sinuses form the **red pulp**, which is 75% of the spleen and has an essential role in monitoring the integrity of red blood cells (see below). A minority of the splenic vasculature is closed in which the arterial and venous systems are connected by capillaries with a continuous endothelial layer.

The central arterioles are surrounded by a core of lymphatic tissue known as **white pulp**, which has an organization similar to lymph nodes (Fig. 10.1). The **periarteriolar lymphatic sheath** (PALS) lies directly around the arteriole and is equivalent to the T zone of the lymph node (p. 104). B cell follicles are found adjacent to the PALS and these

are surrounded by the **marginal** and **perifollicular zones** which are rich in macrophages and dendritic cells. Lymphocytes migrate into white pulp from the sinuses of the red pulp or from vessels that end directly in the marginal and perifollicular zones.

There are both rapid (1–2 min) and slow (30–60 min) blood circulations through the spleen. The slow circulation becomes increasingly important in splenomegaly.

The functions of the spleen

The spleen is the largest filter of the blood in the body and several of its functions are derived from this.

Control of red cell integrity

The spleen has an essential role in the ‘quality control’ of red cells. Excess DNA, nuclear remnants (**Howell–Jolly bodies**) and **siderotic granules** are removed (Fig. 10.2). In the relatively hypoxic environment of the red pulp, and because of plasma skimming in the cords, the membrane flexibility of aged and abnormal red cells is impaired and they are trapped within the sinus where they are ingested by macrophages.

Immune function

The lymphoid tissue in the spleen is in a unique position to respond to antigens filtered from the blood and entering the white pulp. Macrophages and dendritic cells in the marginal zone initiate an immune response and then present antigen to B and T cells to start adaptive immune responses. This arrangement is particularly efficient at mounting an immune response to encapsulated bacteria and explains the susceptibility of hyposplenic patients to these organisms.




Figure 10.1 Schematic representation of the blood circulation in the spleen. Most blood flows in an ‘open’ circulation through splenic cords and regains entry into the circulation through the venous sinuses.




Figure 10.2 Splenic atrophy: peripheral blood film showing Howell–Jolly bodies, Pappenheimer bodies (siderotic granules; see p. XX) and misshapen cells.

Extramedullary haemopoiesis

The spleen, like the liver, undergoes a transient period of haemopoiesis at around 3–7 months of fetal life but is not a site of erythropoiesis in the normal adult. However, haemopoiesis may be re-established in both organs as **extramedullary haemopoiesis**, in disorders such as myelofibrosis or in chronic severe haemolytic and megaloblastic anaemias. Extramedullary haemopoiesis may result either from reactivation of dormant stem cells within the spleen or homing of stem cells from the bone marrow to the spleen.

Imaging the spleen

Ultrasound is the most frequently used technique to image the spleen (Fig. 10.3). This can also detect whether or not blood flow in the splenic, portal and hepatic veins is normal, as well as liver size and consistency. Computed tomography (CT) is

preferable for detecting structural detail and any associated lymphadenopathy (e.g. for lymphoma staging). Magnetic resonance imaging (MRI) also gives improved fine detail structure. Positron emission tomography (PET) is used particularly for initial staging and for detecting residual disease after treatment of lymphoma (Fig. 10.4).

Splenomegaly

Splenic size is increased in a wide range of conditions (Table 10.1). Splenomegaly is usually felt under the left costal margin but massive splenomegaly may be felt as far as the right iliac fossa (see Fig. 15.10). The spleen moves with respiration and a medial splenic notch may be palpable in some cases. In developed countries the most common causes of splenomegaly are infectious mononucleosis, haematological malignancy and portal hypertension, whereas malaria and schistosomiasis are more prevalent on a global scale (Table 10.1). Chronic myeloid




Figure 10.3 Imaging the spleen. (a) Ultrasound of spleen showing splenomegaly (15.3 cm). (b) Normal spleen (10 cm) on computed tomography (CT) scan. (c) CT scan: the spleen is enlarged and shows multiple low-density areas. A diagnosis of diffuse large cell B lymphoma was made histologically after splenectomy. Source: Figures (a) and (b) courtesy of Dr T. Ogunremi.




Figure 10.4 (a) Axial positron emission tomography (PET), (b) fused PET/CT and (c) CT (with barium in stomach) images. These demonstrate a solitary focal area of [18]FDG uptake on the PET image which is seen to localize to the spleen on the fused PET/CT image. Source: Courtesy of Dr V.S. Warbey and Professor G.J.R. Cook.

leukaemia, primary myelofibrosis, lymphoma, Gaucher's disease, malaria, leishmaniasis and schistosomiasis are potential causes of massive splenomegaly.

Tropical splenomegaly syndrome

A syndrome of massive splenomegaly of uncertain aetiology has been found frequently in many malarious zones of the tropics including Uganda, Nigeria, New Guinea and the Congo. Smaller numbers of patients with this disorder are seen in southern Arabia, Sudan and Zambia. Previously, such terms as 'big spleen disease', 'cryptogenic splenomegaly' and 'African macroglobulinaemia' have been used to describe this syndrome.

While it seems probable that malaria is the fundamental cause of tropical splenomegaly syndrome, this disease is not the result of active malarial infection, as parasitaemia is usually scanty and malarial pigment is not found in biopsy material from the liver and spleen. An abnormal host response to the continual presence of malarial antigen, which results in a reactive and relatively benign lymphoproliferative disorder

that predominantly affects the liver and spleen, seems more likely.

Splenomegaly is usually gross and the liver is also enlarged. Portal hypertension may be a feature. The anaemia is often severe and leucopenia is usual; some patients develop a marked lymphocytosis. Serum IgM levels are high and there are high titres of malarial antibody.

Although splenectomy corrects the pancytopenia, there is an increased risk of fulminant malarial infection. Antimalarial therapy has proved successful in the management of many affected patients.

Hypersplenism

Normally, only approximately 5% (30–70 mL) of the total red cell mass is present in the spleen, although up to half of the total marginating neutrophil pool and 30% of the platelet mass may be located there. As the spleen enlarges, the proportion of haemopoietic cells within the organ increases such that up to 40% of the red cell mass, and 90% of platelets (see Fig. 25.9), may be pooled in an enlarged spleen. **Hypersplenism** is a

Table 10.1 Causes of splenomegaly.*Haematological*

Chronic myeloid leukaemia*
 Chronic lymphocytic leukaemia
 Acute leukaemia
 Malignant lymphoma*
 Primary myelofibrosis*
 Polycythaemia vera
 Hairy cell leukaemia
 Thalassaemia major or intermedia*
 Sickle cell anaemia (before splenic infarction)
 Haemolytic anaemias
 Megaloblastic anaemia

Portal hypertension

Cirrhosis
 Hepatic, portal, splenic vein thrombosis

Storage diseases

Gaucher's disease*
 Niemann–Pick disease
 Histiocytosis X

Systemic diseases

Sarcoidosis
 Amyloidosis
 Collagen diseases – systemic lupus erythematosus, rheumatoid arthritis
 Systemic mastocytosis

Infections

Acute: septicaemia, bacterial endocarditis, typhoid, infectious mononucleosis
 Chronic: tuberculosis, brucellosis, syphilis, malaria, leishmaniasis,* schistosomiasis*

*Tropical**

Possibly caused by malaria

* Possible causes of massive (>20 cm) splenomegaly.

clinical syndrome that can be seen in any form of splenomegaly. It is characterized by:

- Enlargement of the spleen;
- Reduction of at least one cell line in the blood in the presence of normal bone marrow function.

Depending on the underlying cause, splenectomy may be indicated if the hypersplenism is symptomatic. It is followed by a rapid improvement in the peripheral blood count.

Hyposplenism

Functional hyposplenism is revealed by the blood film findings of Howell–Jolly bodies or Pappenheimer bodies (siderotic granules on iron staining; Fig. 10.2). The most frequent cause is surgical removal of the spleen, e.g. after traumatic rupture, but hyposplenism can also occur in sickle cell anaemia, gluten-induced enteropathy, amyloidosis and other conditions (Table 10.2).

Table 10.2 Causes of hyposplenism and blood film features.

Causes	Blood film features
Splenectomy	<i>Red cells</i>
Sickle cell disease	Target cells
Essential	Acanthocytes
thrombocythaemia	Irregularly contracted or crenated cells
Adult gluten-induced enteropathy	Howell–Jolly bodies (DNA remnants)
Dermatitis herpetiformis	Siderotic (iron) granules (Pappenheimer bodies)
Amyloidosis	
Rarely	
Inflammatory bowel disease	<i>White cells</i>
Splenic arterial thrombosis	± Mild lymphocytosis, monocytosis
	<i>Platelets</i>
	± Thrombocytosis

Splenectomy

Surgical removal of the spleen may be indicated for treatment of haematological disorders as well as after splenic rupture or for splenic tumours or cysts (Table 10.3). With advances in drug treatment of immune thrombocytopenia, of chemotherapy and immunotherapy for lymphomas and chronic lymphocytic leukaemia, and the introduction of JAK2 inhibitors for treatment of primary myelofibrosis, splenectomy for these conditions is now much less frequently indicated than previously. Splenectomy can be performed by open abdominal laparotomy or by laparoscopic surgery.

The platelet count can often rise dramatically in the early postoperative period, reaching levels of up to $1000 \times 10^9/L$ and peaking at 1–2 weeks. Thrombotic complications are seen in some patients and prophylactic aspirin or heparin are often required during this period. Long-term alterations in the peripheral blood cell count may also be seen, including a persistent thrombocytosis, lymphocytosis or monocytosis.

Table 10.3 Indications for splenectomy.

Splenic rupture
Some cases of:
Chronic immune thrombocytopenia
Haemolytic anaemia, e.g. hereditary spherocytosis, autoimmune haemolytic anaemia, thalassaemia major or intermedia
Chronic lymphocytic leukaemia and lymphomas
Primary myelofibrosis
Tropical splenomegaly

Prevention of infection in hyposplenic patients

Patients with hyposplenism are at lifelong increased risk of infection from a variety of organisms. This is seen particularly in children under the age of 5 years and those with sickle cell anaemia. The most characteristic susceptibility is to the encapsulated bacteriae *Streptococcus pneumoniae*, *Haemophilus influenzae* type B and *Neisseria meningitidis*. *Streptococcus pneumoniae* is a particular concern and can cause a rapid and fulminant disease. Malaria and infection caused by animal bites tend to be more severe in splenectomized individuals.

Measures to reduce the risk of serious infection include the following:

- 1 Patients should be informed about their increased susceptibility to infection and advised to carry a card about their condition. They should be counselled about the increased risk of infection on foreign travel, including that from malaria and tick and animal bites.
- 2 Prophylactic oral penicillin is recommended usually for life. High-risk groups include those aged under 16 years or older than 50 years, splenectomy for a haematological malignancy, history of previous invasive pneumococcal disease. Low-risk adults, if they choose to discontinue penicillin, must be warned to seek immediate medical advice if they develop a high fever. Erythromycin may be prescribed for patients allergic to penicillin. A supply of appropriate antibiotics should also be given for patients to take in the event of onset of fever before medical care is available.
- 3 Vaccination against pneumococcus, haemophilus, meningococcus and influenza infection is recommended (Table 10.4). All types of vaccine, including live vaccines, can be given safely to hyposplenic individuals although the immune response to vaccination may be impaired.

Table 10.4 Recommendations for vaccination of patients with hyposplenism.

Vaccine	Time of vaccination	Revaccination schedule	Comments
1 Pneumococcal polyvalent (23) vaccine (PPV) and/or pneumococcal conjugated vaccine (PCV13)	If possible, at least 2 weeks prior to splenectomy. Alternatively 2 weeks post-splenectomy for all three vaccines	5 yearly	Assessment of antibody response may be useful
2 Combined <i>Haemophilus influenzae</i> Type b conjugate and meningococcal conjugate		Not required	Not required if previously vaccinated
3 Influenza	As soon as available for seasonal protection	Annual	Standard live vaccine

The more immunogenic pneumococcal conjugate vaccine, PCV13, covers fewer (13 strains) than PPV (23 strains) and is used in those with a poor response to PPV or in some protocols as well as PPV.


- The normal adult spleen weighs 150–250g and is 5–13cm in diameter. It has a specialized circulation because the majority of arterioles end in 'cords' which lack an endothelial lining. The blood re-enters the circulation via venous sinuses. The cords and sinuses form the red pulp which monitors the integrity of red blood cells.
- The central arterioles are surrounded by lymphoid tissue called white pulp which is similar in structure to a lymph node.
- The spleen removes aged or abnormal red cells, and excess DNA and siderotic granules, from intact red cells. It also has a specialized immune function against capsulated bacteria, *Pneumococcus*,

Haemophilus influenzae and *Meningococcus*, against which splenectomized patients are immunized. Splenectomy is needed for splenic rupture and in some haematological diseases.

- Enlargement of the spleen (splenomegaly) occurs in many malignant and benign haematological diseases, in portal hypertension and with systemic diseases, including acute and chronic infections.
- Hyposplenism occurs in sickle cell anaemia, gluten-induced enteropathy, amyloidosis and rarely in other diseases.
- Vaccination against capsulated organisms and prolonged antibiotic prophylaxis is needed for patients with absent splenic function.



Now visit www.wileyessential.com/haematology to test yourself on this chapter.




CHAPTER 11

The aetiology and genetics of haematological malignancies

Key topics

■ The incidence of haematological neoplasms	123
■ The aetiology of haemopoietic malignancy	124
■ The genetics of haemopoietic malignancy	125
■ Chromosome nomenclature	127
■ Specific examples of genetic abnormalities in haematological malignancies	129
■ Diagnostic methods used to study malignant cells	131
■ Value of genetic markers in management of haematological malignancy	133






Figure 11.1 Theoretical graph to show the replacement of normal bone marrow cells by a clonal population of malignant cells arising by successive mitotic divisions from a single cell with an acquired genetic alteration.

The haemopoietic malignancies are **clonal diseases** that derive from a single cell in the marrow or peripheral lymphoid tissue that has undergone genetic alteration (Fig. 11.1). In this chapter we discuss the aetiology and genetic basis of haematological malignancy and subsequent chapters discuss the aetiology, diagnosis and management of the individual conditions.

The incidence of haematological neoplasms

Cancer is an important cause of morbidity and mortality and approximately 50% of the UK population will develop cancer in their lifetime. The majority of cancers are epithelial malignancies and haematological cancers represent approximately 7% of all malignant disease (Fig. 11.2). There are major geographical variations in the occurrence of some diseases, most notably chronic lymphocytic leukaemia (CLL) which is the most common leukaemia in white people but rare in the Far East.




Figure 11.2 The relative frequency of the haematological malignancies as a proportion of malignant disease. Source: Smith A. *et al.* (2009) *Br J Haematol* 148: 739–53. Reproduced with permission of John Wiley & Sons.

The aetiology of haemopoietic malignancy

Cancer results from the accumulation of genetic mutations within a cell and the number present varies widely from over 100 in some cancers to about 10 in most haematological malignancies (Fig. 11.3). Factors such as genetic inheritance and environmental lifestyle will influence the risk of developing a malignancy but most cases of leukaemia and lymphoma appear to result simply as a result of the chance acquisition of critical genetic changes.

Inherited factors

The incidence of leukaemia is greatly increased in some genetic diseases such as Down's syndrome where acute leukaemia occurs with a 20- to 30-fold increased frequency. Additional disorders are Bloom's syndrome, Fanconi's anaemia, ataxia telangiectasia, neurofibromatosis, Klinefelter's syndrome and Wiskott–Aldrich syndrome. There is also a weak familial tendency in diseases such as acute myeloid leukaemia (AML), CLL, Hodgkin lymphoma and non-Hodgkin lymphoma (NHL), although the genes predisposing to this increased risk are largely unknown.

Environmental influences

Chemicals

Chronic exposure to industrial solvents or chemicals such as benzene is a known but rare cause of myelodysplasia or AML.

Drugs

Alkylating agents, such as chlorambucil or melphalan, predispose to later development of AML, especially if combined with radiotherapy. Etoposide is associated with a risk of the development of secondary leukaemia associated with balanced translocations including that of the *MLL* gene at 11q23.

Radiation

Radiation, especially to the marrow, is leukaemogenic. This is illustrated by an increased incidence of leukaemia in survivors of the atom bomb explosions in Japan.

Infection

Infections are responsible for around 18% of all cancers and contribute to a range of haematological malignancies.

Viruses

Viral infection is associated with several types of haemopoietic malignancy, especially different subtypes of lymphoma (see Table 20.2). The retrovirus human T-lymphotropic virus type 1 is the cause of adult T-cell leukaemia/lymphoma (see p. 204), although most people infected with this virus do not develop the tumour. Epstein–Barr virus (EBV) is associated with almost all cases of endemic (African) Burkitt lymphoma, post-transplant




Figure 11.3 Average number of somatic mutations found in different types of cancer. Source: Adapted from Vogelstein B *et al.* (2013) *Science* 339: 1546–58. EAC, oesophageal adenocarcinoma; ESCC, oesophageal squamous cell cancer; MSI, microsatellite instability; MSS, microsatellite stable; NSCLC, non-small cell lung cancer; SCLC, small cell lung cancer.

lymphoproliferative disease (see p. 261) and a proportion of patients with Hodgkin lymphoma. Human herpes virus 8 infection (Kaposi's sarcoma-associated virus) causes Kaposi's sarcoma and primary effusion lymphoma (see Table 20.2).

HIV infection is associated with an increased incidence of lymphomas at unusual sites such as the central nervous system. These HIV-associated lymphomas are usually of B-cell origin and of high-grade histology.

Bacteria

Helicobacter pylori infection has been implicated in the pathogenesis of gastric mucosa B-cell (MALT) lymphoma (see p. 221) and antibiotic treatment may even bring about disease remission.

Protozoa

Endemic Burkitt lymphoma occurs in the tropics, particularly in malarial areas. It is thought that malaria may alter host immunity and predispose to tumour formation as a result of EBV infection.

The genetics of haemopoietic malignancy

Malignant transformation occurs as a result of the accumulation of genetic mutations in cellular genes. The genes that are involved in the development of cancer can be divided broadly into two groups: **oncogenes** and **tumour-suppressor genes**.

Oncogenes

Oncogenes arise because of gain-of-function mutations or inappropriate expression pattern in normal cellular genes called **proto-oncogenes** (Fig. 11.4). Oncogenic versions are generated

when the activity of proto-oncogenes is increased or they acquire a novel function. This can occur in a number of ways including translocation, mutation or duplication. In general, these mutations affect the processes of cell signalling, cell differentiation and survival. One of the striking features of haematological malignancies, in contrast to most solid tumours, is their high frequency of chromosomal translocations. Several oncogenes are involved in suppression of apoptosis, of which the best example is *BCL-2* which is overexpressed in follicular lymphoma (see p. 222).

The types of mutations that are detected in a case of cancer fall into two broad groups. **Driver mutations** are those that confer a selective growth advantage to a cancer cell. Recent data suggest that the combinatorial sequence in which different driver mutations occur in a tumour may affect the clinical features of the resulting disease. **Passenger mutations** do not confer a growth advantage and may have already been present in the cell from which the cancer arose or arise as a neutral genetic change in the proliferating cell. It is therefore important that targeted drug treatments are directed against the activity of driver mutations.

Tyrosine kinases

These are enzymes which phosphorylate proteins on tyrosine residues and they are important mediators of intracellular signalling. Mutations of tyrosine kinases underlie a large number of haematological malignancies and they are the targets of many extremely effective new drugs. Common examples discussed in the relevant chapters include **ABL1** in chronic myeloid leukaemia (CML), **JAK2** in myeloproliferative neoplasms, **FLT3** in AML, **KIT** in both systemic mastocytosis and AML, and **Bruton kinase** in chronic lymphocytic leukaemia and other lymphoproliferative disorders.

Tumour-suppressor genes

Tumour-suppressor genes may acquire loss-of-function mutations, usually by point mutation or deletion, which lead to malignant transformation (Fig. 11.4). Tumour-suppressor genes commonly act as components of control mechanisms that regulate entry of the cell from the G₁ phase of the cell cycle into the S phase or passage through the S phase to G₂ and mitosis (see Fig. 1.7). Examples of oncogenes and tumour-suppressor genes involved in haemopoietic malignancies are shown in Table 11.1. The most significant tumour-suppressor gene in human cancer is **p53** which is mutated or inactivated in over 50% of cases of malignant disease, including many haemopoietic tumours.

Clonal progression

Malignant cells appear to arise as a multistep process with acquisition of mutations in different intracellular pathways. This may occur by a **linear evolution**, in which the final clone harbours all the mutations that arose during evolution of the malignancy (Fig. 11.5a), or by **branching evolution**, in which




Figure 11.4 Proliferation of normal cells depends on a balance between the action of proto-oncogenes and tumour-suppressor genes. In a malignant cell this balance is disturbed leading to uncontrolled cell division.

Table 11.1 Some of the more frequent genetic abnormalities within haematological tumours (see also individual chapters).

Disease	Genetic abnormality	Genes involved
AML	t(8;21) translocation t(15;17) translocation Nucleotide insertion Mutation (internal tandem duplication) Mutation	RUNX1-RUNX1T1 (<i>CBFα</i>) <i>PML</i> , <i>RARA</i> <i>NPM</i> <i>FLT3</i> <i>DNMT3A</i>
Secondary AML	11q23 translocations	<i>MLL</i>
Myelodysplasia	-5, del (5q) -7, del (7q)	<i>RPS 14</i> <i>N RAS</i>
CML	t(9;22) translocation	<i>BCR-ABL1</i>
Myeloproliferative disease	Point mutation	<i>JAK-2</i> or <i>CALR</i>
B-ALL	t(12;21) translocation t(9;22) translocation 11q23 translocations	<i>ETV6-RUNX1</i> <i>BCR-ABL1</i> <i>AF4/MLL</i>
T-ALL	Mutation	<i>NOTCH</i>
Follicular lymphoma	t(14;18) translocation	<i>BCL2</i>
Lymphoplasmacytic lymphoma	Mutation	<i>MYD88</i>
Burkitt lymphoma	t(8;14) translocation	<i>MYC</i>
Hairy cell leukaemia	Mutation	<i>BRAF</i>
CLL	17p deletion Mutations	<i>P53</i> <i>NOTCH</i> , <i>SF3B1</i> , <i>ATM</i>

AML, acute myeloid leukaemia; B-ALL, B-acute lymphoblastic leukaemia; CLL, chronic lymphocytic leukaemia; CML, chronic myeloid leukaemia;




Figure 11.5 Multistep origin of a malignant tumour. **(a)** Linear evolution: successive mutations lead to growth advantage of one clone. **(b)** Branching evolution: subclones arise at different stages of the tumour evolution. These subclones share at least one common founder mutation.




Figure 11.6 Examples of different potential patterns of clonal progression between the development, treatment and relapse of leukaemia. Source: Adapted from Bolli N. & Vassiliou G. In Hoffbrand A.V. et al. (eds) (2016) *Postgraduate Haematology 7e*. Reproduced with permission of John Wiley & Sons.

there is more than one clone of cells characterized by different somatic mutations but which share at least one mutation traceable back to a single ancestral cell (Figure 11.5b). During this progression of the disease, one subclone may gradually acquire a growth advantage. Selection of subclones may also occur during treatment which may selectively kill some subclones but allow others to survive and new clones to appear (Fig. 11.6).

Progression of subclinical clonal haematological abnormalities to clinical disease

The use of sensitive immunological and molecular tests has shown many healthy individuals harbour clones of cells which have acquired somatic mutations and from which overt haematological clinical disease may arise (Table 11.2). This is particularly frequent in the elderly. Examples include clones of cells identical to those of chronic lymphocytic leukaemia, which can be present in the blood of individuals with a normal lymphocyte count, and the finding of clones of cells


harbouring mutations, such as of *TET2*, which are characteristic of myeloid malignancy and yet may be present in a normal-appearing bone marrow in nearly a fifth of elderly healthy subjects. Progression of benign monoclonal paraproteinaemia to myeloma has been well recognised for many decades (see Chapter 21).

Chromosome nomenclature

The normal somatic cell has 46 chromosomes and is called *diploid*; ova or sperm have 23 chromosomes and are called *haploid*. The chromosomes occur in pairs and are numbered 1–22 in decreasing size order; there are two sex chromosomes, XX in females, XY in males. *Karyotype* is the term used to describe the chromosomes derived from a mitotic cell which have been set out in numerical order (Fig. 11.7). A somatic cell with more or less than 46 chromosomes is termed *aneuploid*; more than 46 is *hyperdiploid*, less than 46 *hypodiploid*; 46 but with chromosome rearrangements, *pseudodiploid*.

Table 11.2 Examples of clonal abnormalities that may be detected in otherwise healthy individuals and which may or may not progress to clinical disease.

Normal	(Early) Disease	Disease
Increased incidence of CD5+ clones in normal peripheral blood with age	Monoclonal B lymphocytosis	Chronic lymphocytic leukaemia
Increased incidence of clones with mutations, e.g. DNMT3, IDH1 in normal bone marrow with age	Myelodysplasia	Acute myeloid leukaemia
Polyclonal immunoglobulins	IgG monoclonal gammopathy of uncertain significance IgM monoclonal gammopathy of uncertain significance	Smouldering myeloma/symptomatic myeloma Waldenström's macroglobulinaemia
Mutation in utero in fetus, e.g. t(12;21)	Childhood acute lymphoblastic leukaemia	Childhood acute lymphoblastic leukaemia
Presence of t(14;18) clone at low level in peripheral blood	Follicular lymphoma <i>in situ</i>	Follicular lymphoma stage I-IV
Presence of BCR-ABL1 clone at low level in peripheral blood	Chronic myeloid leukaemia	Transformed chronic myeloid leukaemia

**Figure 11.7** A colour-banded karyotype from a normal male. Each chromosome pair shows an individual colour-banding pattern. This involves a cross-species multiple colour chromosome banding technique. Probe sets developed from the chromosomes of gibbons are combinatorially labelled and hybridized to human chromosomes. The success of cross-species colour banding depends on a close homology between host and human conserved DNA, divergence of repetitive DNA and a high degree of chromosomal rearrangement in the host relative to the human karyotype. Source: Courtesy of Professor C.J. Harrison.

Each chromosome has two arms: the shorter called 'p', the longer called 'q'. These meet at the *centromere* and the distal ends of the chromosomes are called *telomeres*. On staining each arm divides into regions numbered outwards from the centromere and each region divides into bands (Fig. 11.8).

When a whole chromosome is lost or gained, a – or + is put in front of the chromosome number. If part of the chromosome is lost it is prefixed with *del* (for deletion). If there is extra material replacing part of a chromosome the prefix *add* (for additional material) is used. Chromosome




Figure 11.8 A schematic representation of a chromosome. The bands may be divided into subbands according to staining pattern.

translocations are denoted by *t*, the chromosomes involved placed in brackets with the lower numbered chromosome first. The prefix *inv* describes an inversion where part of the chromosome has been inverted to run in the opposite direction. An *isochromosome*, denoted by *i*, describes a chromosome with identical chromosome arms at each end; for example, *i*(17q) would consist of two copies of 17q joined at the centromere.

Telomeres

Telomeres are repetitive sequences at the ends of chromosomes. They decrease by approximately 200 base pairs of DNA with every round of replication. When they decrease to a critical length, the cell exits from cell cycle. Germ cells and stem cells, which need to self-renew and maintain a high proliferative potential, contain the enzyme *telomerase*, which can add extensions to the telomeric repeats and compensate for loss at replication, and so enable the cells to continue proliferation. Telomerase is also often expressed in malignant cells but this is probably a consequence of the malignant transformation rather than an initiating factor.

Specific examples of genetic abnormalities in haematological malignancies

The genetic abnormalities underlying the different types of leukaemia and lymphoma are described with the diseases which are themselves increasingly classified according to genetic change rather than morphology. The types of gene abnormality include the following (Fig. 11.9).




Figure 11.9 Types of genetic abnormality which may lead to haemopoietic malignancy. (a) Point mutation; (b) chromosomal translocation; (c) chromosomal deletion or loss; (d) chromosomal duplication; (e) epigenetic changes: DNA methylation or deacetylation of histone tails suppresses gene transcription.

Point mutation

This is illustrated by the Val617Phe mutation in the *JAK2* gene, which leads to constitutive activation of the JAK2 protein in most cases of myeloproliferative disease (see Chapter 15). Mutations within the *RAS* oncogenes or *p53* tumour-suppressor gene are common in many haemopoietic malignancies. The point mutation may involve several base pairs. In 35% of cases of AML the *nucleophosmin* gene shows an insertion of four base pairs, resulting in a frameshift change. Internal tandem duplication or point mutations occur in the *FLT3* gene in 30% of cases of AML.

Translocations

These are a characteristic feature of haematological malignancies and there are two main mechanisms whereby they may contribute to malignant change (Fig. 11.10).




Figure 11.10 The two possible mechanisms by which chromosomal translocations can lead to dysregulated expression of an oncogene.

- 1 Fusion of parts of two genes to generate a chimeric fusion gene that is dysfunctional or encodes a novel '*fusion protein*', e.g. *BCR-ABL1* in t(9;22) in CML (see Fig. 14.1), *RAR α -PML* in t(15;17) in acute promyelocytic leukaemia (see Fig. 13.7) or *ETV6-RUNX1* in t(12; 21) in B-ALL.
- 2 Overexpression of a normal cellular gene, e.g. overexpression of *BCL-2* in the t(14;18) translocation of follicular lymphoma or of *MYC* in Burkitt lymphoma (Fig. 11.11). Interestingly, this class of translocation nearly always involves a *TCR* or immunoglobulin gene locus, presumably as a result of aberrant activity of the recombinase enzyme which is involved in immunoglobulin or *TCR* gene rearrangement in immature B or T cells.

Deletions

Chromosomal deletions may involve a small part of a chromosome, the short or long arm (e.g. 5q-) or the entire chromosome (e.g. monosomy 7). The critical event is probably loss of a tumour-suppressor gene or of a microRNA as in the 13q14 deletion in CLL (see below). Loss of multiple chromosomes is termed hypodiploidy and is seen frequently in ALL.

Duplication or amplification

In chromosomal duplication (e.g. trisomy 12 in CLL) or gene amplification, gains are common in chromosomes 8, 12, 19, 21 and Y. Gene amplification is increasingly recognised within haemopoietic malignancy and an example is that involving the *MLL* gene.




Figure 11.11 The genetic events in one of the three translocations found in Burkitt lymphoma and B-cell acute lymphoblastic leukaemia. The oncogene *c-MYC* is normally located on the long arm (q) of chromosome 8. In the (8; 14) translocation, *c-MYC* is translocated into close proximity to the immunoglobulin heavy-chain gene on the long arm of chromosome 14. Part of the heavy-chain gene (the V region) is reciprocally translocated to chromosome 8. C, constant region; IgH, immunoglobulin heavy-chain gene; J, joining region; V, variable region.

Epigenetic alterations

Gene expression in cancer may be dysregulated not only by structural changes to the genes themselves but also by alterations in the mechanism by which genes are transcribed. These changes are called *epigenetic* and are stably inherited with each cell division so they are passed on as the malignant cell divides. The most important mechanisms are:

- 1 Methylation of cytosine residues in DNA;

- 2 Enzymatic alterations, such as acetylation or methylation, of the histone proteins that package DNA within the cell; and
- 3 Alterations in enzymes that mediate the splicing machinery (see Fig. 16.1).

They are particularly important in the myeloid malignancies. Demethylating agents, such as azacytidine, increase gene transcription and are valuable in treating myelodysplasia (MDS) and AML.

MicroRNAs

Chromosomal abnormalities, both deletions and amplifications, can result in loss or gain of short (micro) RNA sequences. These are normally transcribed but not translated. MicroRNAs (miRNAs) control expression of adjacent or distally located genes. Deletion of the miR15a/miR16-1 locus may be relevant to CLL development with the common 13q14 deletion, and deletions of other microRNAs have been described in AML and other haematological malignancies.

Diagnostic methods used to study malignant cells

Karyotype analysis

Karyotype analysis involves direct morphological analysis of chromosomes from tumour cells under the microscope (see Fig. 14.1). This requires tumour cells to be in metaphase and so cells are cultured to encourage cell division prior to chromosomal preparation.

Fluorescence *in situ* hybridization analysis

Fluorescence *in situ* hybridization (FISH) analysis involves the use of fluorescent-labelled genetic probes which hybridize to specific parts of the genome. It is possible to label each chromosome with a different combination of fluorescent labels (Fig. 11.12). This is a sensitive technique that can detect extra copies of genetic material in both metaphase and interphase (non-dividing) cells or, by using two different probes, reveal chromosomal translocations (see Fig. 14.1e) or reduced chromosome numbers.

Gene sequencing

Gene sequence analysis is used to detect the genetic mutations that can cause malignant disease. **Next generation sequencing (NGS)** can be used to study individual genes of interest; sequencing of the whole exome or genome of the cancer (6.10^9 base pairs) can be performed for moderate cost. This is then compared to the germline sequence of the patient to identify the mutations in the tumour. It is likely that cancer treatment in the future will be based on assessment of the patient's germline genome and the genome of their tumour. The bioinformatic analysis of the information




Figure 11.12 An example of FISH analysis showing the t(12;21) translocation. The green probe hybridizes to the region of the *ETV6* gene on chromosome 12 and the red probe hybridizes to the region of the *RUNX1* gene on chromosome 21. The arrows point to the two derived chromosomes resulting from the reciprocal translocation. Source: Courtesy of Professor C.J. Harrison.

that arises from NGS can be challenging and uses sophisticated computer programmes. Gene sequencing identifies point mutations such as of *JAK2* in the myeloproliferative diseases, *KIT* in systemic mastocytosis (see Chapter 15) and *FLT3* in AML (see Chapter 13).

DNA microarray platforms

DNA microarrays allow a rapid and comprehensive analysis of the pattern of cellular transcription within a cell or tissue by hybridizing labelled cellular mRNA to DNA probes which are immobilized on a slide or microchip (Fig. 11.13). It is valuable in research but is not widely used for diagnosis. An alternative approach to assessing the profile of RNA within a cell is to use NGS to sequence all RNA transcripts ('RNASeq').

Flow cytometry

In this technique, antibodies labelled with different fluorochromes recognize the pattern and intensity of expression of different antigens on the surface of normal and leukaemic cells (Fig. 11.14). Normal cells each have a characteristic profile but malignant cells often express an aberrant phenotype that can be useful in allowing their detection (see Figs 13.6 and 17.8). In the case of B-cell malignancies such as CLL, expression of only one light chain, κ or λ , by the tumour




Figure 11.13 Microarray analysis of genes distinguishing acute lymphoblastic leukaemia (ALL) from acute myeloid leukaemia (AML). The 50 genes most highly correlated on gene-expression microarrays with each of these leukaemias are shown. Each row corresponds to a gene; each column corresponds to the expression value in a particular sample. Expression for each gene is normalized across the samples such that the mean is 0 and the SD is 1. Expression greater than the mean is shaded in red, and that below the mean is shaded in blue. Although the genes as a group appear correlated with the type of leukaemia under study, no single gene is uniformly expressed across the class, illustrating the value of a multigene prediction method. Source: Reproduced courtesy of Golub and colleagues.




Figure 11.14 FACS analysis of acute lymphoblastic leukaemia, T lineage. The blast cells express cCD3, TdT, CD34, CD7 and CD2. Source: Courtesy of Immunophenotyping Laboratory, Royal Free Hospital, London.




Figure 11.15 Immunohistological identification of Reed–Sternberg cells in Hodgkin lymphoma. The binucleate cells stain positively for (a) CD15 and (b) CD30.

cells distinguishes them from a normal polyclonal population which express both κ and λ chains, usually in a κ : λ ratio of 2:1 (see Fig. 20.4). The commonly used markers for the diagnosis of the malignant haematological diseases are listed in the relevant chapters.

Immunohistology (immunocytochemistry)

Antibodies can also be used to stain tissue sections. The fixed sections are incubated with an antibody, washed, and incubated with a second antibody linked to an enzyme, usually peroxidase. A substrate is added that the enzyme converts to a coloured precipitate, usually brown. The presence and architecture of tumour cells can be identified by visualization of stained tissue sections under the microscope (Fig. 11.15). The clonal nature of B-cell malignancies can be shown in tissue sections by staining for κ or λ chains. A malignant clonal population (e.g. in B-cell NHL) will express one or other light chain but not both (see Fig. 20.4).

Value of genetic markers in management of haematological malignancy

The detection of genetic abnormalities is important in several aspects of the management of patients with leukaemia or lymphoma.

Initial diagnosis

Many genetic abnormalities are so specific for a particular disease that their presence determines that diagnosis. An example is the $t(11;14)$ translocation which defines mantle cell lymphoma. Clonal immunoglobulin or TCR gene rearrangements are useful in establishing clonality and determining the lineage of a lymphoid malignancy.

For establishing a treatment protocol

Each major type of haematological malignancy can be further subdivided on the basis of detailed genetic information. For instance, AML is a diverse group of disorders with characteristic genotypes. Individual subtypes respond differently to standard treatment. The $t(8;21)$ and $inv(16)$ subgroups have a favourable prognosis, whereas monosomy 7 carries a poor prognosis. In those with normal cytogenetics, molecular analysis may show FLT3 internal tandem duplication, an unfavourable marker, or NPM1 mutation, which is favourable. The pattern of genetic changes detected by molecular studies in a new case of AML may distinguish those cases with preceding myelodysplasia (and an unfavourable prognosis) from those without myelodysplasia (see p. 145). Treatment strategies are now tailored for the individual and in some instances knowledge of the underlying genetic abnormality can lead to more rational treatment, e.g. the use of all-*trans* retinoic acid in acute promyelocytic leukaemia with $t(15;17)$ (see p. 149). Genetic information is also valuable for giving a prognosis. For instance, hyperdiploidy in ALL is a favourable finding, whereas for most haematological malignancies p53 mutations usually predict for poor prognosis.

Monitoring the response to therapy

The detection of minimal residual disease (disease that cannot be seen by conventional microscopy of the blood or bone marrow) when the patient is in remission after chemotherapy




Figure 11.16 Sensitivity of detection of leukaemic cells in bone marrow using four different techniques. 10^1 to 10^6 = 1 cell in 10 to 1 cell in 10^6 detected.

or stem cell transplantation is possible using the following techniques (in increasing order of sensitivity; Fig. 11.16).

- 1 Cytogenetic analysis.
- 2 Fluorescence-activated cell sorting to detect tumour cells using immunological markers that detect 'leukaemia-specific' combinations of antigens (see Fig. 17.8).

- 3 PCR and/or sequence analysis to detect tumour-specific translocations or mutations specific to the original clone (Fig. 11.17).

These approaches have an important role in determining the treatment of many forms of haemopoietic malignancy.




Figure 11.17 Real-time quantitative polymerase chain reaction (PCR) in acute B-lineage lymphoblastic leukaemia for minimal residual disease using the immunoglobulin heavy chain as target. Primers are designed based on DNA from sequence analysis of the presenting leukaemic clone. Bone marrow samples taken in clinical remission are amplified by PCR using these primers and fluorescent labelled using Sybergreen. The intensity of the signal measures the total DNA molecules amplified in successive cycles. In this example, the intensities of amplification of DNA from two follow-up bone marrow samples (FU1 and FU2) are compared with serial deletions of (10^{-1} to 10^{-4}) of the DNA from the presentation bone marrow. FU1 shows a level of residual disease of approximately 1 in 5000 (0.02%) and FU2 of 1 in 12000 (0.008%). Source: Courtesy of Dr L. Foroni.

SUMMARY


- The haemopoietic malignancies are clonal diseases that derive from a single cell in the marrow or peripheral lymphoid tissue which has undergone genetic alteration.
- They represent approximately 7% of all malignant disease.
- Inherited and environmental factors both predispose to tumour development but the relative contribution of these is usually unclear.
- Infections (viral and bacterial), drugs, radiation and chemicals can all increase the risk of developing a haemopoietic malignancy.
- Haematological malignancies occur because of genetic alterations that lead to increased activation of

oncogenes or decreased activity of tumour suppressor genes. They usually show about 10 acquired genetic mutations and progress in a linear or branching manner.

- These genetic alterations may occur through a variety of mechanisms such as point mutation, chromosomal translocation or gene deletion.
- Important investigations include study of the chromosomes (karyotype analysis), molecular genetics, FISH, mutation analysis, flow cytometry and immunohistochemistry.
- These investigations guide the diagnosis, treatment and monitoring for residual disease of individual cases.



Now visit www.wileyessential.com/haematology to test yourself on this chapter.


A vertical strip on the left side of the page showing a microscopic view of blood cells. It features several large, bright red erythrocytes (red blood cells) and a few smaller, pale pink leukocytes (white blood cells) against a dark blue background.

CHAPTER 12

Management of haematological malignancy

Key topics

■ General support therapy	136
■ Insertion of a central venous catheter	136
■ Blood product support	136
■ Prophylaxis and treatment of infection	138
■ Specific therapies for haematological malignancy	140
■ Drugs used in the treatment of haemopoietic malignancies	140



The treatment of haematological malignancy has improved greatly over the last 40 years. This has resulted from developments in *supportive therapy* and in *specific treatment*. Details of specific treatment are discussed in relation to individual diseases in the appropriate chapter. Support care and general aspects of the agents used in the treatment of haematological malignancy are described here.

General support therapy

Patients with haematological malignancies often present with medical problems related to suppression of normal haemopoiesis and this problem is compounded by the treatments that are given to eradicate the tumour. When considering treatment it is valuable to assess the normal daily living abilities of the patient and here the **ECOG (Eastern Cooperative Oncology Group) Performance Status** is valuable (Table 12.1). It is also important to check for **co-morbidity** such as cardiac, pulmonary and renal disease.

General supportive therapy for patients undergoing intensive treatment often includes the following.

Insertion of a central venous catheter

A central venous catheter is usually inserted for those patients who will need intensive treatment, and is done via a skin tunnel from the chest into the superior vena cava (Fig. 12.1). This gives ease of access for administering chemotherapy, blood products, antibiotics and intravenous feeding. In addition, blood may be taken for laboratory tests.

Blood product support (see Chapter 30)

Red cell and platelet transfusions are used to treat anaemia and thrombocytopenia. A number of particular issues apply to the support of patients with haematological malignancy:


Table 12.1 ECOG performance status.

Grade	ECOG
0	Fully active, able to carry on all pre-disease performance without restriction
1	Restricted in physically strenuous activity but ambulatory and able to carry out work of a light or sedentary nature, e.g. light house work, office work
2	Ambulatory and capable of all selfcare but unable to carry out any work activities. Up and about more than 50% of waking hours
3	Capable of only limited selfcare, confined to bed or chair more than 50% of waking hours
4	Completely disabled. Cannot carry on any selfcare. Totally confined to bed or chair
5	Dead

- 1 The threshold haemoglobin for transfusion will depend on clinical factors such as symptoms and speed of onset of anaemia but most units give red cell support for a Hb less than 80 g/L, with a higher threshold in older patients. In patients needing both red cells and platelets, platelets are given first to reduce the risk of a further fall in the platelet count. Red cell transfusions should be avoided if at all possible in patients with a very high white cell count (above $100 \times 10^9/\text{L}$)



(a)



(b)

Figure 12.1 (a) A central venous line in a patient undergoing intensive chemotherapy. (b) Chest X-ray showing correct placement of a central venous line, in this case a tunnelled triple lumen left internal jugular line. Source: Courtesy of Dr P. Wylie.

because of the hyperviscosity and the risk of precipitating thrombotic episodes as a result of white cell stasis.

- 2 Large volume transfusions, such as 3 units of blood or more, can precipitate pulmonary oedema in older patients and should be given slowly and with clinical monitoring. Diuretics such as furosemide are often given.
- 3 The trigger for platelet transfusion is typically a platelet count below $10 \times 10^9/L$ but this should be doubled in the presence of active bleeding or infection.
- 4 Fresh frozen plasma (FFP) may be needed to reverse coagulation defects.
- 5 Cytomegalovirus (CMV) negative or leucodepleted blood should be given to all patients until it has been shown that they are either CMV seropositive or that they will never be candidates for stem cell transplantation (SCT). This is to prevent transmission of CMV to uninfected patients, as the virus is a significant problem in stem cell transplant recipients (see p. 259).
- 6 Febrile reactions with blood products are not uncommon and should be managed by slowing the infusion and administration of drugs such as antihistamines, pethidine or hydrocortisone. The dosage of steroids should be limited because of concerns with immunosuppression.
- 7 Granulocyte transfusions may be given for severely neutropenic patients with serious infection, not responding to antibiotics but their efficacy is not proven.
- 8 Blood products given to highly immunosuppressed patients (such as those who have received fludarabine, with aplastic anaemia, Hodgkin lymphoma or post-allogeneic SCT) should be irradiated prior to administration to prevent graft-versus-host disease (see p. 257).
- 9 The use of recombinant erythropoietin to reduce the need for blood transfusion and improve patient well-being (e.g. in myeloma or myelodysplasia) is discussed on p. 16.

Haemostasis support

A coagulation screen should be performed regularly on patients undergoing intensive chemotherapy and support with vitamin K or FFP may be required. Cryoprecipitate or antithrombin may be needed for coagulation factor deficiency, such as that precipitated by asparaginase in the management of acute lymphoblastic leukaemia. Antiplatelet drugs such as aspirin or clopidogrel are usually discontinued in patients undergoing intensive chemotherapy and patients on long-term warfarin can be switched to low molecular weight heparin, which can then itself be stopped if the platelet count falls below $50 \times 10^9/L$. Progesterones are given to premenopausal women undergoing intensive chemotherapy to prevent menstruation. Tranexamic acid can be given to reduce haemorrhage in patients with chronic low-grade blood loss.

Anti-emetic therapy

Nausea and vomiting are common side-effects of chemotherapy. A key objective is to try to prevent nausea occurring early in the treatment, as it is more difficult to control once problems have

arisen. The 5-HT₃ (serotonin) receptor antagonists, such as ondansetron or granisetron, can control nausea from intensive chemotherapy in over 60% of cases and the addition of dexamethasone can increase this by approximately 20%. Metoclopramide, prochlorperazine or cyclizine, benzodiazepines (e.g. lorazepam), domperidone or cannabinoids (e.g. nabilone) can all have a role.

Tumour lysis syndrome

Chemotherapy may trigger an acute rise in plasma uric acid, potassium and phosphate and cause hypocalcaemia because of rapid lysis of tumour cells. This syndrome is seen most commonly with rapidly dividing tumours such as lymphoblastic lymphoma or acute leukaemia and can cause acute renal failure. Allopurinol, intravenous fluids and electrolyte replacement are the mainstay of prevention and alkalinization of the urine is sometimes used. Rasburicase, an enzyme that oxidises uric acid to allantoin, is highly effective in controlling hyperuricaemia.

Psychological support

Patients with a diagnosis of malignant disease commonly feel concerns about such issues as the discomfort of treatment, finance, sexuality and fear of mortality. Even when patients achieve a clinical remission there is understandable concern about the chance of disease relapse. Psychological support should be an integral part of the relationship between physician and patient, and patients should be allowed to express their fears and concerns at the earliest opportunity. Most patients value the opportunity to read more about their disorder and many excellent booklets or websites are now available. Teamwork is also crucial and the nursing staff and trained counselors have a vital role in offering support and information during inpatient and outpatient care. Many units have specialist input from clinical psychologists and psychiatric help may occasionally be required. Inadequate communication is perhaps the most common failing of medical teams. The immediate family should be kept informed of the patient's progress whenever possible and appropriate.

Reproductive issues

Men who are to receive cytotoxic drugs should be offered sperm storage, ideally before treatment commences or, if impossible, within a short period of time thereafter. Ethical issues relating to storage or potential usage of tissue in the event of treatment failure will need to be addressed. Permanent infertility in women is less common after chemotherapy although premature menopause may occur. Storage of fertilized ova is often impractical and specialist advice should be obtained in relation to storage of eggs.

Nutritional support

Some degree of weight loss is virtually inevitable in patients undergoing inpatient chemotherapy because of the combination of a poor nutritional intake, malabsorption caused by

neutropenia, topical antiseptics for bathing and chlorhexidine mouthwashes and a 'clean diet' are recommended. The patient is sometimes nursed in a reverse-barrier room. The severity and length of mucositis may be reduced by treatment with recombinant human keratinocyte growth factor (Palifermin). Oral non-absorbed antimicrobial agents, such as neomycin and colistin, reduce gut commensal flora but many units do not use them in order to avoid bacterial resistance. Regular surveillance cultures can be taken to document the patient's bacterial flora and its sensitivity.

Treatment of bacterial infection

Fever is the main indication that infection is present because if neutropenia is present pus will not be formed and infections are often not localized. Fever may be caused by blood products or drugs, but infection is the most common cause and fever of over 38°C in neutropenic patients should be investigated and treated very quickly. Cultures should be taken from any likely focus of infection including blood from central venous lines and peripheral veins, from urine and mouth swabs. The mouth and throat, intravenous catheter site, and perineal and perianal areas are particularly likely foci. A chest X-ray is indicated as chest infections are frequent.

Antibiotic therapy must be started immediately after blood and other cultures have been taken; in many febrile episodes no organisms are isolated.

There are many different antibiotic regimes in use and a close link with the microbiology team is essential. A typical regimen might be based on a single agent, such as a broad-spectrum penicillin (e.g. Tazocin®), meropenem or a broad-spectrum cephalosporin. *Staphylococcus epidermidis* is a common source of fever in patients with intravenous lines and an agent such as teicoplanin, vancomycin or linezolid may be needed. If an infective agent and its antibiotic sensitivities become known, appropriate changes in the regimen are made. If no response occurs within 48–72 hours, changing the antibiotics or treating a fungal or viral infection are considered.

Viral infection

Prophylaxis and treatment of viral infection

Herpes viruses such as herpes simplex, varicella zoster, CMV and Epstein–Barr virus (EBV) undergo latency following primary infection and are never eradicated from the host. Most patients with haematological malignancy have already been infected with these agents and viral reactivation is therefore the most common problem. Aciclovir or valaciclovir is frequently given prophylactically. Herpes simplex is a common cause of oral ulcers but is usually controlled easily by aciclovir. Varicella zoster frequently reactivates in patients with lymphoproliferative diseases to cause shingles, which requires treatment with high doses of aciclovir or valaciclovir. Primary infection, usually in children, can be very serious and immunoglobulin can be used to prevent infection following recent exposure.

Reactivation of CMV infection is particularly important following SCT (see Chapter 23) but may occur following intensive chemotherapy. Failure of immune control of EBV following allogeneic transplantation can lead to outgrowth of a B-cell tumour known as post-transplant lymphoproliferative disease (see p. 261).

Fungal infection

Prophylaxis and treatment of fungal infection

Because of the intensity of current chemotherapy, fungal infections are a major cause of morbidity and mortality. The two major subtypes are yeasts, such as *Candida* species, and moulds, of which *Aspergillus fumigatus* is the most common.

Invasive aspergillosis is a common cause of infectious death in intensively immunocompromised patients (Fig. 12.3). Infection occurs through inhalation of *Aspergillus* spores (conidia) and air filtration systems are used in many haematology wards. The major risk factor is neutropenia – nearly 70% of patients become infected if they are neutropenic for over 34 days. Steroid use is also important, as is age, chemotherapy and antimicrobial history.

The diagnosis of invasive aspergillosis can be difficult. Definitive diagnosis requires demonstration of invasive growth on a biopsy specimen but such evidence is rarely available. Polymerase chain reaction for fungal DNA or enzyme-linked immunosorbent assay (ELISA) for *Aspergillus* galactomannan or β 1–3 d-glucan are useful. High-resolution computed tomography (HRCT) chest scan is valuable and early features are nodular lesions with a 'ground glass' halo appearance. Later on, wedge lesions and the air crescent sign are seen (Fig. 12.4). A high index of suspicion for fungal infection should be maintained and treatment is often started empirically for a fever that has failed to resolve after 2–4 days of antibiotic treatment.




Figure 12.3 Cytology of sputum illustrates the branching septate hyphae of *Aspergillus* (methenamine silver stain).




Figure 12.4 (a) Chest X-ray of patient with pulmonary aspergillosis which shows an area of cavitation containing a central fungal ball (arrow) leading to the typical 'air-crescent' sign. CT scans in *aspergillosis* show hazy ground-glass shadowing with bronchiolar dilatation: (b) and (c). Nodules are seen in early aspergillosis, whereas a fungal ball with surrounding air is typical of more advanced disease (d).

Prophylaxis for patients at risk of *Aspergillus* infection is usually performed with itraconazole, posaconazole or lipid formulation amphotericin. Treatment of established *Aspergillus* infection is with voriconazole, lipid formulation amphotericin, posaconazole or caspofungin. Surgery to remove lung lesions may be needed.

Candida species are a common hospital pathogen and frequently cause oral infection. *Candida* can be significant when isolated from normally sterile body fluids such as blood or urine. Prophylaxis or treatment is usually with fluconazole, itraconazole or caspofungin. Anidulafungin and micafungin are also licensed. *Pneumocystis jirovecii* (*carinii*) is an important cause of pneumonitis. Prophylaxis with co-trimoxazole or nebulized pentamidine is highly effective and is given to those who have received intensive (combination) chemotherapy or fludarabine. Treatment is with high-dose co-trimoxazole.

Specific therapies for haematological malignancy

Specific therapy is aimed at reducing the tumour cell burden by the use of drugs or radiotherapy. The hope in some diseases is to eradicate the tumour completely, and cure rates for haematological malignancy are gradually improving. However, cure is often not achievable, so palliation can also be an important aim.

A wide variety of drugs is used in the management of haemopoietic malignancies and several drugs acting at different sites (Fig. 12.5) are often combined together in regimens that minimize the potential for resistance to occur against a single agent. Many act specifically on dividing cells and their selectivity is dependent on the high proliferation rate within the tumour. Not all tumour cells will be killed by a single course of treatment and it is usual to give several courses of treatment which gradually eradicate the tumour burden. This 'log kill' hypothesis also gives the residual normal haemopoietic cells the opportunity to recover between treatment courses.

Drugs used in the treatment of haemopoietic malignancies

Cytotoxic drugs (Table 12.2)

Alkylating agents, such as chlorambucil, cyclophosphamide and melphalan, are activated to expose reactive alkyl groups which make covalent bonds to molecules within the cell. These have a particular affinity for purines and are thus able to cross-link DNA strands and impair DNA replication, resulting in a block at G₂ and death of the cell by apoptosis (see Fig. 1.9). Bendamustine is a unique drug in this class as it also appears to have activity associated with purine analogue function.




Figure 12.5 The site of action of drugs used in the management of haemopoietic malignancies. ATRA, all-*trans* retinoic acid.

Antimetabolites block metabolic pathways used in DNA synthesis. There are four major groups:

- 1 **Inhibitors of *de novo* DNA synthesis.** Hydroxycarbamide (hydroxyurea) is used widely in the treatment of myeloproliferative disorders. It inhibits the enzyme ribonucleotide reductase which converts ribonucleotides to deoxyribonucleotides. It is not thought to permanently damage DNA and is used in non-malignant disorders such as sickle cell anaemia (p. 85).
- 2 **Folate antagonists**, such as methotrexate (see Fig. 5.5). Methotrexate is widely used alone or in combination with cytosine arabinoside as intrathecal prophylaxis of CNS disease in patients with ALL, acute myeloid leukaemia (AML) or

high-grade non-Hodgkin lymphoma. High systemic doses may also penetrate the CNS. Folinic acid (formyl THF) is able to overcome the activity of methotrexate and is sometimes administered to 'rescue' normal cells after high-dose methotrexate therapy.

- 3 **Pyrimidine analogues** include cytosine arabinoside (cytarabine; ara-C) which is an analogue of 2'-deoxycytidine and is incorporated into DNA, where it inhibits DNA polymerase and blocks replication.
- 4 **Purine analogues** include fludarabine (which inhibits DNA synthesis in a manner similar to ara-C), mercaptopurine, azathioprine, bendamustine, clofarabine and pentostatin.

Table 12.2 Drugs used in the treatment of leukaemia and lymphoma.

	Mechanism of action	Particular side-effects*
<i>Alkylating agents</i>		
Bendamustine	Alkylating agent and purine analogue	Myelosuppression
Cyclophosphamide	Cross-link DNA, impede RNA formation	Haemorrhagic cystitis, cardiomyopathy, loss of hair
Chlorambucil		Marrow aplasia, hepatic toxicity, dermatitis
Melphalan		Marrow aplasia
<i>Antimetabolites</i>		
Hydroxycarbamide (hydroxyurea)	Inhibits ribonucleotide reductase	Pigmentation, nail dystrophy, skin keratosis, epitheliomas
Methotrexate	Inhibit pyrimidine or purine synthesis or incorporation into DNA	Mouth ulcers, gut toxicity
Cytosine arabinoside	Inhibits DNA synthesis	CNS especially cerebellar toxicity and conjunctivitis at high doses
6-Mercaptopurine [†] , 6-thioguanine [†]	Purine analogue	Jaundice, gut toxicity
Clofarabine	Purine analogue	Myelosuppression
Fludarabine 2-Chlorodeoxyadenosine Deoxycoformycin	Inhibit adenosine deaminase or other purine pathways	Immunosuppression (low CD4 counts); renal and neurotoxicity (at high doses)
<i>Cytotoxic antibiotics</i>		
Anthracyclines (e.g. daunorubicin) Hydroxodaunorubicin (Adriamycin) Mitoxantrone Idarubicin	Bind to DNA and interfere with mitosis	Cardiac toxicity, hair loss
Bleomycin	DNA breaks	Pulmonary fibrosis, skin pigmentation
<i>Plant derivatives</i>		
Vincristine (Oncovin [®]), vinblastine	Spindle damage	Neuropathy (peripheral or bladder or gut)
Etoposide	Mitotic inhibitor	Hair loss, oral ulceration
<i>Demethylating agents</i>		
Azacitidine, decitabine	Inhibit DNA methyltransferase	Myelosuppression
<i>Signal transduction inhibitors</i>		
Imatinib, dasatinib, nilotinib, bosutinib	Inhibit BCR-ABL tyrosine kinase	Myelosuppression, fluid retention
Ibrutinib	Inhibit BTK protein	Bleeding
Idelalisib	Inhibit PI3Kdelta	Colitis
Ruxolitinib	Inhibit JAK2	Marrow suppression
<i>Miscellaneous</i>		
Corticosteroids	Lymphoblast lysis	Diabetes, osteoporosis, psychosis,
Trans-retinoic acid	Induces differentiation	Skin hyperkeratosis, leucocytosis and pleural effusion

	Mechanism of action	Particular side-effects*
Arsenic	Induces differentiation or apoptosis	Hyperleucocytosis, cardiac
α -Interferon	Activation of RNAase and natural killer activity	Flu-like symptoms, thrombocytopenia, leucopenia, weight loss
Bortezomib, carfilzomib	Proteasome inhibition	Neuropathy
L-Asparaginase	Deprive cells of asparagine	Hypersensitivity, low albumin and coagulation factors, pancreatitis
Thalidomide, lenalidomide, pomalidomide	Immunomodulation	Neuropathy, constipation, thrombosis
<i>Monoclonal antibodies</i>		
Rituximab, ofatumumab, obinutuzumab (anti-CD20)	Induction of apoptosis	Infusion reactions, immunosuppression
Alemtuzumab (anti-CD52)	Lysis of target cell by complement fixation	Infusion reactions, immunosuppression
Ibritumomab (Zevalin®) (anti-CD20 ^{+90Y})	Toxicity to bound cell	Myelosuppression, nausea
Mylotarg® (anti-CD33)	Kills myeloid cells	Myelosuppression

* Many of the drugs cause nausea, vomiting, mucositis and bone marrow toxicity, and in large doses infertility. Tissue necrosis is a problem if the drugs are extravasated during infusion.

† Allopurinol potentiates the action and side-effects of 6-mercaptopurine.

Cytotoxic antibiotic drugs include the anthracyclines, such as doxorubicin, hydroxodaunorubicin, epirubicin and mitozantrone. These are able to intercalate into DNA and then bind strongly to topoisomerases which are critical for relieving torsional stress in replicating DNA by nicking and resealing DNA strands. If topoisomerase activity is blocked, DNA replication cannot take place.

Bleomycin is a metal chelating antibiotic that generates superoxide radicals within cells that degrade preformed DNA. It is active on non-cycling cells.

Plant derivatives include the vinca alkaloids such as vincristine, which is derived from the periwinkle plant. It binds to tubulin and prevents its polymerization to microtubules. This blocks cell division in metaphase. Etoposide inhibits topoisomerase action.

Targeted drugs

A wide range of targeted drugs which block specific proteins are now in use and are likely to eventually replace the cytotoxic agents described above.

BCR-ABL inhibitors such as imatinib and nilotinib bind to the BCR-ABL1 fusion protein. They block binding of adenosine triphosphate (ATP) and thus prevent the tyrosine kinase from phosphorylating substrate proteins leading to apoptosis of the cell (see Fig. 14.4). They are used in CML and BCR-ABL+ ALL.

Inhibitors of B cell signalling pathways Ibrutinib, which blocks Bruton kinase (BTK), or idelalisib which inhibits PI3Kdelta, are valuable in the management of a range of B cell disorders (Fig. 9.4).

A wide range of additional kinase inhibitors is being introduced into therapy, such as JAK2 inhibitors, effective in primary myelofibrosis and polycythaemia vera, crizotinib, which blocks ALK activity, and inhibitors of the FLT3 kinase for treatment of AML.

Bortezomib and **carfilzomib** are proteasome inhibitors used widely in the treatment of myeloma and some lymphomas.

Monoclonal antibodies are highly effective and are particularly well established against B-cell malignancies. Rituximab binds to CD20 on B cells and mediates cell death, primarily through direct induction of apoptosis and opsonization (see p. 221). Other anti-CD20 and CD22 antibodies are available. Alemtuzumab binds to CD52 and is highly efficient at fixing complement which lyses the target B and T cells. Anti-CD30 (Brentuximab) is effective in Hodgkin lymphoma. Antibodies may also carry attached toxins (e.g. Mylotarg®, anti-CD33) or radioactive isotopes (e.g. Zevalin®, anti-CD20).

Other agents

Corticosteroids have a potent lymphocytotoxic activity and have an important role in many chemotherapeutic regimens used in the treatment of lymphoid malignancy and myeloma.

All-trans retinoic acid (ATRA) is a vitamin A derivative that acts as a differentiation agent in acute promyelocytic leukaemia (APML). Tumour cells in APML are arrested at the promyelocyte stage as a result of transcriptional repression resulting from the PML-RARA fusion protein (see p. 152). ATRA relieves this block and may lead to a brisk neutrophilia within a few days of treatment, with other side-effects known as the 'ATRA' or 'differentiation' syndrome (see p. 153).

Demethylation agents (e.g. azacytidine, decitabine) act to increase transcription by reducing methylation on cytosine residues within DNA.

Interferon- α is an antiviral and antimitotic substance produced in response to viral infection and inflammation. It has proven useful in chronic myeloid leukaemia, myeloma and myeloproliferative diseases.

Immunomodulatory drugs include thalidomide, lenalidomide and pomalidomide. They are effective in myeloma and in some types of myelodysplasia.

Asparaginase is an enzyme derived from bacteria that breaks down the amino acid asparagine within the circulation. ALL cells lack asparagine synthase and thus need a supply of exogenous asparagine for protein synthesis. Intramuscular asparaginase is an important agent in the treatment of ALL, although hypersensitivity reactions are not uncommon and blood clotting may be disturbed.

Platinum derivatives (e.g. cisplatin) are used in combinations for treating lymphoma.


Arsenic is useful in treatment of acute promyelocytic leukaemia. It induces differentiation and apoptosis.

SUMMARY

- Progress in the treatment of haemopoietic malignancies has been the result of improvements in both supportive therapy and specific tumour treatments.
- Initial assessment includes a performance score and tests for co-morbidities.
- Supportive treatments often include:
 - insertion of a central venous catheter;
 - appropriate use of red cell and platelet transfusions;
 - early administration of drugs to treat infection;
 - optimization of the blood coagulation system;
 - drugs to reduce side effects such as nausea or pain;
 - psychological support.
- Gram-positive skin organisms such as *Staphylococcus* are common infections and often colonize central venous catheters.
- Gram-negative bacteria are usually derived from the gut and can cause severe septicaemia.
- The use of air filters, handwashing and prophylactic antibiotics can reduce infection rates.
- Neutropenic patients who develop a fever should be treated urgently with broad-spectrum antibiotics.
- Herpes viruses are a common cause of infection in patients who are significantly immunosuppressed.
- Fungal infections are a major clinical problem for patients undergoing chemotherapy. Oral and intravenous antifungal drugs may be used for either prevention or treatment of disease.
- A wide range of drugs is now available for the treatment of haemopoietic malignancy:
 - alkylating agents;
 - antimetabolites;
 - anthracyclines;
 - signal transduction inhibitors, including tyrosine kinase inhibitors;
 - monoclonal antibodies;
 - immune modulators;
 - proteasome inhibitors;
 - others, e.g. corticosteroids, ATRA, demethylating agents, interferon, asparaginase, arsenic, platinum derivatives.



Now visit www.wileyessential.com/haematology to test yourself on this chapter.


A vertical strip on the left side of the page showing a microscopic view of blood cells. It features several large, bright red erythrocytes and some smaller, pinkish leukocytes against a dark blue background.

CHAPTER 13

Acute myeloid leukaemia

Key topics

■ Classification of leukaemia	146
■ Diagnosis of acute leukaemia	146
■ Acute myeloid leukaemia	147
■ Classification	147
■ Clinical features	148
■ Investigations	148
■ Treatment	149
■ Outcome	154



The leukaemias are a group of disorders characterized by the accumulation of malignant white cells in the bone marrow and blood. These abnormal cells cause symptoms because of: (i) bone marrow failure (e.g. anaemia, neutropenia, thrombocytopenia); and (ii) infiltration of organs (e.g. liver, spleen, lymph nodes, meninges, brain, skin or testes).

Classification of leukaemia

The main classification is into four types: acute and chronic leukaemias, which are further subdivided into lymphoid or myeloid.

Acute leukaemias are usually aggressive diseases in which malignant transformation occurs in the haemopoietic stem cell or early progenitors. Genetic damage is believed to involve several key biochemical steps resulting in (i) an increased rate of proliferation, (ii) reduced apoptosis and (iii) a block in cellular differentiation. Together these events cause accumulation in the bone marrow of early haemopoietic cells known as **blast cells**. The dominant clinical feature of acute leukaemia is usually bone marrow failure caused by accumulation of blast cells, although organ infiltration also occurs. If untreated, acute leukaemias are usually rapidly fatal, although with modern treatments most younger patients are ultimately cured of their disease.

Diagnosis of acute leukaemia

Acute leukaemia is normally defined as the presence of over 20% of blast cells in the bone marrow at clinical presentation. However, it can be diagnosed with less than 20% blasts if specific leukaemia-associated cytogenetic or molecular genetic abnormalities are present (Table 13.1; see Table 17.1).

The **lineage of the blast cells** is defined by microscopic examination (morphology) (Fig. 13.4; see Fig. 17.3), immunophenotypic (flow cytometry) (Fig. 13.1), cytogenetic and

Table 13.1 Classification of acute myeloid leukaemia (AML) according to the WHO classification 2008 (modified; see also Appendix).

<i>Acute myeloid leukaemia with recurrent genetic abnormalities</i>
AML with t(8;21)
AML with inv(16)
AML with t(15;17)(q22;q12); <i>PML-RARA</i>
<i>Acute myeloid leukaemia with myelodysplasia-related changes</i>
<i>Therapy-related myeloid neoplasms (t-AML)</i>
<i>Acute myeloid leukaemia, not otherwise specified</i>
AML with minimal differentiation
AML without differentiation
AML with maturation
Acute myelomonocytic leukaemia
Acute monoblastic/monocytic leukaemia
Acute erythroid leukaemia
Acute megakaryoblastic leukaemia
Acute basophilic leukaemia
Acute panmyelosis with myelofibrosis
<i>Myeloid sarcoma</i>
<i>Myeloid proliferations related to Down syndrome</i>
Transient abnormal myelopoiesis
Myeloid leukaemia

molecular analysis (Table 13.4 and see Table 11.1). This will define whether the blasts are of myeloid or lymphoid lineage and also localize the stage of cellular differentiation (Table 13.2). The typical 'myeloid immunophenotype' is CD13, CD33⁺ and TdT⁻ (Table 13.2; Fig. 13.1). Special antibodies are helpful in the diagnosis of the rare undifferentiated, erythroid or megakaryoblastic subtypes (Table 13.2).

Cytogenetic and **molecular** analysis is essential and is usually performed on marrow cells, although blood may be




Figure 13.1 Development of three cell lineages from pluripotential stem cells giving rise to the three main immunological subclasses of acute leukaemia. CD34 is expressed on most stem cells whereas TdT expression defines a lymphoid lineage. Three surface markers which are seen characteristically in T-ALL, B-ALL and AML are shown. AML, acute myeloid leukaemia; B-ALL, B-cell acute lymphoblastic leukaemia; c, cytoplasmic; HLA, human leucocyte antigen; T-ALL, T-cell acute lymphoblastic leukaemia; TdT, terminal deoxynucleotidyl transferase.

Table 13.2 Specialized tests for acute myeloid leukaemia.*Immunological markers (flow cytometry)*

CD13, CD33, CD117	+
CD11c, 14, 64	+ (monocytic)
Glycophorin (CD235a)	+ (erythroid)
Platelet antigens (e.g. CD41, CD42, CD61)	+ (megakaryoblastic)
Myeloperoxidase	+ (undifferentiated)

*Chromosome and genetic analysis (see Tables 13.1 & 13.4)**Cytochemistry*

Myeloperoxidase	+ (including Auer rods)
Sudan black	+ (including Auer rods)
Non-specific esterase	+ in M ₄ , M ₅

used if the blast cell count is particularly high. Cytochemistry can be useful in determining the blast cell lineage (see Fig. 13.6) but is no longer performed in centres where the newer and more definitive tests are available.

Acute myeloid leukaemia

Pathogenesis

The AML genome contains an average of about 10 mutations within protein-coding genes, amongst the smallest number of any adult cancer (see Fig. 11.3). Many AML 'driver mutations' have been identified, with the most common being within *FLT3*, *NPM1* and *DNMT3A* (Fig. 13.2). Some other mutations, e.g. of *ASXL1*, are frequent in myelodysplasia and when found in AML suggest that it is secondary to myelodysplasia. The mutations usually occur on only one of the two alleles for the gene and may be 'loss of function' or 'gain of function'. The average AML at presentation contains less than one gene-fusion event, which usually arise from translocations, with the most common being *PML-RARA*, *CBFB-MYH11*, *RUNX1-RUNX1T1* (Table 11.1) which are found in around 15%, 12% and 8% of cases respectively. The wide variety of cytogenetic abnormalities and molecular mutations are such that each case usually has a unique pattern of mutations.

Incidence

Acute myeloid leukaemia (AML) is the most common form of acute leukaemia in adults and becomes increasingly common with age, with a median onset of 65 years. It forms only a minor fraction (10–15%) of the leukaemias in childhood. Cytogenetic abnormalities and response to initial treatment have a major influence on prognosis (see Table 13.4).

Classification

AML is classified according to the World Health Organization (2008) scheme. There is an increasing focus on the genetic




Figure 13.2 The most commonly mutated genes within an analysis of 200 cases of acute myeloid leukaemia. Source: Adapted from The Cancer Genome Atlas Research Network. *NEJM* (2013) 368 (22): 2059–74.

abnormalities within the malignant cells and it is likely that ultimately almost all AML cases will be classified by specific genetic subtype. Currently this is not possible but many genetic subtypes have been determined. Approximately 60% of tumours exhibit karyotypic abnormalities on cytogenetic analysis and many cases with a normal karyotype carry mutations in genes such as nucleophosmin *FLT3*, (*NPM1*), *CEBPA*, *DNMT3A* (see below) detected only by molecular methods and which have prognostic significance.

Six main groups of AML are recognized (Table 13.1) and these are discussed below.

- 1 AML with recurrent genetic abnormalities** encompasses subtypes with specific chromosomal translocations or gene mutations. The detection of these abnormalities defines the tumour as AML and so the diagnostic criteria for this subgroup are relaxed in that the bone marrow blast cell count does not need to exceed 20% in order to make a diagnosis. In general these disorders have a good prognosis.
- 2 AML with myelodysplasia-related changes.** In this group the AML is associated with microscopic features of dysplasia in at least 50% of cells in at least two lineages. The clinical outcome of these patients is impaired in relation to the first subgroup.
- 3 Therapy-related myeloid neoplasms(t-AML)** arise in patients who have been previously treated with drugs such as etoposide or alkylating agents. They commonly exhibit mutations in the *MLL* gene and the clinical response is usually poor.
- 4 AML, not otherwise specified.** This group is defined by the absence of cytogenetic abnormalities and com-

prises around 30% of all cases. Mutations in the *NPM1* and *FLT3* genes are more frequent in those with normal cytogenetics.

5 Myeloid sarcoma is rare but refers to a disease that resembles a solid tumour but is composed of myeloid blast cells.

6 Myeloid proliferations related to Down's syndrome. Children with Down's syndrome have a greatly increased risk of acute leukaemia. Two myeloid variants are recognized: (i) transient abnormal myelopoiesis in which there is a self-limiting leucocytosis; and (ii) AML.

Mixed phenotype acute leukaemia

These rare cases express two markers for both myeloid and lymphoid differentiation either on the same blast cells or on two different cell populations. They usually have a poor prognosis.

Clinical features

The clinical features of AML are dominated by the pattern of bone marrow failure caused by the accumulation of malignant

cells within marrow (Fig. 13.3). Infections are frequent, and anaemia and thrombocytopenia are often profound. A bleeding tendency caused by thrombocytopenia and disseminated intravascular coagulation (DIC) is characteristic of the promyelocytic variant of AML. Tumour cells can infiltrate a variety of tissues. Gum hypertrophy and infiltration (Fig. 13.4), skin involvement and CNS disease are characteristic of the myelomonocytic and monocytic subtypes.

Investigations

Table 13.3 lists the initial clinical and laboratory tests to be performed in newly diagnosed cases of AML and similar work-up is needed for all new haematological malignancies.

Haematological investigations reveal a normochromic normocytic anaemia with thrombocytopenia in most cases. The total white cell count is usually increased and blood film examination typically shows a variable numbers of blast cells. The bone marrow is hypercellular and typically contains many leukaemic blasts (Fig. 13.5). Blast cells are characterized by morphology, immunological (flow cytometric) cytogenetic and molecular genetic analysis for confirming the diagnosis,



Figure 13.3 (a) An orbital infection in a female patient (aged 68 years) with acute myeloid leukaemia and severe neutropenia (haemoglobin 83 g/L, white cells $15.3 \times 10^9/L$, blasts 96%, neutrophils 1%, platelets $30 \times 10^9/L$). (b) Acute myeloid leukaemia: top: plaque *Candida albicans* on soft palate; lower: plaque *Candida albicans* in the mouth, with lesion of herpes simplex on the upper lip. (c) Skin infection (*Pseudomonas aeruginosa*) in a female patient (aged 33 years) with acute lymphoblastic leukaemia receiving chemotherapy and with severe neutropenia (haemoglobin 101 g/L, white cells $0.7 \times 10^9/L$, neutrophils $<0.1 \times 10^9/L$, lymphocytes $0.6 \times 10^9/L$, platelets $20 \times 10^9/L$).



Figure 13.4 Monocytic acute myeloid leukaemia: the gums are swollen and haemorrhagic because of infiltration by leukaemic cells.

determining prognosis and developing a treatment plan (Table 13.4). Cytochemistry (Fig 13.6) is no longer used in most centres.

Tests for DIC are often positive in patients with the promyelocytic variant of AML (see below). Biochemical tests are

Table 13.3 The initial work up of a newly diagnosed patient with AML.

Assessment of medical history, examination and performance status (see Chapter 12)
Full blood count and differential
Bone marrow aspirate and trephine biopsy
Immunophenotyping of bone marrow (and/or blood if blast cells present)
Cytogenetic analysis by karyotype
Mutation analysis
Cytochemical analysis (performed in some countries instead of immunophenotyping)
Biochemistry (liver, renal, uric acid, calcium, LDH)
Coagulation
Pregnancy test
Information on oocyte or sperm storage
Assessment of eligibility for stem cell transplantation
Hepatitis B, C and HIV test
CXR with ECG and ECHO in older patients
CXR, chest X-ray; ECG, electrocardiography; ECHO, echocardiography; HIV, human immunodeficiency virus; LDH, lactate dehydrogenase.

performed as a baseline before treatment begins and may reveal raised uric acid or lactate dehydrogenase.

Cytogenetics and molecular genetics

Genetic abnormalities are used to classify the majority of cases of AML (Table 13.1). Two of the most common $t(8;21)$ and $inv(16)$ are associated with a good prognosis. **Acute promyelocytic leukaemia** is a variant of AML that contains the $t(15;17)$ translocation in which the gene *PML* on chromosome 15 is fused to the retinoic acid receptor α gene, *RAR α* , on chromosome 17 (Fig. 13.7). The resultant PML-RAR α fusion protein functions as a transcriptional repressor whereas normal (wild-type) *RAR α* is an activator. Normally, the PML protein forms homodimers with itself, whereas the RAR α protein forms heterodimers with the retinoid X receptor protein, RXR. The PML-RAR α fusion protein binds to PML and RXR, preventing them from linking with their natural partners. This results in the cellular phenotype of arrested differentiation.

Point mutations affecting the genes *FLT3*, *NPM1*, *DNMT3A*, *CEBPA*, *TET2*, *WT1*, *IDH1*, *RUNX1*, and others are frequent in AML, especially in those cases without a cytogenetic abnormality (Fig. 13.2). They may be used to subclassify the disease (see Table 11.1) and have prognostic significance (Table 13.4). Some of these genes are involved in DNA methylation or acetylation (see Fig 16.1) and are also mutated in cases of myelodysplasia and myeloproliferative diseases (see Chapters 15 and 16). The presence in *de novo* AML of a myelodysplasia 'specific' mutation, e.g. of *ASXL1* or *SF3B1* is unfavourable.

Treatment

Management is both supportive and specific.

- 1 General supportive therapy** for bone marrow failure is described in Chapter 12 and includes the insertion of a central venous cannula, blood product support and prevention of tumour lysis syndrome. The platelet count is generally maintained above $10 \times 10^9/L$ and the haemoglobin above 80 g/L. Any episode of fever must be treated promptly. Acute promyelocytic leukaemia needs special support as described below.
- 2 The aim of treatment** in acute leukaemia is to induce complete remission (less than 5% blasts in the bone marrow, normal blood counts and clinical status) and then to consolidate this with intensive therapy, hopefully eliminating the disease (Fig. 13.8). Allogeneic stem cell transplantation is considered in poor prognosis cases (Table 13.4) or for patients who have relapsed.
- 3 Specific therapy of AML** is determined by the age and performance status of the patient as well as the genetic lesions within the tumour. In younger patients treatment is primarily with the use of intensive chemotherapy. This is usually given in three or four blocks, each




Figure 13.5 Morphological examples of acute myeloid leukaemia. (a) Blast cells without differentiation show few granules but may show Auer rods, as in this case; (b) cells in differentiation show multiple cytoplasmic granules or (c) M_3 blast cells contain prominent granules or multiple Auer rods; (d) myelomonocytic blasts have some monocytoïd differentiation; (e) monoblastic leukaemia in which >80% of blasts are monoblasts; (f) monocytic with <80% of blasts monoblasts; (g) erythroid showing preponderance of erythroblasts; (h) megakaryoblastic showing cytoplasmic blebs on blasts.




Figure 13.6 Cytochemical staining in acute myeloid leukaemia. **(a)** Sudan black B shows black staining in the cytoplasm. **(b)** Myelomonocytic: non-specific esterase/chloracetate staining shows orange-staining monoblast cytoplasm and blue-staining (myeloblast) cytoplasm.

of approximately 1 week, and the most commonly used drugs are cytosine arabinoside and daunorubicin (both in conventional or high doses). Idarubicin, mitoxantrone and etoposide are also used in various regimens (Figs 13.8 and 13.9).

A typical good response in AML is shown in Figure 13.10. The drugs are myelotoxic with limited selectivity between

leukaemic and normal marrow cells, so marrow failure resulting from the chemotherapy is severe and prolonged, and intensive supportive care is required. Maintenance therapy is of no value except in promyelocytic AML with ATRA. CNS prophylaxis is not usually given. New drugs such as FLT3 inhibitors are now being introduced for tumours with *FLT3* mutations. Monoclonal immunoconjugates targeted against

Table 13.4 Prognostic factors in acute myeloid leukaemia (AML).

	Favourable	Intermediate	Unfavourable
Cytogenetics	t(15;17) t(8;21) inv(16)	Normal Other non-complex changes	Deletions of chromosome 5 or 7, TP53 mutated Complex rearrangements (>3 unrelated abnormalities)
Molecular genetics	<i>NPM</i> mutation <i>CEBPA</i> mutation	Wild type	<i>FLT3</i> internal tandem repeat
Bone marrow response to remission induction	<5% blasts after first course		>20% blasts after first course
Age	Child	<60 years	>60 years
Performance status	Good		Bad
Co-morbidities	Absent		Present
White cell count	<10 × 10 ⁹ /L		>100 × 10 ⁹ /L
Secondary leukaemia	Absent		Present, e.g. to previous chemotherapy or marrow disease
Minimal residual disease in remission	Absent		Present (>0.1% of cells)




Figure 13.7 Generation of the t(15;17) translocation. The *PML* gene at 15q22 may break at one of three different breakpoint cluster regions (BCR-1, -2 and -3) and joins with exons 3–9 of the *RARα* gene at 17q12. Three different fusion mRNAs are generated (termed long (L), variable (V) or short (S)) and these give rise to fusion proteins of different size. In this diagram only the long version resulting from a break at BCR-1 is shown.

CD33 (e.g. Mylotarg®) or CD45 provide an additional therapeutic option for initial or consolidation AML therapy.

Acute promyelocytic leukaemia (APML) has its own treatment protocol. An haemorrhagic syndrome can lead to catastrophic haemorrhage and may be present either at

diagnosis or develop in the first few days of treatment. It is treated as for disseminated intravascular coagulation (DIC) with multiple platelet transfusions and replacement of clotting factors with fresh frozen plasma (see p. 358). In addition, all-*trans* retinoic acid (ATRA) therapy is given for this




Figure 13.8 Acute leukaemia: principles of therapy for AML or ALL (acute lymphoblastic leukaemia); SCT, stem cell transplantation; TBI, total body irradiation. The decision for SCT in remission is based on prognostic factors as well as tests for minimal residual disease.




Figure 13.9 Acute myeloid leukaemia: flow chart illustrating typical treatment regimen.

disease subtype and is combined initially with either arsenic trioxide or anthracycline. The arsenic combination appears to give a better clinical response with fewer side-effects. The **differentiation syndrome** (also known as ATRA syndrome) is a specific complication that may arise after ATRA treatment. Clinical problems, which result from the neutrophilia that follows differentiation of promyelocytes, include fever, hypoxia with pulmonary infiltrates and fluid overload. Treatment is with steroids and ATRA is only discontinued in very severe cases.

Prognosis and treatment stratification

The outcome for an individual patient with AML will depend on a number of factors including age and white cell count at presentation (Table 13.4). However, the genetic abnormalities in the tumour are the most important determinant.

Complete remission is defined as less than 5% blasts without Auer rods, neutrophil count greater than $1.0 \times 10^9/L$, platelets greater than $100 \times 10^9/L$, independence of red cell transfusions and no extramedullary disease.

An important development in AML therapy is that of basing treatment according to the individual patient's **risk group**. Favourable cytogenetics and remission after one course of chemotherapy both predict for a better prognosis. In contrast, monosomy 5 or 7 abnormalities, blast cells with the *FLT3* internal tandem duplication mutation or poorly responsive disease places patients into poor risk groups which need more intensive treatments (Table 13.4).

Monitoring of **minimal residual disease** during and after chemotherapy, for example detection of a positive *FLT3* mutation test in otherwise complete remission, is being investigated

as a means to guide appropriate treatment. It may be performed by molecular or cytogenetic tests or flow cytometry of the abnormal 'leukaemia-associated immunophenotype' seen in over 90% of cases.

Stem cell transplantation

Allogeneic stem cell transplantation (SCT) reduces the rate of AML relapse and is offered in selected intermediate- and high-risk cases in first remission. It carries risk of morbidity and mortality so is not used for patients in the favourable risk group unless they have disease relapse. Reduced-intensity conditioning regimens have raised the age at which patients may be considered for SCT. Potential donors are discussed in Chapter 23. Autologous transplantation confers no benefit above that of post remission chemotherapy.

Patients over 70 years of age

The median age for presentation of AML is approximately 65 years and treatment outcomes in the elderly are poor because of primary disease resistance and poor tolerability of intensive treatment protocols. Death from haemorrhage, infection or failure of the heart, kidneys or other organs is more frequent than in younger patients. In elderly patients with serious disease of other organs, the decision may be made to use supportive care with or without gentle single-drug chemotherapy, e.g. with low-dose cytarabine, azacytidine or hydroxycarbamide. However, in those otherwise well, combination chemotherapy similar to that used in younger patients may produce long-term remissions and reduced-intensity SCT is increasingly being offered.




Figure 13.10 Typical flow chart for the management with chemotherapy of acute myeloid leukaemia. WBC, white blood cells.

Treatment of relapse

Most patients suffer relapse and the outlook will then depend on age, the duration of the first remission and the cytogenetic risk group. In addition to further chemotherapy, allogeneic SCT with either standard or reduced-intensity conditioning is usually performed in those patients who can tolerate the procedure and who have a suitable donor. Arsenic trioxide is useful in management of relapse in the promyelocytic variant.

Outcome

The prognosis for patients with AML has been improving steadily, particularly for those under 60 years of age, and approximately one-third of this group can expect to achieve long-term cure (Fig. 13.11a). For the elderly the situation is poor and less than 10% of those over 70 years of age achieve long-term remission (Fig. 13.11b).





Figure 13.11 Improvement in survival over time for patients with AML in UK MRC AML Trials (a) and in relation to age 0–14 years (b), age 15–59 years (c) and age 60+ years (d). Source: Courtesy of Dr Robert Hills, University of Cardiff.

- The leukaemias are a group of disorders characterized by the accumulation of malignant white cells in the bone marrow and blood. They can be classified into four subtypes on the basis of being either *acute* or *chronic*, and *myeloid* or *lymphoid*.
- Acute leukaemias are aggressive diseases in which transformation of a haemopoietic stem cell leads to accumulation of >20% blast cells in the bone marrow.
- The clinical features of acute leukaemia result from bone marrow failure and include anaemia, infection and bleeding. Tissue infiltration can also occur.
- AML is rare in childhood but becomes increasingly common with age with a median onset of 65 years.
- The diagnosis is made by analysis of blood and bone marrow using microscopic examination (morphology) as well as immunophenotypic, cytogenetic and molecular studies.
- Cytogenetic and molecular abnormalities are used to classify and indicate prognosis in the majority of cases of AML.
- In younger patients treatment is primarily with the use of intensive chemotherapy. This is usually given in three or four blocks, each of approximately 1 week, using drugs such as cytosine arabinoside and daunorubicin. The blood count must be given time to recover before the next block of therapy can be given and this can typically take 4–6 weeks.
- Acute promyelocytic leukaemia is a variant of AML that carries a t(15;17) chromosomal translocation. It commonly presents with bleeding and is treated with retinoic acid (ATRA) and arsenic or chemotherapy.
- The prognosis for patients with AML has been improving steadily, particularly for those under 60 years of age, and approximately one-third of this group can expect to achieve long-term cure. The outcome for elderly people remains disappointing.
- Allogeneic stem cell transplantation is useful in treating some subsets of patients and may also be curative for patients with relapsed disease.



Now visit www.wileyessential.com/haematology to test yourself on this chapter.

A vertical strip on the left side of the page showing a microscopic view of various blood cells, including red blood cells and white blood cells, in shades of red, pink, and yellow against a dark blue background.

CHAPTER 14

Chronic myeloid leukaemia

Key topics

■ Chronic myeloid leukaemia	157
■ Clinical features	159
■ Laboratory findings	159
■ Treatment	160
■ Accelerated phase disease and blastic transformation	162
■ Chronic neutrophilic leukaemia	164
■ Chronic eosinophilic leukaemia	164




Table 14.1 Chronic myeloid leukaemia (CML) and myelodysplastic/myeloproliferative neoplasms (see Chapter 16; see also Appendix).

Type
<i>BCR-ABL1</i> rearrangement positive CML
<i>BCR-ABL1</i> rearrangement negative CML
Chronic neutrophilic leukaemia
Chronic eosinophilic leukaemia
Chronic monocytic leukaemia
Chronic myelomonocytic leukaemia
Juvenile myelomonocytic leukaemia
Refractory anaemia with ringed sideroblasts and thrombocytosis
NB All the diseases except <i>BCR-ABL1</i> positive CML are rare.

The chronic leukaemias are distinguished from acute leukaemias by their slower progression. Chronic leukaemias can be broadly subdivided into myeloid (Table 14.1) and lymphoid groups (see Chapter 18).

Chronic myeloid leukaemia

Chronic myeloid leukaemia *BCR-ABL1*+ (CML) is a clonal disorder of a pluripotent stem cell. The disease accounts for around 15% of leukaemias and may occur at any age. The diagnosis of CML is rarely difficult and is assisted by the characteristic presence of the **Philadelphia (Ph) chromosome**. This results from the t(9;22) (q34;q11) translocation between chromosomes 9 and 22, as a result of which part of the onco-gene *ABL1* is moved to the *BCR* gene on chromosome 22 (Fig. 14.1a) and part of chromosome 22 moves to chromosome 9. The abnormal chromosome 22 is the Ph chromosome. In the Ph translocation 5' exons of *BCR* are fused to the 3' exons of *ABL1* (Fig. 14.1b, c). The resulting chimeric *BCR-ABL1* gene codes for a fusion protein of size 210 kDa (p210). This has tyrosine kinase activity in excess of the normal 145-kDa *ABL1* product. The Ph translocation is also seen in a minority of cases of acute lymphoblastic leukaemia (ALL) and in some of these the breakpoint in *BCR* occurs in the same region as in CML. However, in other cases the breakpoint in *BCR* is further upstream, in the intron between the first and second exons, leaving only the first *BCR* exon intact. This chimeric *BCR-ABL1* gene is expressed as a p190 protein which, like p210, has enhanced tyrosine kinase activity.





Figure 14.1 The Philadelphia chromosome. **(a)** There is translocation of part of the long arm of chromosome 22 to the long arm of chromosome 9 and reciprocal translocation of part of the long arm of chromosome 9 to chromosome 22 (the Philadelphia chromosome). This reciprocal translocation brings most of the *ABL1*-gene into the *BCR* region on chromosome 22 (and part of the *BCR* gene into juxtaposition with the remaining portion of *ABL* on chromosome 9). **(b)** The breakpoint in *ABL1* is between exons 1 and 2. The breakpoint in *BCR* is at one of the two points in the major breakpoint cluster region (M-BCR) in chronic myeloid leukaemia (CML) or in some cases of Ph+ acute lymphoblastic leukaemia (ALL). **(c)** This results in a 210-kDa fusion protein product derived from the *BCR-ABL1* fusion gene. In other cases of Ph+ ALL, the breakpoint in *BCR* is at a minor breakpoint cluster region (m-BCR) resulting in a smaller *BCR-ABL1* fusion gene and a 190-kDa protein.

(Continued)




(d)



(i)

(e)



(ii)

Figure 14.1 (Continued) (d) Karyotype showing the t(9;22) (q34;q11) translocation. The Ph chromosome is arrowed. (e) Visualization of the Ph chromosome on: (i) dividing (metaphase); and (ii) quiescent (interphase) cells by fluorescence *in situ* hybridization (FISH) analysis (ABL probe in red and BCR probe in green) with fusion signals (red/green which shows as yellow) on the Ph (BCR-ABL1) and der(9) (ABL1-BCR) chromosomes. Source: Courtesy of Dr Ellie Nacheva.

In most patients the Ph chromosome is seen by karyotypic examination of tumour cells (Fig. 14.1d), but in a few the Ph abnormality cannot be seen under the microscope but the same molecular rearrangement is detectable by more sensitive techniques: fluorescence *in situ* hybridization (FISH)

(Fig. 14.1e) or by reverse transcriptase PCR (RT-PCR) for BCR-ABL1 transcripts. This Ph-negative *BCR-ABL1*-positive CML behaves clinically like Ph-positive CML. As the Ph chromosome is an acquired abnormality of haemopoietic stem cells it is found in cells of both the myeloid (granulocytic, erythroid

and megakaryocytic) and lymphoid (B and T cell) lineages. The main cause of death in CML is transformation to a blast phase, which may be preceded by an accelerated phase. These are discussed later in this chapter.

BCR-ABL1-negative chronic myeloid leukaemia is classified with the myelodysplastic/myeloproliferative syndromes (see Chapter 16).

Clinical features

This disease occurs in either sex (male:female ratio of 1.4:1), most frequently between the ages of 40 and 60 years. However, it may occur in children and neonates, and in the very old. In up to 50% of cases the diagnosis is made incidentally from a routine blood count. In those cases where the disease presents clinically, the following features may be seen:

- 1 Symptoms related to hypermetabolism (e.g. weight loss, lassitude, anorexia or night sweats).
- 2 Splenomegaly is nearly always present and may be massive. In some patients splenic enlargement is associated with considerable discomfort, pain or indigestion.
- 3 Features of anaemia may include pallor, dyspnoea and tachycardia.
- 4 Bruising, epistaxis, menorrhagia or haemorrhage from other sites because of abnormal platelet function.
- 5 Gout or renal impairment caused by hyperuricaemia from excessive purine breakdown may be a problem.
- 6 Rare symptoms include visual disturbances and priapism.

Laboratory findings

- 1 Leucocytosis is the main feature and may reach levels greater than $200 \times 10^9/\text{L}$ (Fig. 14.2). A complete spectrum of myeloid cells is seen in the peripheral blood. The levels of neutrophils and myelocytes exceed those of blast cells and promyelocytes (Fig. 14.3).
- 2 Increased circulating basophils are a characteristic feature.
- 3 Normochromic normocytic anaemia is usual.
- 4 Platelet count may be increased (most frequently), normal or decreased.
- 5 Bone marrow is hypercellular with granulopoietic predominance.
- 6 Presence of the *BCR-ABL1* gene fusion by RT-PCR analysis and in 98% of cases Ph chromosome on cytogenetic analysis (Fig. 14.1d).
- 7 Serum uric acid is usually raised.

Prognostic scores (stages)

Attempts have been made to stage CML at presentation in order to predict prognosis. The most frequently used is the Sokal score, which takes account of age, blast cell percentage, spleen size and platelet count. However the rate of response to a tyrosine kinase inhibitor is now a more useful measure.



Figure 14.2 Chronic myeloid leukaemia: peripheral blood film showing a vast increase in buffy coat. The white cell count was $532 \times 10^9/\text{L}$.




Figure 14.3 Chronic myeloid leukaemia: peripheral blood film showing various stages of granulopoiesis including promyelocytes, myelocytes, metamyelocytes and band and segmented neutrophils.

Table 14.2 The tyrosine kinase inhibitors used for treatment of CML.

	Action	Side-effects	Role in clinical therapy
Imatinib	The first TKI, designed as a specific inhibitor of the BCR-ABL1 fusion protein. It blocks tyrosine kinase activity by competing with adenosine triphosphate (ATP) binding (Fig. 14.4).	Rash, nausea, myelosuppression, fluid retention, muscle cramps	First-line therapy. Occasional use in resistance/intolerance to other TKI
Nilotinib	Second-generation inhibitor that has increased affinity for BCR-ABL1. Achieves molecular remission more rapidly than imatinib	Myelosuppression, headache, nausea, prolonged QT interval	First-line or resistance/intolerance to first line TKI
Dasatinib	Inhibits BCR-ABL and SRC family kinases, which play a role in driving CML progression	Headache, pleural effusion, cough, prolonged QT interval	First-line or resistance/intolerance to first-line TKI.
Bosutinib	BCR-ABL1 inhibitor with additional inhibitory effect on SRC family kinases	Diarrhoea, nausea, low platelet count	Resistance/intolerance to first-line TKI
Ponatinib	Multi-targeted TKI inhibitor	Low platelets, rash, dry skin, arterial thrombosis	Resistance/intolerance to first-line TKI. Able to treat CML with the T315I mutation

Treatment

Treatment of chronic phase

Tyrosine kinase inhibitors

Tyrosine kinase inhibitors (TKI) are the mainstay of the treatment of CML and several different drugs are now available (Table 14.2).

First-line therapy for patients with chronic phase CML is usually imatinib, nilotinib or dasatinib. Most experience exists with imatinib (Fig. 14.4), which is also the cheapest. Overall around 60% of patients given imatinib achieve an excellent response, whereas 40% proceed to a second-line agent due to intolerance or inadequate response.

Nilotinib and dasatinib achieve more rapid responses with first-line therapy and are therefore used in some centres, although side-effects are somewhat more common. They are also used as second-line therapy after imatinib (Fig. 14.5). Bosutinib is also an effective second line therapy, whilst ponatinib has the unique advantage that it is effective against tumours that carry the T315I mutation within BCR-ABL1.

Monitoring of response to tyrosine kinase inhibitors (TKI)

TKI are highly effective at reducing the number of tumour cells in the bone marrow and should be monitored by RT-PCR analysis for *BCR-ABL1* transcripts in marrow or blood and/or karyotypic analysis of the bone marrow, typically at 3, 6 and 12 months. Molecular response is assessed as the ratio of

BCR-ABL1 transcripts to *ABL1* transcripts and it is expressed as *BCR-ABL1*% on a log scale, where 10%, 1%, 0.1%, 0.01%, and 0.001% correspond to a decrease of 1, 2, 3, 4, and 5 logs, respectively, below the standard baseline. A complete cytogenetic response (CCyR) is defined as the absence of Ph-positive metaphases within bone marrow.




Figure 14.4 Mode of action of the tyrosine kinase inhibitor imatinib. It blocks the enzyme's adenosine triphosphate (ATP) binding site.




Figure 14.5 Example of an optimal haematological and cytogenetic response in a patient with CML who achieves complete remission with imatinib therapy. **(a)** The white cell count returns to normal within days. **(b)** Karyotypic examination of the bone marrow reveals a gradual reduction in the number of Ph chromosomes over the 6 months. **(c)** PCR analysis of the bone marrow or blood shows a reduction in the number of BCR-ABL1 transcripts in comparison with the normal ABL1 transcript. BCR-ABL1 transcripts continue to be detected at a very low level but can become negative in some patients. In this case analysis was performed on bone marrow for the first 6 months and on peripheral blood thereafter.

Treatment responses can be defined as *Optimal* or *Failure*, with an intermediate area termed *Warning* (Table 14.3). Patients with optimal responses continue their original treatment whereas those with treatment failure are treated with second-generation TKI therapy or stem cell transplantation (SCT). Patients in the Warning zone should be monitored more regularly and might be considered for an early change in therapy or increase in the dose of imatinib.

BCR-ABL1 mutation screening

One mechanism of disease resistance to TKI therapy is selection for clones carrying mutations within the *BCR-ABL1* fusion gene e.g. T315I. These mutations may be detected by sequencing the *BCR-ABL1* gene. The pattern of mutation can be useful for determining which treatment to choose as second-line therapy.

Response to TKI therapy

TKI therapy is highly effective and after 5 years of treatment the progression-free survival is 85–90%, with overall survival over 90% (Fig. 14.6). For patients becoming BCR-ABL1 transcript negative, about 60% will remain negative on stopping TKI therapy or remain in remission with a stable low level of the transcripts. For those who do become BCR-ABL1 positive again with a rising level of transcripts, treatment with the TKI will usually return them to BCR-ABL1 negative status. It is therefore possible that TKI can cure some patients with CML.

Additional forms of treatment for CML

Chemotherapy

Hydroxycarbamide (hydroxyurea) treatment can control and maintain the white cell count in the chronic phase but does not

Table 14.3 Definition of the response to TKIs as first-line treatment in CML. As examples, BCR-ABL1 $\leq 10\%$ refers to the fact that the *BCR-ABL1* transcript level has been reduced to less than 10% of the *ABL1* level. Ph+ $\leq 35\%$ indicates that the number of Philadelphia-positive chromosomes in cells in the bone marrow is less than 35% of the total number that are examined. Source: From Baccarani *et al.* (2013) *Blood*, 122: 872–884.

	Optimal	Warning	Failure
3 months	BCR-ABL1 $\leq 10\%$ and/or Ph+ $\leq 35\%$	BCR-ABL1 $> 10\%$ and/or Ph+ 36–95%	Non-CHR and/or Ph+ $> 95\%$
6 months	BCR-ABL1 $< 1\%$ and/or Ph+ 0	BCR-ABL1 1–10% and/or Ph+ 1–35%	BCR-ABL1 $> 10\%$ and/or Ph+ $> 35\%$
12 months	BCR-ABL1 $\leq 0.1\%$	BCR-ABL1 > 0.1 –1%	BCR-ABL1 $> 1\%$ and/or Ph+ > 0
Then, and at any time	BCR-ABL1 $\leq 0.1\%$	CCA/Ph– (–7, or 7q–)	Loss of complete haematological response Loss of complete cytogenetic response Loss of major molecular response (BCR-ABL1 expression of $\leq 0.1\%$) Mutations Clonal chromosome abnormalities in Ph+ cells

CCA, complex cytogenetic abnormalities; CHR, complete haematological response.

reduce the percentage of *BCR-ABL1*-positive cells. TKIs have now largely replaced its use.

α -Interferon

This was often used after the white cell count had been controlled by hydroxycarbamide but has now been replaced by

imatinib. Almost all patients have symptoms of a ‘flu-like’ illness in the first few days of treatment which responds to paracetamol and gradually wears off. More serious complications include anorexia, depression and cytopenias (see Table 12.2). A minority (approximately 15%) of patients may achieve long-term remission with loss of the Ph chromosome on cytogenetic analysis although the *BCR-ABL1* fusion gene can usually still be detected by PCR.

Stem cell transplantation

Allogeneic SCT is a potentially curative treatment for CML but because of the risks associated with the procedure, it is usually reserved for TKI failures (Fig. 14.7), or patients presenting in accelerated or acute phases. The results are, however, better when it is performed in chronic rather than these later phases. The 5-year survival is approximately 50–70%. Relapse of CML after the transplant is a significant problem but donor leucocyte infusions are highly effective in CML (see p. 261), particularly if relapse is diagnosed early by molecular detection of the *BCR-ABL1* transcript.

Accelerated phase disease and blastic transformation

Acute transformation (greater than 20% blasts in blood or marrow) may occur rapidly over days or weeks (Fig. 14.8).




Figure 14.6 Clinical outcome of patients given imatinib therapy for treatment of CML in chronic phase. After 7 years only 6% of patients died due to CML and overall survival is 86%. Source: O'Brien S.G. *et al.* (2008) *Blood* 112: 76a, with permission.




Figure 14.7 A potential algorithm for the management of young patients with CML in chronic phase (CML-CP). Imatinib is suggested as first line therapy. The role of stem cell transplantation (SCT) is here emphasized in the management of young patients. Source: Adapted from de la Fuente *et al.* (2014) *Br J Haematol*, 167(1): 33–47. 2G TKI, second generation tyrosine kinase inhibitor.

More commonly there is an **accelerated phase** with anaemia, thrombocytopenia (platelets less than $100 \times 10^9/L$), increase in blood basophils to greater than 20% or marrow blast cells 10–19% with blast cells in the blood. The spleen may be enlarged despite control of the blood count and the marrow may be fibrotic. New clonal chromosomal or molecular abnormalities may appear. The patient may be in this phase for several months, during which the disease is less easy to control than in the chronic phase. In approximately one-fifth of cases of acute transformation this is lymphoblastic and patients may be treated in a similar way to ALL, with a number of patients returning to the chronic phase for months or even a year or two. In the majority, transformation is into acute myeloid leukaemia or mixed types. These are more difficult to treat and survival is rare beyond 1 year. TKI are valuable in the management of blastic transformation but resistance to treatment usually occurs within a few weeks. Allogeneic SCT is a valuable option where possible.




Figure 14.8 Chronic myeloid leukaemia: acute myeloblastic transformation. Peripheral blood film showing frequent myeloblasts.

Chronic neutrophilic leukaemia

These patients have no inflammatory or other causes of neutrophilia and no evidence for any other myeloproliferative disease. They may have mild splenomegaly. Activating mutations in the gene encoding the receptor for colony stimulating factor 3 (*CSF3R*) are found in most patients and kinase inhibitors may be useful. The prognosis is variable.

Chronic eosinophilic leukaemia

Chronic eosinophilic leukaemia is a clonal persistent eosinophilia (greater than $1.5 \times 10^9/L$). There may be an interstitial lesion in chromosome 4 resulting in *FIP1L1-PDGFR* fusion


gene (in which case there is a response to tyrosine kinase inhibitors) or other less frequent cytogenetic or molecular defects. There may be greater than 5% but less than 20% blasts in the marrow. The cells may infiltrate various organs causing damage (e.g. endomyocardial fibrosis, lung, CNS, skin and gastrointestinal tract). If clonality cannot be shown and blasts are less than 5%, the condition is diagnosed as hypereosinophilic syndrome (see p. 96).

SUMMARY

- Chronic myeloid leukaemia is a clonal disorder of a pluripotent stem cell. The disease accounts for around 15% of leukaemias and may occur at any age.
- All cases of CML have a translocation between chromosomes 9 and 22. This leads to the oncogene *ABL1* being moved to the *BCR* gene on chromosome 22 and generates the Philadelphia (Ph) chromosome.
- The resulting chimeric *BCR-ABL1* gene codes for a fusion protein with increased tyrosine kinase activity.
- In most patients the Ph chromosome is seen by karyotypic examination of blood or bone marrow cells but the molecular rearrangement may rarely only be detected by FISH or PCR.
- The disease can occur at any age but is most common between the ages of 40 and 60 years.
- The clinical features include loss of weight, sweating, anaemia, bleeding and splenomegaly. There is usually a marked neutrophilia with myelocytes and basophils seen in the blood film.
- Transformation to an accelerated phase or acute leukaemia may occur.
- Treatment is with tyrosine kinase inhibitors such as imatinib, dasatinib or nilotinib. Tumour cells can acquire resistance to treatment and drug therapy is tailored in response to this.
- Stem cell transplantation can be curative and may also be useful for advanced disease.
- The clinical outlook is now very good and 90% of patients can expect long-term control of disease.
- Chronic eosinophilic and neutrophil leukaemias are much rarer.



Now visit www.wileyessential.com/haematology to test yourself on this chapter.


A vertical strip on the left side of the page showing a microscopic view of blood cells. It features several large, bright red erythrocytes and a few smaller, yellowish leukocytes against a dark blue background.

CHAPTER 15

Myeloproliferative disease

Key topics

■ Polycythaemia	168
■ Primary polycythaemia (erythrocytosis)	168
■ Polycythaemia vera	168
■ Secondary polycythaemia	172
■ Apparent polycythaemia	172
■ Differential diagnosis of polycythaemia	172
■ Essential thrombocythaemia	172
■ Primary myelofibrosis	173
■ Mastocytosis	175



The term myeloproliferative neoplasms (MPN) describes a group of conditions arising from marrow stem cells and characterized by clonal proliferation of one or more haemopoietic components in the bone marrow and, in many cases, the liver and spleen. They are often called the myeloproliferative diseases. The three major non-leukaemic disorders included in this classification are:

- 1 **Polycythaemia vera (PV);**
- 2 **Essential thrombocythaemia (ET); and**
- 3 **Primary myelofibrosis.**

Mastocytosis is also discussed in this chapter. *BCR-ABL1*-positive chronic myeloid leukaemia is described in Chapter 14 and the myelodysplastic syndromes and mixed myelodysplastic/myeloproliferative diseases in Chapter 16.

Myeloproliferative disorders are closely related to each other and transitional forms can occur with evolution from one entity into another during the course of the disease (Fig. 15.1). These diseases are associated with acquired mutations of the genes that encode the tyrosine kinase proteins, Janus-associated kinase 2 (**JAK2**), **MPL** (the receptor for thrombopoietin), or calreticulin (**CALR**) (Table 15.1). Mutation of *JAK2* (*JAK2V617F*) occurs in a heterozygous or homozygous state, in the marrow and blood of almost all patients with PV and in approximately 60% of those with ET and primary myelofibrosis, showing the common aetiology of these three diseases (Fig. 15.2). The mutation occurs in a highly conserved region of the pseudokinase domain, which is believed to negatively regulate *JAK2* signalling. *JAK2* has a major role in normal myeloid development by transducing signals from cytokines and growth factors including erythropoietin and thrombopoietin

(see Fig. 1.7). Why the same mutation is associated with different myeloproliferative diseases is unclear but depends partly on the dosage of the mutant allele, this being usually higher in PV than ET. A minority of PV patients show a variant *JAK2* mutation in exon 12. In those patients who do not demonstrate a *JAK2* mutation a mutation in the calreticulin (*CALR*) is observed in most cases. *CALR* is a multifunction protein involved in signal transduction and gene transcription. Mutations in the *MPL* gene are found in 5–10% of ET and MF cases. A mutation in one of these three genes is seen in 99% of cases of PV and around 85–90% of cases of ET or MF (Fig. 15.3).

Table 15.1 Genetic mutations in myeloproliferative diseases and other myeloid neoplasms.

Disease	Gene mutations
Chronic myeloid leukaemia	<i>ABL1</i>
Polycythaemia vera	<i>JAK2</i>
Primary myelofibrosis	<i>JAK2</i> , <i>CALR</i> , <i>MPL</i>
Essential thrombocythaemia	<i>JAK2</i> , <i>CALR</i> , <i>MPL</i>
Mastocytosis	<i>KIT</i>
Myeloid neoplasm with eosinophilia	<i>PDGFRA</i> , <i>PDGFRB</i> , <i>FGFR1</i>




Figure 15.1 Relationship between the three myeloproliferative diseases. They may all arise by somatic mutation in the pluripotent stem and progenitor cells. Many transitional cases occur showing features of two conditions and, in other cases, the disease transforms during its course from one of these diseases to another or to acute myeloid leukaemia. The three diseases, polycythaemia rubra vera, essential thrombocythaemia and primary myelofibrosis, are characterized by *JAK2* or *CALR* mutation in a varying proportion of cases.




Figure 15.2 The role of *JAK2* mutation in the generation of myeloproliferative diseases. **(a)** (i) Most haemopoietic growth factor receptors do not have intrinsic kinase activity but associate with a protein kinase such as *JAK2* in the cytoplasm. (ii) When the receptor binds a growth factor (e.g. erythropoietin) the cytoplasmic domains move closer together and the *JAK2* molecules can activate each other by phosphorylation and phosphorylate downstream proteins e.g. *STATS* (Fig. 1.7). (iii) The V617F *JAK2* mutation allows the *JAK* protein to become activated even when no growth factor is bound. **(b)** A model for the development of myeloproliferative disease following *JAK2* mutation. The primary event appears to predispose to an acquired heterozygous mutation of *JAK2* (V617F). This leads to a survival advantage. In some patients, a mitotic recombination event leads to a homozygous *JAK2* mutation state.




Figure 15.3 Frequency of *JAK2*, *CALR* and *MPL* mutations in the three subtypes of myeloproliferative neoplasm. Source: Adapted from Nangalia J & Green TR (2014) *Hematology*. American Society of Haematology Education Program 287–96.

Some patients also show mutations of genes, e.g. TET2, found in myelodysplasia and acute myeloid leukaemia (see Fig. 16.1), but their prognostic significance is not yet established.

There is a five-fold increased incidence of myeloproliferative disease in close relatives of patients, implying a genetic predisposition to the diseases.

Polycythaemia

Polycythaemia is defined as an increase in the haemoglobin concentration above the upper limit of normal for the patient's age and sex.

Classification of polycythaemia

Polycythaemia is classified according to its pathophysiology but the major subdivision is into **absolute polycythaemia** or erythrocytosis, in which the red cell mass (volume) is raised to greater than 125% of that expected for body mass and gender, and the much more frequent **relative** or **pseudopolycythaemia** in which the red cell volume is normal but the plasma volume is reduced. If the haematocrit is higher than 0.60 there will always be a raised red cell mass. Hb greater than 185 g/L or haematocrit above 0.52 in men, and Hb greater than 165 g/L or haematocrit above 0.48 in women, indicate that erythrocytosis is likely but isotope studies may be required (Table 15.2).

Once established, absolute polycythaemia can be subdivided into **primary polycythaemia** or **polycythaemia vera** (in which there is an intrinsic overactivity in the bone marrow) or **secondary polycythaemia** in which the bone marrow is driven by increase in erythropoietin as a result of factors such as smoking or altitude (Table 15.3).

Table 15.2 Radiodilution methods for measuring red cell and plasma volume.

	Normal	Primary or secondary polycythaemia	Relative polycythaemia
Total red cell volume (^{51}Cr)	Men 25–35 mL/kg Women 22–32 mL/kg	Increased	Normal
Total plasma volume (^{125}I -albumin)	40–50 mL/kg	Normal	Decreased

Table 15.3 Causes of polycythaemia (erythrocytosis).

Primary erythrocytosis

Congenital

Erythropoietin receptor mutations

Acquired

Polycythaemia vera

Secondary erythrocytosis

Congenital

Defects of the oxygen-sensing pathway

VHL gene mutation (Chuvash erythrocytosis)

PHD2 mutations

HIF-2 α mutations

Other congenital defects

High oxygen-affinity haemoglobin

Acquired

Erythropoietin-mediated

Central hypoxia

Chronic lung disease

Right-to-left cardiopulmonary vascular shunts

Carbon monoxide poisoning

Smoking

Obstructive sleep apnoea

High altitude

Local hypoxia

Renal artery stenosis

End-stage renal disease

Hydronephrosis

Renal cysts (polycystic kidney disease)

Post-renal transplant erythrocytosis

Pathological erythropoietin production

Tumours – cerebellar haemangioblastoma, meningioma, parathyroid tumours, hepatocellular carcinoma, renal cell cancer, pheochromocytoma, uterine leiomyoma

Drug-associated

Erythropoietin administration

Androgen administration

Primary polycythaemia (erythrocytosis)

Congenital

(See below.)

Acquired

This is caused in nearly all cases by the acquisition of mutations in the *JAK2* gene.

Polycythaemia vera (PV)

In PV the increase in red cell volume is caused by a clonal malignancy of a marrow stem cell. The disease results from somatic mutation of a single haemopoietic stem cell which gives

Table 15.4 Criteria for diagnosis of polycythaemia vera.
Source: McMullin M.F. *et al.* (2007) *B J Haem* 138: 821.
Reproduced with permission of John Wiley & Sons.

JAK2-positive polycythaemia vera

A1	High haematocrit (>0.52 in men, >0.48 in women) or raised red cell mass (>25% above predicted)*
A2	Mutation in <i>JAK2</i>

Diagnosis requires both criteria to be present

JAK2-negative polycythaemia vera

A1	Raised red cell mass (>25% above predicted) or haematocrit >0.60 in men, >0.56 in women.
A2	Absence of mutation in <i>JAK2</i>
A3	No cause of secondary erythrocytosis
A4	Palpable splenomegaly
A5	Presence of an acquired genetic abnormality (excluding <i>BCR-ABL</i>) in the haematopoietic cells
B1	Thrombocytosis (platelet count $>450 \times 10^9/L$)
B2	Neutrophil leucocytosis (neutrophil count $>10 \times 10^9/L$ in non-smokers; $>12.5 \times 10^9/L$ in smokers)
B3	Radiological evidence of splenomegaly
B4	Endogenous erythroid colonies or low serum erythropoietin

Diagnosis requires A1 + A2 + A3 + either another A or two B criteria

* WHO (2008) uses haemoglobin >185 g/L in men and 165 g/L in women as a major criterion, in *JAK2*+ cases and hypercellular marrow as a minor criterion as well as criteria A2 and B4 above.

its progeny a proliferative advantage. The *JAK2V617F* mutation is present in haemopoietic cells in about 97% of patients and a mutation in exon 12 is seen in some of the remainder. Although the increase in red cells is the diagnostic finding, in many patients there is also an overproduction of granulocytes and platelets. Some families have an inherited predisposition to myeloproliferative disease but *JAK2* or *CALR* mutations are not present in the germline.

Diagnosis

Making the diagnosis of PV in a patient who presents with polycythaemia can be difficult (Table 15.4).

Clinical features

PV is usually a disease of older people and has an equal sex incidence. Clinical features result from hyperviscosity, hypervolaemia, hypermetabolism or thrombosis.

- 1 Headaches, dyspnoea, blurred vision and night sweats. Pruritus, characteristically after a hot bath, can be a severe problem.
- 2 Plethoric appearance: ruddy cyanosis (Fig. 15.4), conjunctival suffusion and retinal venous engorgement.
- 3 Splenomegaly in 75% of patients (Fig. 15.5).



Figure 15.4 Polycythaemia vera: facial plethora and conjunctival suffusion in a 63-year-old woman. Haemoglobin 180 g/L; total red cell volume 45 mL/kg.



Figure 15.5 Splenomegaly: enlarged liver and spleen in a patient with polycythaemia vera.




Figure 15.6 (a) The feet of a 72-year-old man with polycythaemia rubra vera. There is inflammation of the right metatarsophalangeal and other joints caused by uric acid deposits. (b) Gangrene of the left fourth toe in essential thrombocythaemia.

- 4 Haemorrhage or thrombosis, either arterial or venous, may be seen.
- 5 Gout (as a result of raised uric acid production; Fig. 15.6a).

Laboratory findings

- 1 The haemoglobin, haematocrit and red cell count are increased. The total red cell volume (Table 15.2) is increased.
- 2 A neutrophil leucocytosis is seen in >50% and some have increased basophils.
- 3 A raised platelet count is present in about half of patients.
- 4 The *JAK2* mutation is present in the bone marrow and peripheral blood granulocytes in over 95% of patients.
- 5 The bone marrow is hypercellular with trilineage growth, as assessed by trephine biopsy (Fig. 15.7a).

- 6 Serum erythropoietin is low.
- 7 Plasma urate is often increased; the serum lactate dehydrogenase (LDH) is normal or slightly raised.
- 8 Circulating erythroid progenitors (erythroid colony-forming unit, CFU_E, and erythroid burst-forming unit, BFU_E; see p. 13) are increased compared to normal and grow *in vitro* independently of added erythropoietin (endogenous erythroid colonies).
- 9 Chromosome abnormalities (e.g. deletions of 9p or 20q) are found in a minority of subjects and mutations in *TET-2* or other epigenetic genes (see Fig. 16.1) occur in 10–20%.

Treatment

Treatment is aimed at maintaining a normal blood count. The haematocrit should be maintained at around 0.45 and the platelet count below $400 \times 10^9/L$.

Venesection

Venesection to reduce the haematocrit to less than 0.45 is particularly useful when a rapid reduction of red cell volume is required (e.g. at the start of therapy). It is indicated especially in younger patients and those with mild disease. The resulting iron deficiency may limit erythropoiesis. Unfortunately, venesection does not control the platelet count.

Hydroxycarbamide (hydroxyurea)

This is used if there is poor control or intolerance of venesection, or symptomatic or progressive splenomegaly, thrombocytosis, weight loss or night sweats. Daily treatment can control the blood count and may need to be continued for many years (Fig. 15.8). The *JAK2* mutation affects platelet function leading to thrombosis or haemorrhage and it is therefore necessary to control both the platelet count and haematocrit to reduce the risk. Side-effects of hydroxycarbamide include myelosuppression, nausea and toxicity to the skin areas exposed to ultraviolet light.




Figure 15.7 Iliac crest trephine biopsies. (a) Polycythaemia vera: fat spaces are almost completely replaced by hyperplastic haemopoietic tissue. All haemopoietic cell lines are increased with megakaryocytes particularly prominent. (b) Primary myelofibrosis: normal marrow architecture is lost and haemopoietic cells are surrounded by increased fibrous tissue and intercellular substance.




Figure 15.8 Haematological response to therapy with hydroxycarbamide (hydroxyurea) in polycythaemia vera. Hb, haemoglobin; WBC, white blood cells.

JAK inhibitors

Drugs such as ruxolitinib (lestaurtinib, pacritinib and momelotinib are in trials) inhibit JAK2 activity and are effective in many patients. At present ruxolitinib is they are used in patients who are not controlled adequately or have side-effects with hydroxurea therapy, but the drugs may eventually come to be first-line therapy.

Interferon

α -Interferon suppresses excess proliferation in the marrow and has produced good haematological responses. It is less convenient than the oral agents and side-effects are frequent. It may be particularly valuable in controlling itching and is often used for younger patients to avoid early exposure to other drugs.

Aspirin

Low-dose aspirin reduces thrombotic complications without significant increased risk of major haemorrhage and is used in almost all patients.

Course and prognosis

Typically, the prognosis is good with a median survival over 10 years. Thrombosis and haemorrhage are the major clinical problems. Increased viscosity, vascular stasis and high platelet levels and altered platelet function may all contribute to thrombosis, whereas defective platelet function may promote haemorrhage. Transition from PV to myelofibrosis occurs in approximately 30% of patients and approximately 5% of patients progress to acute myeloid leukaemia (AML).

Congenital causes of primary polycythaemia

Congenital causes are relatively rare and include cases caused by mutations in the genes that regulate oxygen sensing (*VHL*, *PHD2* or *HIF2A*) (see Chapter 2) as well as mutation of the erythropoietin receptor and haemoglobin mutations that lead to high oxygen-affinity variants with subsequent tissue hypoxia. These patients often have a family history of polycythaemia and present at a young age.

Secondary polycythaemia

The causes of secondary polycythaemia are listed in Table 15.3.

Acquired causes are due to an increase in the erythropoietin level. Hypoxia caused by smoking or chronic obstructive airways disease is a common cause, and measurement of arterial oxygen saturation is a valuable test. Renal and tumour causes of inappropriate erythropoietin secretion are less common.

There is little evidence on which to guide a treatment plan but one approach is to advise venesection if the haematocrit is above 0.54 with the aim of reducing to a target around 0.5. A lower target for venesection may be used if there is hypertension, diabetes, dyspnoea, angina or a previous thrombotic episode. Low-dose aspirin may be of benefit for many patients.

Apparent polycythaemia

Apparent polycythaemia, also known as pseudopolycythaemia, is the result of plasma volume contraction. By definition, the red cell mass is normal. The cause is uncertain but it is far more common than PV. It occurs particularly in young or middle-aged men and may be associated with cardiovascular problems such as hypertension or cerebral transient ischaemic attacks. Diuretic therapy, heavy smoking, obesity and alcohol consumption are frequent associations. Venesection is recommended for patients with a recent history of thrombosis or with cardiovascular risk factors, with the aim of maintaining a haematocrit around 0.45–0.47.

Differential diagnosis of polycythaemia

The identification of the *JAK2* mutation has rationalized the approach to diagnosis of polycythaemia. A three-stage approach to diagnosis has been suggested:

1 Stage 1 History and examination

Full blood count/film

JAK2 mutation

Serum ferritin

Renal and liver function tests

If JAK2 is negative and there is no clear secondary cause, consider investigations in stage 2.

2 Stage 2 Red cell mass

Arterial oxygen saturation

Abdominal ultrasound

Serum erythropoietin level

Bone marrow aspirate and trephine

Cytogenetic analysis

BFU_E culture

Specialized tests may then be required.

3 Stage 3 Arterial oxygen dissociation

Sleep study

Lung function studies

Gene mutations *EPOR*, *VHL*, *PHD2*, *HIFs*

Essential thrombocythaemia

In this condition there is a sustained increase in the platelet count due to megakaryocyte proliferation and overproduction of platelets. The haematocrit is normal and the Philadelphia chromosome or *BCR-ABL1* rearrangement are absent. The bone marrow shows no collagen fibrosis. A persisting platelet count of greater than $450 \times 10^9/L$ is the central diagnostic feature, but other causes of a raised platelet count (particularly iron deficiency, inflammatory or malignant disorder and myelodysplasia) need to be fully excluded before the diagnosis can be made.

Fifty to 60% of patients show the *JAK2* (*V617F*) mutation and these cases tend to resemble more closely PV with higher haemoglobin and white cell counts than *JAK2*-negative cases (Table 15.5). The *JAK2* mutation also affects platelet function leading to a pro-thrombotic state. Mutations in the *CALR* gene are seen in around 75% of *JAK2*-negative ET patients, in total representing about a third of all patients. These patients tend to be younger and have higher platelet counts but a lower incidence of thrombosis (Table 15.5). Mutations within the *MPL* gene are seen in 4% of cases. Rare primary familial cases in children have been associated with mutations in the genes for thrombopoietin or its receptor *MPL*.

Diagnosis

This used to be based on the exclusion of other causes of chronic thrombocytosis but now that specific genetic lesions have been identified a positive diagnosis can be made in most cases.

Table 15.5 Typical clinical and laboratory features of essential thrombocythaemia associated with *JAK2* or *CALR* mutation

	<i>JAK2</i> mutated	<i>CALR</i> mutated
Age	Older	Younger
Haemoglobin	Higher	Lower
White cell count	Higher	Lower
Platelet count	Lower	Higher
Serum erythropoietin	Lower	Higher
Thrombosis risk	Higher	Lower
Transformation to PRV	Yes	No
Risk of transformation to myelofibrosis	Equal	Equal
Approximate survival	9 years	17 years

The suggested diagnostic criteria for essential thrombocythaemia are:

- A1** Sustained platelet count above $450 \times 10^9/L$.
 - A2** Presence of an acquired pathogenetic mutation (e.g. in *JAK2* or *CALR*).
 - A3** No other myeloid malignancy, PV, primary myelofibrosis, chronic myeloid leukaemia (CML) or myelodysplastic syndrome.
 - A4** No reactive cause for thrombocytosis and normal iron stores.
 - A5** Bone marrow trephine histology showing increased megakaryocytes with prominent large hyperlobulated forms; reticulin is generally not increased.
- Diagnosis requires A1–A3 or A1 + A3–A5

Clinical and laboratory findings

The dominant clinical features are thrombosis and haemorrhage. Most cases are asymptomatic and diagnosed on a routine blood count. Thrombosis may occur in the venous or arterial systems (Fig. 15.6b), whereas haemorrhage, as a result of abnormal platelet function, may cause either chronic or acute bleeding. Some patients, particularly those with the *JAK2* mutation, present with Budd–Chiari syndrome when the platelet count may be normal because of splenomegaly. A characteristic symptom is erythromelalgia, a burning sensation felt in the hands or feet and promptly relieved by aspirin. Up to 40% of patients will have palpable splenomegaly, whereas in others there may be splenic atrophy because of infarction.

Abnormal large platelets and megakaryocyte fragments may be seen on the blood film (Fig. 15.9). The bone marrow is similar to that in PV but an excess of abnormal megakaryocytes is typical. Cytogenetics and molecular analysis are performed to exclude *BCR-ABL1*+ CML. The condition must be distinguished from other causes of a raised platelet count




Figure 15.9 Peripheral blood film in essential thrombocythaemia showing increased numbers of platelets and a nucleated megakaryocytic fragment.

Table 15.6 Causes of a raised platelet count.

Reactive

Haemorrhage, trauma, postoperative

Chronic iron deficiency

Malignancy

Chronic infections

Connective tissue diseases (e.g. rheumatoid arthritis)

Post-splenectomy

Endogenous

Essential thrombocythaemia (*JAK2* mutation + or –)

Some cases of polycythaemia vera, primary myelofibrosis, *BCR-ABL1*+ chronic myeloid leukaemia, myelodysplasia (5q- or refractory anaemia with ring sideroblasts)

(Table 15.6). Platelet function tests are rarely needed, but are consistently abnormal, with failure of aggregation with adrenaline being particularly characteristic.

Prognosis and treatment

The principle is to reduce the risk of the major clinical problems of thrombosis or haemorrhage. Standard cardiovascular risk factors, such as cholesterol, smoking, diabetes, obesity and hypertension, should be identified and treated. Low-dose aspirin at 75 mg/day is generally recommended in all cases.

Patients at **high risk** include those over 60 years of age, and/or with previous thrombosis and/or with platelet count over $1500 \times 10^9/L$, and this group should be treated with hydroxycarbamide or anagrelide to reduce the platelet count. **Low-risk** patients are those aged below 40 years and here aspirin alone is sufficient. Optimum control of the **medium-risk** group (age 40–60 years) is uncertain.

Hydroxycarbamide is the most widely used treatment and is well tolerated, although after prolonged therapy some patients develop skin keratosis, epitheliomas, ulceration or pigmentation. Anagrelide is a good second-line treatment but has cardiovascular side-effects, and a possible increased risk of myelofibrosis is also of concern. These two drugs can be combined at low doses to reduce side-effects. α -Interferon is also effective and is often used in younger patients or during pregnancy. A long-acting PEGylated preparation is preferred. *JAK2* inhibitors are also being assessed.

Course

Often the disease is stationary for 10–20 years or more. The disease may transform after a number of years to myelofibrosis but the risk of transformation to AML is relatively low (less than 5%).

Primary myelofibrosis

The predominant feature of primary myelofibrosis is a progressive generalized reactive fibrosis of the bone marrow in association with the development of haemopoiesis in the spleen and liver (known as myeloid metaplasia). Clinically this leads to anaemia and massive splenomegaly (Fig. 15.10). In some




Figure 15.10 (a) Peripheral blood film in primary myelofibrosis. Leucoerythroblastic change with 'tear-drop' cells and an erythroblast. **(b)** Massive splenomegaly in a patient with myelofibrosis.

patients there is osteosclerosis. The fibrosis of the bone marrow is secondary to hyperplasia of abnormal megakaryocytes. Fibroblasts are stimulated by platelet-derived growth factor and other cytokines secreted by megakaryocytes and platelets.

The *JAK2*, *CALR* and *MPL* mutations occur in approximately 55%, 25% and 10% of patients respectively (Fig. 15.2). One-third of patients with similar features have a previous history of PV or ET and some patients present with clinical and laboratory features of both disorders.

Clinical features

- 1 An insidious onset in older people is usual with symptoms of anaemia.
- 2 Symptoms resulting from massive splenomegaly are frequent and include abdominal discomfort, pain or indigestion. Splenomegaly is the main physical finding (Fig. 15.5b).
- 3 Hypermetabolic symptoms such as loss of weight, anorexia, fever and night sweats are common.
- 4 Bleeding problems, bone pain or gout occur in a minority of patients.

Laboratory findings

- 1 Anaemia is usual but a normal or increased haemoglobin level may be found in some patients.
- 2 The white cell and platelet counts are frequently high at the time of presentation. Later in the disease leucopenia and thrombocytopenia are common.
- 3 A leucoerythroblastic blood film is found. The red cells show characteristic 'tear-drop' poikilocytes (Fig. 15.10).
- 4 Bone marrow is usually unobtainable by aspiration. Trephine biopsy (Fig. 15.7b) shows a fibrotic, hypercellular marrow. Increased megakaryocytes are frequently seen. In 10% of cases there is increased bone formation with increased bone density on X-ray.
- 5 *JAK2* is mutated in approximately 55% of cases and *CALR* mutations occur in 25%. The *CALR*-mutated patients have lower white cell and higher platelet counts and longer survival.

6 High serum urate and LDH levels reflect increased, although ineffective, haemopoiesis.

7 Transformation to AML occurs in 10–20% of patients.

Treatment

This is aimed at reducing the effects of anaemia and splenomegaly. Useful prognostic information may be obtained by analysis of the IPSS score (Table 15.7). Blood transfusions and regular folic acid therapy are useful in severely anaemic patients. Ruxolitinib is an oral JAK2 inhibitor that can reduce spleen size, improve constitutional symptoms and quality of life and increase survival. Hydroxycarbamide may also reduce

Table 15.7 A dynamic international prognostic scoring system (DIPSS) for patients with primary myelofibrosis. Median survival times are 185, 78, 35, and 16 months for the low (0 points), intermediate-1 (1 point), intermediate-2 (2–3 points), and high-risk categories (4–6 points) respectively. Source: From Gangat *et al.* (2011) *J Clin Oncol* 29(4): 392–7.

Variable	IPSS*	DIPSS	DIPSS Plus
Age >65 years	✓	✓	✓
Constitutional symptoms	✓	✓	✓
Hb <100 g/L	✓	✓	✓
Leucocyte count >25 × 10 ⁹ /L	✓	✓	✓
Circulating blasts ≥1%	✓	✓	✓
Platelet count <100 × 10 ⁹ /L			✓
RBC transfusion need			✓
Unfavourable karyotype			✓
	1 point each	1 point each but Hb = 2	1 point each

*Unfavourable karyotype includes +8, +–7/7q, –5/5q, complex and others

splenomegaly and hypermetabolic symptoms. Trials of thalidomide, lenalidomide, azacytidine and histone deacetylase inhibitors are in progress. Erythropoietin can be tried but may cause splenic enlargement.

Splenectomy is considered for patients with severe symptomatic splenomegaly: mechanical discomfort, thrombocytopenia, portal hypertension, excessive transfusion requirements. Allopurinol may be indicated to prevent gout and urate nephropathy from hyperuricaemia. Allogeneic stem cell transplantation may be curative for young patients.

The median survival is around 5 years and causes of death include heart failure, infection and leukaemic transformation.

Mastocytosis

Mastocytosis is a clonal neoplastic proliferation of mast cells that accumulate in one or more organ systems. Mast cells are similar to basophils and survive for months or years in vascular tissues and most organs. Systemic mastocytosis is a clonal myeloproliferative disorder involving usually the bone marrow, heart, spleen, lymph nodes and skin.

The somatic *KIT* mutation Asp816Val is detected in the majority of patients and may be partly responsible for autonomous growth and enhanced survival of the neoplastic mast cells.

Symptoms are related to histamine and prostaglandin release and include flushing, pruritus, abdominal pain and bronchospasm. The skin usually shows urticaria pigmentosa (Fig. 15.11). Serum tryptase is increased and can be used to monitor treatment. Antihistamine drugs are valuable and interferon, chlorodeoxyadenosine and tyrosine kinase inhibitors can be helpful. In many patients the disease runs a chronic indolent course. In others an aggressive course may be associated with AML, mast cell leukaemia or other haemopoietic conditions (see Appendix 2).



Figure 15.11 Systemic mastocytosis: female 72 years; widespread erythematous, confluent plaques of urticaria pigmentosa over chest, abdomen and upper arms. Source: Courtesy of Professor M. Rustin.


- Myeloproliferative neoplasms are a group of conditions arising from marrow stem cells and characterized by clonal proliferation of one or more haemopoietic components in the bone marrow. The three major subtypes are: polycythaemia vera (PV); essential thrombocythaemia (ET); and primary myelofibrosis.
- These subtypes are closely related to each other and mutation of the *JAK2* gene is detected in almost all patients with PV and approximately 50% of those with ET and primary myelofibrosis. Mutations in *CALR* or *MPL* are present in most cases without a *JAK2* mutation.
- Polycythaemia is defined as an increase in the haemoglobin concentration and the major subdivision is into *absolute polycythaemia*, in which the red cell mass is raised, and *relative polycythaemia*, in which the red cell volume is normal but the plasma volume is reduced.
- Absolute polycythaemia is divided into primary polycythaemia, known as polycythaemia vera (PV), or secondary polycythaemia.
- The diagnosis of PV is made by finding polycythaemia together with a *JAK2* mutation. It occurs in older patients and the increase in blood viscosity leads to headaches, plethoric appearance and splenomegaly.

(Continued)

- Treatment aims to maintain the haematocrit around 0.45. Useful approaches include venesection, hydroxyurea and *JAK2* inhibitors. Aspirin is also given. Survival is usually over 10 years but there may be progression to leukaemia or myelofibrosis.
- Secondary polycythaemia can arise from rare congenital causes or acquired disorders such as lung disease or tumours that secrete erythropoietin. Venesection may be needed.
- Essential thrombocythaemia is diagnosed by persistent raised platelet count in the absence of other causes. *JAK2*, *CALR* or *MPL* genes are mutated in most cases.
- The predominant feature of primary myelofibrosis is a progressive generalized reactive fibrosis of the bone marrow in association with the development of haemopoiesis in the spleen and liver. Symptoms usually result from anaemia and a grossly enlarged spleen.
- Diagnosis is made on blood film, which shows a leucoerythroblastic appearance, together with bone marrow biopsy and *JAK2*, *CALR* and *MPL* mutation screen. Treatment is with red cell transfusion and *JAK2* inhibition.
- Systemic mastocytosis is a clonal proliferation of mast cells with involvement of bone marrow, skin (as urticaria pigmentosa) and other organs.



Now visit www.wileyessential.com/haematology to test yourself on this chapter.


A vertical strip on the left side of the page showing a microscopic view of blood cells. It features several red blood cells (erythrocytes) in shades of red and pink, and a few white blood cells (leukocytes) with distinct nuclei, set against a dark blue background.

CHAPTER 16

Myelodysplasia

Key topics

■ Myelodysplasia (myelodysplastic syndromes)	178
■ Classification	179
■ Laboratory findings	180
■ Treatment	182
■ Myelodysplastic/myeloproliferative neoplasms	184



Myelodysplasia (myelodysplastic syndromes, MDS)

This is a group of clonal disorders of haemopoietic stem cells characterized by increasing bone marrow failure in association with dysplastic changes in one or more cell lineages (Table 16.1). A hallmark of the disease is simultaneous proliferation and apoptosis of haemopoietic cells (*ineffective haemopoiesis*) leading to the paradox of a hypercellular bone marrow but pancytopenia in peripheral blood. There is a tendency to progress to acute myeloid leukaemia (AML), although death often occurs before this develops.

In most cases, the disease is **primary** but in a significant proportion of patients it is **secondary** to chemotherapy and/or radiotherapy that has been given for treatment of another malignancy. This latter type is termed **therapy-related MDS** (*t-MDS*) and is now classified with therapy-related AML.

Pathogenesis

The pathogenesis of the MDS is unclear but is presumed to start following genetic change in a multipotent haemopoietic progenitor cell. The immune system may have a minor role in suppressing bone marrow function and immunosuppression is sometimes used in treatment (see below).

Chromosome abnormalities are frequent (Table 16.2). In addition, molecular analysis shows that MDS typically carry two or three point mutations. These driver mutations usually involve genes involved in epigenetic processes, DNA methylation (*TET2*, *DNMT3A* and *IDH1/2*) and chromatin modification (*ASXL1* and *EZH2*) as well as in RNA splicing (Fig 16.1). Some of these gene mutations, e.g. *TET2*, may be found in other myeloid malignancies, including acute myeloid leukaemia (AML) and the myeloproliferative diseases (MPD), while others (e.g. *ASXL1*, *SF3B1*) are relatively specific to MDS. A striking example is mutation in the

Table 16.1 The World Health Organization (2008) classification of myelodysplasia (modified; see also Appendix).

Subtype	Peripheral blood	Bone marrow	(% blasts)	Relative proportion (%)
<i>Refractory cytopenia with unilineage dysplasia (RCUD)</i>				
Refractory anaemia (RA)	Anaemia	Unilineage erythroid dysplasia (in >10% cells), <5% blasts	<5	10–20
Refractory neutropenia (RN)	Neutropenia	Unilineage granulocytic dysplasia, <5% blasts	<5	<1
Refractory thrombocytopenia (RT)	Thrombocytopenia	Unilineage megakaryocytic dysplasia, <5% blasts	<5	<1
<i>Refractory anaemia with ring sideroblasts (RARS)</i>	Anaemia	Unilineage erythroid dysplasia, >15% erythroid precursors are ring sideroblasts, <5% blasts	<5	3–10
<i>Refractory cytopenia with multilineage dysplasia (RCMD)</i>	Cytopenia(s)	Multilineage dysplasia +/- ring sideroblasts, <5% blasts. No Auer rods	<5	30
<i>Refractory anaemia with excess blasts type 1 (RAEB-1)</i>	Cytopenia(s)	Unilineage or multilineage dysplasia	5–9	20
<i>Refractory anaemia with excess blasts type 2 (RAEB-2)</i>	Cytopenia(s)	Unilineage or multilineage dysplasia	10–19	20
<i>Myelodysplastic syndrome associated with isolated del(5q)</i>	Anaemia Normal or high platelet count	5q31 deletion Anaemia, hypolobulated megakaryocytes	<5	<5%
<i>Childhood myelodysplastic syndrome</i>	Pancytopenia	Multilineage dysplasia	<5	<1%

Table 16.2 Examples of cytogenetic abnormalities in myelodysplasia.

Prognosis	Cytogenetic abnormality
Very good	–Y or del(11q)
Good	Normal or del(5q)
Intermediate	del(7q) or double independent clones
Poor	inv(3) or double including -7 or 1del(7q),
Very poor	Complex: >3 abnormalities

SF3B1 gene, involved in splicing RNA, which is seen in almost all cases of the RARS subtype of MDS but not in *de novo* AML or myeloproliferative diseases. Some of the epigenetic mutations can also be detected in apparently normal marrow in up to 20% of the elderly (Table 11.2). On the other hand, mutations in *FLT3* and *NPM1* are relatively specific for *de novo* AML and when MDS transforms to AML these AML-specific mutations appear.


Other genes frequently mutated in MDS (and AML) are involved in transcription regulation (*RUNX1*), DNA repair (*TP53*), signal transduction (*NRAS* and *KRAS*), and in the cohesin complex involved in mitosis (*STAG2*).

Classification

The myelodysplastic syndromes are classified on the basis of the blood count, their morphological appearance, and the number of blast cells in blood or bone marrow and genetic analysis (Table 16.1). The classification of MDS can be confusing and flow charts are helpful in the approach to diagnosis (Fig. 16.2). Although the classification appears complex, the principles are as follow:


- Dysplasia may be present solely in a single lineage – red cells (**refractory anaemia**), neutrophils or platelets – or present in two or more myeloid lineages (**refractory cytopenia with multilineage dysplasia**; RCMD).

1. Methylation of DNA (regulates gene expression)



2. Modification of histones (regulates gene transcription)

ASXL1 and **EZH2** involved in modification of histone and chromatin structure, enhancing or inhibiting gene expression



3. Messenger RNA splicing




Figure 16.1 Genes involved in DNA methylation, histone modification and in mRNA splicing that are frequently mutated (loss of function or gain in function) in myelodysplasia, acute myeloid leukaemia or myeloproliferative diseases.

Table 16.3 IPSS-R prognostic score values. Source: Based on Greenberg PL, Tuechler H, Schanz J et al. 2012 Blood 120: 2454–65)

Prognostic variable	0	0.5	1	1.5	2	3	4
Cytogenetics (see Table 16.2)	Very Good		Good		Intermediate	Poor	Very Poor
Bone marrow blast %	≤2		>2–<5		5–10	>10	
Hb concentration (g/L)	≥100		80–<100	<80			
Platelet count	≥100	50–<100	<50				
Neutrophil count (× 10 ⁹ /L)	≥0.8	<0.8					

Table 16.4 IPSS-R prognostic risk categories/scores and clinical outcomes. Time taken for 25% of cases to evolve to AML. Source: Based on Greenberg PL, Tuechler H, Schanz J *et al.* (2012) *Blood* 120: 2454–65.)

Risk category	Risk score	Survival (median in years)	Median time to evolution of AML in 25% + of cases (years)
Very low	≤1.5	8.8	Not reached
Low	>1.5–3	5.3	10.8
Intermediate	>3–4.5	3.0	3.2
High	>4.5–6	1.6	1.4
Very high	>6	0.8	0.73

- Erythroid dysplasia can also be associated with **ring sideroblasts** to define a unique subtype. The definition of a pathological ring sideroblast is an erythroid precursor with five or more iron granules encircling at least one-third of the nucleus.
- If the blast cell count is increased in the bone marrow, the diagnosis is made of **refractory anaemia with excess blasts** (RAEB) and these subtypes have a poor prognosis, especially if the blast count is between 10 and 19% rather than 5–9% (Table 16.1).
- **5q-Syndrome** receives its own classification. The key gene that is deleted is *RPS14*, encoding a ribosomal protein (see Fig. 22.3). It is more common in women and typically there is a macrocytic anaemia with thrombocytosis in 50% of cases. It has a particularly good prognosis (Tables 16.3 and 16.4).
- Where the monocyte count is over $1 \times 10^9/L$ the diagnosis becomes **chronic myelomonocytic leukaemia**.

Clinical features

The disease has an incidence of 4 in 100 000 and a slight male predominance. Over half of patients are over 70 years and fewer than 25% are less than 50. The evolution is often slow and the disease may be found by chance when a patient has a blood count for some unrelated reason. The symptoms, if present, are those of anaemia, infections or of easy bruising or bleeding (Fig. 16.3). In some patients transfusion-dependent anaemia dominates the course, while in others recurring infections or spontaneous bruising and bleeding are the major clinical problems. The function of the neutrophils, monocytes and platelets is often impaired so that infections and bleeding may occur out of proportion to the severity of the cytopenia. The spleen is not usually enlarged.

It is important to remember that dysplastic features in bone marrow may be seen in a wide range of conditions, such as excess alcohol intake, megaloblastic anaemia, parvovirus, recovery from cytotoxic chemotherapy and granulocyte colony-stimulating factor (G-CSF) therapy. These must be ruled out before making a diagnosis of MDS and diagnostic tests may need to be repeated over time in some patients.

Laboratory findings

Peripheral blood

Pancytopenia is a frequent finding. The red cells are usually macrocytic but occasionally hypochromic; normoblasts may be present. The reticulocyte count is low. Granulocytes are often reduced in number and frequently show lack of granulation (Fig. 16.4). Their chemotactic, phagocytic and adhesive functions are impaired. The Pelger abnormality (single or bilobed nucleus) is often present. The platelets may be unduly large or small and are usually decreased in number, but in 10% of cases are elevated. In poor prognosis cases, variable numbers of myeloblasts are present in the blood.

Bone marrow

The cellularity is usually increased. A small number of dysplastic cells may be seen in marrow from healthy elderly individuals so at least 10% of the cells in a lineage should be dysplastic in order to consider the diagnosis of MDS. Typical dysplastic features are:

- **Erythroid:** Multinucleate normoblasts, internuclear bridges and nuclear budding (Fig. 16.4). Ring sideroblasts are seen in a specific subset of MDS and are caused by iron deposition in the mitochondria of erythroblasts.
- **Myeloid:** The granulocyte precursors often show defective granulation and may be difficult to distinguish from monocytes.




Figure 16.2 Approach to the diagnosis of myelodysplasia. MP, myeloproliferative. Source: Based on Bennett J.M. & Komrokji R.S. (2006) *Hematology* 10: Suppl 1:258–69.



(a)



(b)

Figure 16.3 Myelodysplasia. **(a)** A 78-year-old male patient with refractory anaemia had recurring infections of the face and maxillary sinuses associated with neutropenia (haemoglobin 98 g/L; white cells 1.3×10^9 /L; neutrophils 0.3×10^9 /L; platelets 38×10^9 /L). **(b)** Purpura in a 58-year-old female with refractory anaemia (haemoglobin 105 g/L; white cells 2.3×10^9 /L; platelets 8×10^9 /L).

■ **Megakaryocytes:** Dysplastic megakaryocytes may show micronuclear, small binuclear or polynuclear forms (Fig. 16.4). In a minority of cases (about 20%) the marrow is hypocellular and may resemble aplastic anaemia; in others there is fibrosis. A sample of bone marrow should be sent for genetic analysis.

Genetic abnormalities

Cytogenetic analysis is essential and common abnormalities include 5q-, partial or total loss of chromosomes 5 or 7, or trisomy 8 (Table 16.2). Several cytogenetic abnormalities are considered so diagnostic of MDS that they allow diagnosis of MDS even in the absence of morphological abnormalities within cells.

The mutations that may be found by molecular testing have been described on p.178 and Figure 16.1 illustrates those involved in epigenetic processes.

Treatment

The classification of MDS can be confusing and flow charts are helpful in the approach to diagnosis (Fig. 16.2). Treatment for MDS has improved significantly in recent years. A key subdivision is into patients with low-risk or high-risk disease. The Revised International Prognostic Scoring System (IPSS-R) classifies the patients according to the percentage of marrow blasts, the type of karyotype abnormality, and the number and severity of cytopenias (Table 16.3). For ease, 'clinical calculators' are now available online, e.g. at www.ukmidsforum.org. At this stage it is not clear if patients with 'intermediate-risk' prognosis should be treated according to low-risk or high-risk regimens.

Low-risk myelodysplastic syndromes

Patients with less than 5% of blasts in the marrow and only one cytopenia and favourable cytogenetics are defined as having low-grade MDS. Intensive chemotherapy is rarely used for these patients. Instead they are not treated or, if necessary, attempts may be made to improve marrow function with haemopoietic growth factors, either singly or in combination. Erythropoietin may improve anaemia although the haemoglobin should not be raised above 120 g/L. G-CSF shows synergy with erythropoietin and may increase the response rate. Ciclosporin or anti-thymocyte globulin occasionally helps, particularly for those with a hypocellular bone marrow.

Transfusion support with red cells or platelets is often needed. Thrombomimetics, such as eltrombopag or romiplostim, can be given in the event of bleeding, but trials to determine whether they increase the risk of AML in this setting are in progress. Antibiotics may be required. In the long term, iron overload may be a problem after multiple transfusions; iron chelation therapy should be considered after 20–50 units have been transfused or if the ferritin rises above 1000 µg/L. Measurement of liver and cardiac iron by MRI is helpful in deciding whether or not chelation therapy is indicated. Lenalidomide is particularly effective in MDS associated with del(5q) where it can often reduce the size of the del(5q) clone and reduce transfusion requirements.

In selected patients standard or low-intensity allogeneic stem cell transplantation (SCT) offers a permanent cure.




Figure 16.4 Myelodysplasia: appearances of the peripheral blood and bone marrow. **(a)** Multinucleate polychromatic erythroblasts. **(b)** Perl's stain showing iron overload in macrophages of a bone marrow fragment. **(c)** Ring sideroblasts. **(d)** White cells showing pseudo-Pelger cells, agranular myelocytes and neutrophils. **(e)** Monocytoid cells and an agranular neutrophil. **(f)** Mononuclear megakaryocyte.

High-risk myelodysplastic syndromes

In these patients a variety of treatments have been attempted to improve the overall prognosis, with varying degrees of success.

Stem cell transplantation

SCT offers the prospect of complete cure for MDS and the advent of non-myeloablative conditioning is increasing the age range of patients that may be treated. Chemotherapy should be given prior to the transplant in patients with >10% blasts in the marrow.

Intensive chemotherapy

Chemotherapy as given in AML (see p. 149) is often tried in high-risk patients. Although the majority may obtain a remission, relapse is likely and frequently occurs within a few months. The risks of intensive chemotherapy are great because prolonged pancytopenia may occur in some cases without normal haemopoietic regeneration, presumably because normal stem cells are not present.

Hypomethylating agents

Azacitidine (azacitidine) and decitabine are DNA methyltransferase inhibitors which inhibit methylation of newly formed DNA. They improve blood counts in a minority of high-risk MDS patients. Azacitidine is given for 7 days every month and is continued for as long as the patient is responding. Studies have shown that it can improve survival by approximately 9 months.

General support care only

This is most suitable in elderly patients with other major medical problems. Transfusions of red cells and platelets, and therapy with antibiotics and antifungals, are given as needed (see Chapter 12).

Myelodysplastic/myeloproliferative neoplasms

These disorders are classified between myelodysplasia and myeloproliferative disease as they show the presence of dysplastic features but also increased number of circulating cells in one or more lineage (Table 16.5). They share common clinical and genetic features such as mutations of the *TET2* tumour suppressor gene in about 20% of cases and of *JAK2* in a smaller proportion of patients.

Chronic myelomonocytic leukaemia

This is defined by a persistent monocytosis of more than $1.0 \times 10^9/L$ with blasts <20% in the marrow, dysplasia in other lineages and negative for the *BCR-ABL1* translocation. The total white cell count is usually raised and may exceed $100 \times 10^9/L$. Patients may develop skin rashes and around half have splenomegaly. Bruising is frequent and gum hypertrophy

Table 16.5 Classification of myelodysplastic/myeloproliferative neoplasms.

	Diagnostic features
Chronic myelomonocytic leukaemia	Monocytosis $>1 \times 10^9/L$
Atypical chronic myeloid leukaemia BCR-ABL1 negative	WBC $>13 \times 10^9/L$ BCR-ABL1 absent
Juvenile myelomonocytic leukaemia	
Myelodysplastic/myeloproliferative neoplasm, unclassifiable	
Refractory anaemia with ring sideroblasts associated with marked thrombocytosis	Platelet count $>450 \times 10^9/L$ Large atypical megakaryocytes

and lymphadenopathy may also be present. *TET-2*, *ASXL1* and other mutations (Fig 16.1) are frequent. Treatment is difficult, although oral hydroxycarbamide or etoposide may be useful. SCT may be tried in younger patients. Median survival is approximately 2 years, with increased marrow blasts a predictor of poor outcome.

Atypical chronic myeloid leukaemia

These patients have an increased white cell count with mainly granulocytes and granulocyte precursors in the blood and hypercellular bone marrow but the Philadelphia chromosome and *BCR-ABL1* fusion gene are not present. There are usually some morphological features in the blood or bone marrow of myelodysplasia. Treatment is difficult and the outlook is poor.

Juvenile myelomonocytic leukaemia (JMML)


This presents in the first 4 years of life and has features of both myelodysplasia and a myeloproliferative disease. There is often an eczematous skin rash, hepatosplenomegaly and lymphadenopathy. There is a monocytosis to more than $1.0 \times 10^9/L$ and clonal cytogenetic change. The only curative treatment is allogeneic SCT. If untreated, death usually occurs within 4 years, often from acute transformation with leukaemic infiltration (e.g. of the lungs). Co-mutations in the *TET2* and *SRSF2* genes are commonly found.

Children with two genetic disorders, Noonan's syndrome and neurofibromatosis, are at increased risk of JMML and mutations in the genes *PTPN11* and *NF1*, which underlie these genetic disorders, are frequent in cases of JMML not associated with these disorders.

- Myelodysplasia includes a group of clonal disorders of haemopoietic stem cells that lead to bone marrow failure and low blood cell counts. A hallmark of the disease is the increased proliferation and apoptosis of haemopoietic cells leading to the paradox of a hypercellular bone marrow but pancytopenia. There is a tendency to progress to acute myeloid leukaemia.
- In most cases, the disease is *primary* but it may be *secondary* to chemotherapy or radiotherapy given for treatment of another malignancy.
- The main clinical features of anaemia, infection and bleeding, are caused by reduction in the blood count. Most patients are over 70 years of age.
- Diagnosis is made by examination of the blood and bone marrow together with cytogenetic and molecular genetic studies of the tumour cells. They are classified into eight major subtypes.
- Scoring systems can divide patients into those with low-grade or high-grade disease.
- Low-grade disease may not need treatment. Haemopoietic growth factors, lenalidomide or blood product support are useful when required.
- High-grade myelodysplasia may be treated by intensive chemotherapy, demethylating drugs or stem cell transplantation. Allogeneic transplantation is the only curative procedure.
- Myelodysplastic/myeloproliferative neoplasms are a group of disorders classified between myelodysplasia and myeloproliferative disorders and show the presence of dysplastic features but also increased number of circulating white cells.



Now visit www.wileyessential.com/haematology to test yourself on this chapter.


A vertical strip on the left side of the page showing a microscopic view of blood cells. It features several large, red, biconcave disc-shaped cells (erythrocytes) and a few smaller, white, irregularly shaped cells (leukocytes) with prominent nuclei. The background is a deep blue.

CHAPTER 17

Acute lymphoblastic leukaemia

Key topics

■ Incidence and pathogenesis	187
■ Classification	188
■ Investigations	188
■ Treatment	192
■ Minimal residual disease	192
■ Specific therapy of ALL in adults	194



Acute lymphoblastic leukaemia (ALL) is caused by an accumulation of lymphoblasts in the bone marrow and is the most common malignancy of childhood. The definition of acute leukaemia and differentiation of ALL from acute myeloid leukaemia are described in Chapter 13.

Incidence and pathogenesis

The incidence of ALL is highest at 3–7 years, with 75% of cases occurring before the age of 6. There is a secondary rise after the age of 40 years. Eighty-five per cent of cases are of B-cell lineage and have an equal sex incidence; there is a male predominance for the 15% of T-cell ALL (T-ALL).

The pathogenesis is varied. **A proportion of cases of childhood ALL are initiated by genetic mutations that occur during development *in utero* (Fig. 17.1).** Studies in identical twins have shown that both may be born with the same chromosomal abnormality (e.g. t(12;21) *ETV6-RUNX1* translocation). This has presumably arisen spontaneously in a progenitor cell that has passed from one twin to the other as a result of the shared placental circulation. Environmental exposure during pregnancy may be important for this first event. One twin may develop ALL early (e.g. at age 4) because of a second transforming event affecting the copy numbers of several genes including those in B-cell development (see

below). The other remains well or develops ALL later. The *ETV6-RUNX1* translocation is present in the blood of approximately 10% of newborn infants but only 1 in 100 of these go on to develop ALL at a later date. The mechanism of the 'second genetic hit' within the tumour cell is unclear but an abnormal response of the immune system to infection is suggested by epidemiological studies. In other cases, the disease seems to arise as a postnatal mutation in an early lymphoid progenitor cell.

Children with a high level of social activity, notably those attending early nursery daycare, have a reduced incidence of ALL, whereas those living in more isolated communities and who have a reduced exposure to common infections in the first years of life have a higher risk.

Certain germline polymorphisms in a group of genes mainly involved in B-cell development (e.g. *IKZF1*) appear to predispose to ALL since they are more frequent in children with B-cell ALL (B-ALL) than controls. Interestingly, *IKZF1* is also deleted in the leukaemic cells in 30% of high risk B-ALL and 95% of ALL BCR-ABL1 positive cases. In general the genomic landscape in ALL is characterized by primary chromosomal abnormalities and a wide range of secondary deletions and mutations involving key pathways implicated in leukaemogenesis. For childhood ALL an average of 11 somatically acquired structural variations are present.




Figure 17.1 Prenatal origin of acute lymphoblastic leukaemia (ALL) in a pair of identical twins. Both tumours had an identical t(12;21) translocation. ALL was diagnosed in the first twin at age 5 years and in the second at age 14 years, indicating probable origin of the leukaemic clone *in utero* and dissemination to both twins via a shared placental blood supply. Because of the prolonged latency of the ALL it is presumed that a secondary event is required to initiate the development of frank leukaemia. At the time of the diagnosis of ALL in twin 1 the t(12;21) translocation could be detected in the bone marrow of twin 2. It is likely that such a 'fetal origin' of childhood ALL occurs in a significant number of sporadic ALL cases. Source: Adapted from J.L. Wiemels *et al.* (1999) *Blood*, 94(3): 1057–62.

Table 17.1 Classification of acute lymphoblastic leukaemia (ALL) according to WHO (modified from WHO 2008; see also Appendix).

B acute lymphoblastic leukaemia with recurrent genetic abnormalities

ALL with t(12;21)

ALL with t(9;22)

ALL with t(11q23; variable)

Hyperdiploidy (>50 chromosomes)

Hypodiploidy (<45 chromosomes)

T- acute lymphoblastic leukaemia

N.B. A minority of patients present with nodal or extranodal masses and <20% blasts in the marrow and are called lymphoblastic lymphoma if the tumour cells resemble those of ALL.

Classification

Acute lymphoblastic leukaemia, B cell or T cell, is subclassified by WHO (2008) according to the underlying genetic defect (Table 17.1). Within B-ALL there are several specific genetic subtypes such as those with the t(9;22) or t(12;21) translocations, rearrangements of the *MLL* gene or alteration in chromosome number (diploidy) (Table 17.1). The subtype is an important guide to the optimal treatment protocol and to prognosis. In T-ALL an abnormal karyotype is found in 50–70% of cases and the NOTCH signalling pathway is activated in most cases (see below).

Clinical features

Clinical features are a result of the following.

Bone marrow failure

- Anaemia (pallor, lethargy and dyspnoea);
- Neutropenia (fever, malaise, features of mouth, throat, skin, respiratory, perianal or other infections);
- Thrombocytopenia (spontaneous bruises, purpura, bleeding gums and menorrhagia).


Organ infiltration

This causes tender bones, lymphadenopathy (Fig. 17.2), moderate splenomegaly, hepatomegaly and meningeal syndrome (headache, nausea and vomiting, blurring of vision and diplopia). Fundal examination may reveal papilloedema and sometimes haemorrhage. Many patients have a fever which usually resolves after starting chemotherapy. Less common manifestations include testicular swelling or signs of mediastinal compression in T-ALL (Fig. 17.3).

If lymph node or solid extranodal masses predominate with <20% blasts in the marrow, the disease is called lymphoblastic lymphoma but is treated as ALL.

Investigations

Haematological investigations reveal a normochromic normocytic anaemia with thrombocytopenia in most cases. The total white cell count may be decreased, normal or increased to $200 \times 10^9/L$ or more. The blood film typically shows a variable numbers of blast cells. The bone marrow is hypercellular



(a)



(b)

Figure 17.2 Acute lymphoblastic leukaemia. (a) Marked cervical lymphadenopathy in a boy. (b) Facial asymmetry in a 59-year-old man due to a right lower motor neurone seventh nerve palsy resulting from meningeal leukaemic infiltration. Source: Hoffbrand V.A., Pettit J.E. & Vyas P. (2010) Color Atlas of Clinical Hematology, 4th edn. Reproduced with permission of John Wiley & Sons.

Table 17.2 Specialized tests for acute lymphoblastic leukaemia (ALL).


<i>Immunological markers</i> (flow cytometry)	See Table 17.3; Fig. 11.14
<i>Immunoglobulin and TCR genes</i>	B-ALL: clonal rearrangement of immunoglobulin genes T-ALL: clonal rearrangement of TCR genes
<i>Chromosomes and genetic analysis</i>	See Table 17.1
B-ALL, B-cell acute lymphoblastic leukaemia; T-ALL, T-cell acute lymphoblastic leukaemia; TCR, T-cell receptor.	

with >20% leukaemic blasts. The blast cells are characterized by morphology (Fig. 17.4), immunological tests (Table 17.3) and cytogenetic analysis (Table 17.1). Identification of the immunoglobulin or T-cell receptor (TCR) gene rearrangement (Table 17.2), (aberrant) immunophenotype and molecular genetics of the tumour cells is important to determine treatment and to detect minimal residual disease (MRD) during follow-up.


Lumbar puncture for cerebrospinal fluid (CSF) examination is not generally performed as it may promote the spread of tumour cells to the CNS. Initial assessment of the CSF should always be combined with the concurrent administration of intrathecal chemotherapy. Biochemical tests may reveal a raised serum uric acid, serum lactate dehydrogenase or, less commonly, hypercalcaemia. Liver and renal function tests are performed as a baseline before treatment begins. Radiography may reveal lytic bone lesions and a mediastinal mass caused by enlargement of the thymus and/or mediastinal lymph nodes characteristic of T-ALL (Fig. 17.3).

Table 17.3 Immunological markers for classification of acute lymphoblastic (ALL) leukaemia (see also Fig. 11.14).

Marker	ALL	
	B	T
<i>B lineage</i>		
CD19	+	–
cCD22	+	–
cCD79a	+	–
CD10	+ or –	–
clg	+ (pre-B)	–
slg	–	–
TdT	+	+
<i>T lineage</i>		
CD7	–	+
cCD3	–	+
CD2	–	+
TdT	+	+
c, cytoplasmic; s, surface.		



(a)



(b)

Figure 17.3 Chest X-ray of a boy aged 16 years with acute lymphoblastic leukaemia (T-ALL). (a) There is a large mediastinal mass caused by thymic enlargement at presentation. (b) After 1 week of therapy with prednisolone, vincristine and daunorubicin the mass has resolved.

The differential diagnosis includes acute myeloid leukaemia (AML), aplastic anaemia (with which ALL sometimes presents), marrow infiltration by other malignancies (e.g. rhabdomyosarcoma, neuroblastoma and Ewing's sarcoma), infections such as infectious mononucleosis and pertussis, juvenile rheumatoid arthritis and immune thrombocytopenic purpura.

Cytogenetics and molecular genetics

Cytogenetic analysis shows differing frequencies of abnormalities in infants, children and adults which partly explains the different prognoses of these groups (Fig. 17.5). Cases are




Figure 17.4 Morphology, and immunophenotyping of acute lymphoblastic leukaemia (ALL). **(a)** Lymphoblasts show scanty cytoplasm without granules. **(b)** Lymphoblasts are large and heterogeneous with abundant cytoplasm. **(c)** Lymphoblasts are deeply basophilic with cytoplasmic vacuolation. **(d)** Acute lymphoblastic leukaemia: bone marrow cells staining positive for TdT by immunoperoxidase. Source: Hoffbrand A.V., Pettit J.E. & Vyas P. (2010) Color Atlas of Clinical Hematology, 4th edn. Reproduced with permission of John Wiley & Sons.




Figure 17.5 Cytogetic subsets of acute lymphoblastic leukaemia (ALL). The incidence of different cytogenetic abnormalities in infants, children and adults.

stratified according to the number of chromosomes in the tumour cell (**ploidy**) or by specific molecular abnormalities. The two parameters define good- and poor-prognosis disease.

Hyperdiploid cells have more than 50 chromosomes and generally have a good prognosis whereas **hypodiploid** cases (less than 44 chromosomes) carry a poor prognosis. The most common specific abnormality in childhood B-ALL is the t(12;21)(p13;q22) *ETV6-RUNX1* translocation. The RUNX1 protein plays an important part in transcriptional control of haemopoiesis and is repressed by the ETV6-RUNX1 fusion protein.

The frequency of the Ph translocation t(9;22) increases with age and carries a poor prognosis (although improving with the addition of BCR-ABL1 kinase inhibitors to therapy). Translocations of chromosome 11q23 involve the *MLL* gene and are seen particularly in cases of infant leukaemia. Using more sensitive molecular genetic tests, as well as fluorescence

in situ hybridization (FISH) analysis, some cases, normal by conventional cytogenetic testing, are found to have fusion genes, e.g. *BCR-ABL1* or a Ph-like pattern of kinase and cytokine expression without the *BCR-ABL1* fusion gene or other genetic abnormalities. These molecular genetic changes carry prognostic significance whether or not a corresponding chromosomal change is present.

T-ALL accounts for 15% of childhood and 25% of adult ALL and the clinical picture is often dominated by a very high white cell count, mediastinal mass or pleural effusion. TCR (and in 20% the *IGH* gene) show clonal rearrangement. Cytogenetic changes often involve the TCR loci with different partner genes. The majority of cases have acquired genetic abnormalities that lead to constitutive activation of the *NOTCH* signalling pathway and drugs that target these abnormalities are being developed (Fig. 17.6).




Figure 17.6 The molecular basis of activation of NOTCH signalling in T-ALL. **(a)** The molecular basis of NOTCH signalling. NOTCH is expressed at the cell membrane and after binding to a ligand (Delta-like or Jagged) on a neighbouring cell, the protein is cleaved in two places – first by extracellular ADAM 10 and then by an intracellular γ -secretase complex. The portion of intracellular NOTCH that is released is then translocated to the nucleus where it leads to activation of NOTCH1 target genes. **(b)** Several types of genetic abnormalities are seen in the NOTCH signalling pathway in patients with T-ALL. These include (1) mutations in the extracellular cleavage site, (2) insertion of an internal tandem duplications in the juxtamembrane region or (3) deletion of the intracellular PEST domain. The net result of all these mutations is to increase the rate of cleavage and nuclear translocation of the NOTCH domain.

Treatment

This may be conveniently divided into *supportive* and *specific* treatment.

General supportive therapy

General supportive therapy for bone marrow failure is described in Chapter 12 and includes the insertion of a central venous cannula, blood product support and prevention of tumour lysis syndrome. The risk of tumour lysis syndrome is highest in children with a high white cell count, T-cell disease or concurrent renal impairment at presentation. Any episode of fever must be treated promptly.

Specific therapy of ALL in children

Specific therapy of ALL is with chemotherapy and sometimes radiotherapy (Fig. 17.6), and treatment protocols are extremely complex. There are several phases in a treatment course, which usually has four components (Fig. 17.7). The protocols are *risk adjusted* to reduce the treatment given to patients with good prognosis. The factors that guide treatment include age, gender and white cell count at presentation. The initial response to therapy is also important, as slow clearance of blood or marrow blasts after a week or two of induction therapy or persistence of MRD (see below) is associated with a relatively high risk of relapse. ALL in infants (<1 year) has a worse clinical outcome with cure rates of only 20–50%. The disease is associated with chromosomal translocation involving the *MLL* gene in 80% of cases and is treated by unique protocols.

Minimal residual disease

Even when the blood and bone marrow appear to be clear of leukaemia, small numbers of tumour cells may sometimes be detected by fluorescence activated cell sorter (FACS) analysis or molecular methods (Fig. 17.8). A positive result indicates *minimal residual disease* (MRD) and the analysis of children for the presence of MRD at day 29 or adults after 3 months of treatment has prognostic significance and is now being used in planning therapy (see below) (Fig. 17.9). The value of tests for MRD at the end of induction or during consolidation continues to be explored in trials, in which the intensity of consolidation or maintenance therapy is reduced in those who rapidly become MRD negative, whereas more intensified therapy, or even allogeneic stem cell transplantation, is given to those with persistent MRD. Good-prognosis children become MRD negative as early as day 29 of therapy whereas in adults the finding at 3 months has greater prognostic importance (Table 17.4). One example being the randomised reduction in the number of intensification blocks in children with no evidence of MRD at day 29 within the current UK ALL 2003 trial. This reduction has not impaired the excellent prognosis in these low-risk children and is now standard therapy.




Figure 17.7 Flow chart illustrating typical treatment regimen of acute lymphoblastic leukaemia (ALL).

Remission induction

At presentation, the patient with acute leukaemia has a very high tumour burden and is at great risk from the complications of bone marrow failure and leukaemic infiltration (Fig. 17.1). The aim of *remission induction* is to rapidly kill most of the tumour cells and get the patient into remission. This is defined as less than 5% blasts in the bone marrow, normal peripheral blood count and no other symptoms or signs of the disease. Steroids (dexamethasone or prednisolone), vincristine and asparaginase are the drugs usually used and they are very effective – achieving remission in over 95% of children and in 80–90% of adults (in whom daunorubicin is also usually added). In remission, however, a patient may still be harbouring large numbers of tumour cells and, without further chemotherapy, virtually all patients will relapse (see Fig. 13.8). Nevertheless, achievement of remission is a valuable first step in the treatment course. Patients who fail to achieve remission need to change to a more intensive protocol.

Intensification (consolidation)

These courses use high doses of multidrug chemotherapy in order to eliminate the disease or reduce the tumour burden to very low levels. The doses of chemotherapy are near the limit of patient tolerability and during intensification blocks patients may need a great deal of support. Typical protocols involve the use of vincristine, cyclophosphamide, cytosine arabinoside, daunorubicin, etoposide or mercaptopurine given as blocks in different




Figure 17.8 Detection of minimal residual disease (MRD) by four-colour flow cytometry in: normal bone marrow mononuclear cells (BM), BM from a patient with B lineage ALL at diagnosis and in remission 6 weeks after diagnosis. The cells were detected with four different antibodies (anti-CD10, anti-CD19, anti-CD34, anti-CD38) attached to fluorescent labels abbreviated as PE, APC, PerCP and FITC, respectively. The tridimensional plot shows the immunophenotype of CD19⁺ lymphoid cells in the three samples. MRD of 0.03% of cells expressing the leukaemia-associated phenotype (CD10⁺, CD34⁺, CD38⁻) were detected at 6 weeks, confirmed by polymerase chain reaction (PCR) analysis. Source: Campana D. and Coustan-Smith E. (1999) *Cytometry. Commun Clin Cytometry* 38: 139–52. Reproduced with permission of John Wiley & Sons.

combinations. The numbers of blocks given in children depends on the child's risk category and varies between one and three.

Central nervous system-directed therapy

Few of the drugs given systemically are able to reach the CSF and specific treatment is required to prevent or treat central nervous system (CNS) disease. Intrathecal methotrexate is usually used. CNS relapses still occur and present with headache, vomiting, papilloedema and blast cells in the CSF. Treatment is with

intrathecal methotrexate, cytosine arabinoside and hydrocortisone, with or without cranial irradiation and systemic chemotherapy because bone marrow disease is usually also present.

Maintenance

This is given with daily oral mercaptopurine and once-weekly oral methotrexate. Intravenous vincristine with a short course (5 days) of oral dexamethasone is added at monthly or 3-monthly (in adults) intervals. There is a high risk of varicella or measles




Figure 17.9 Cumulative incidence of relapse according to minimal residual disease (MRD) levels at the end of remission induction in children with acute lymphoblastic leukaemia (ALL) treated at St Jude Children's Research Hospital. Source: Courtesy of Dr D. Campana.

Table 17.4 Prognosis in acute lymphoblastic leukaemia (ALL).

	Good	Poor
WBC	Low	High (e.g. $>30 \times 10^9/L$ B-ALL, $>100 \times 10^9/L$ T-ALL)
Sex	Girls	Boys
Immunophenotype	B-ALL	T-ALL (in children)
Age	Child	Adult (or infant <1 year)
Cytogenetics	Normal or hyperdiploidy; EVI6 rearrangement	t(9;22), 11q23 (MLL) rearrangements Hypodiploidy (<44 chromosomes)
Time to clear blasts from blood	<1 week	>1 week
Time to remission	<4 weeks	>4 weeks
CNS disease at presentation	Absent	Present
Minimal residual disease (MRD)	Negative at 1 month (children); 3 months (adults)	Still positive at 3–6 months
CNS, central nervous system; WBC, white blood cell count.		

during maintenance therapy in children who lack immunity to these viruses. If exposure to these infections occurs, prophylactic immunoglobulin should be given. In addition, oral co-trimoxazole is given to reduce the risk of *Pneumocystis jirovecii*. Maintenance may in some protocols begin soon after remission induction and is interrupted by the periods of intensification chemotherapy. In typical regimens intensive chemotherapy finishes around 38 weeks and then maintenance begins again and lasts until week 112 in girls or 164 in boys.

Treatment of relapse

If relapse, detected by MRD tests or by reappearance of ALL cells in the blood or marrow, occurs during or soon after maintenance chemotherapy the outlook is poor. Re-induction with combination chemotherapy, including novel drugs such as clofarabine, may help. Another approach to targeting B cells is to develop an antibody against a B cell-associated protein such as CD19, clone the immunoglobulin genes from this antibody and then combine these with a signalling molecule and transfer the chimeric gene into T cells taken from the patient. These T cells are then programmed to kill B cells and this approach of using 'chimeric antigen receptors' (CAR) is proving of value for patients with refractory ALL or CLL. Anti-CD22 monoclonal antibodies epratuzumab or inotuzumab may also help. Chemotherapy is usually followed, where possible, by allogeneic SCT. If relapse occurs after years off all therapy, the outlook is better and reinduction, consolidation and maintenance therapy are given. Allogeneic SCT may also be indicated. MRD levels are also proving valuable in defining risk and directing therapy in this group.

Toxicity

The long-term toxicity of therapy has always been a major concern of those treating children. The increasing success in terms of disease cure means that the mortality and morbidity

of therapy assumes a larger percentage of treatment failures. Specific concerns are the high risk of avascular bone necrosis seen in teenagers and young adults in association with dexamethasone, the potential long-term cardiac risk of even modest anthracycline doses, the impact of alkylating agents on fertility and the small risk of second tumours. The thrust of new-generation clinical trials in children is very much about reducing the risk of toxicity whilst preserving antileukaemic activity.

Specific therapy of ALL in adults

Treatment for ALL in adults has proven challenging compared to the great successes that have been observed in childhood therapy. The initial control of the leukaemia (remission induction) is comparable in both groups but the rate of disease relapse is much higher in adults. Cranial prophylaxis in adults usually involves intrathecal methotrexate, with or without cytosine arabinoside and steroids, and high-dose methotrexate and cytosine arabinoside given systemically. Although cure rates in children now approach 90%, no more than 40% of adult patients remain free of leukaemia after 5 years and this rate is much lower in older patients. A significant factor is that the genetic subtypes of the ALL differ according to age. Hyperdiploidy and t(12;21), which carry a good prognosis and together make up 50% of childhood cases, are both rare in adult patients. In contrast, the presence of the Ph⁺ disease becomes more common with age (Fig. 17.5).

An additional factor that has contributed to the relatively poor outcome for ALL in adults is the lower doses of chemotherapy that have traditionally been used in adult patients. This is now being addressed in younger adult patients where many are now being treated within 'childhood' protocols with high-intensity chemotherapy regimens. The presence of MRD after 3 months or more of therapy in adults is an unfavourable prognostic sign. Many patients are treated by allogeneic SCT if a suitable sibling or matched unrelated donor is available.

Treatment of BCR-ABL1 positive ALL

The introduction of the specific tyrosine kinase inhibitor imatinib has transformed the management of patients with BCR-ABL1+ (Ph+) ALL. Imatinib is used in combination with chemotherapy and is able to obtain remission of the disease in most patients. However, relapse is common because of the appearance of resistant subclones containing mutations in the *BCR-ABL1* gene. All-ogeneic SCT is therefore recommended wherever possible once

remission has been obtained. Second generation TKIs are now in use but their long-term value is uncertain at present.

Prognosis

There is a great variation in the chance of individual patients achieving a long-term cure based on a number of biological variables (Table 17.4). Approximately 25% of children relapse after first-line therapy and need further treatment but overall 90%




Figure 17.10 Overall survival rates for **(a)** adult and **(b)** paediatric ALL. Source: Courtesy of Dr Adele Fielding, Professor Ajay Vora and the UK NCRI Childhood Leukaemia Sub-group.

of children can expect to be cured (Fig. 17.10). The cure rate in adults drops significantly to less than 5% over the age of 70 years. The challenge in both adults and children over the next few years is to better define risk groups and individualise therapy.


This may be accomplished using both existing MRD technology and also newer assessment of molecular patterns; these patterns may, for example, define subgroups with specific drug sensitivities or children at particular risk of a specific drug toxicity.

SUMMARY

- Acute lymphoblastic leukaemia (ALL) is caused by an accumulation of lymphoblasts in the bone marrow. It is the most common malignant disease of childhood – 75% of cases occur before the age of 6 years. Eighty-five per cent of cases are of B-cell lineage with the rest being of T-cell lineage.
- The first genetic mutation occurs in many cases *in utero*, with a secondary genetic event occurring later in childhood, possibly as a reaction to an infection.
- The clinical presentation is with the features of bone marrow failure (anaemia, infection and bleeding) together with symptoms of tissue infiltration by tumour cells, leading to bone pain, enlarged lymph nodes, meningeal and testicular infiltration in some cases.
- Diagnosis is by examination of blood and bone marrow. Important tests include microscopic examination of the tumour cells, immunophenotyping and genetic analysis.
- ALL is subclassified according to the underlying genetic defect and a wide variety of genetic lesions are seen. The number of chromosomes in the tumour cell has prognostic importance: *Hyperdiploid* cells have >50 chromosomes and generally have a good prognosis whereas *hypodiploid* cases (<44 chromosomes) carry a poor prognosis. Other cytogenetic abnormalities are also of prognostic significance, e.g. t(9;22), increasingly frequent with age, which is unfavourable.
- Treatment protocols for ALL are extremely complex and usually have four components – remission induction, intensification, CNS-directed therapy and maintenance.
- Treatment is *risk adjusted* to reduce the treatment given to patients with good prognosis. This is based on age, gender, white cell count and cytogenetics at presentation.
- Small numbers of tumour cells may sometimes be detected by FACS or molecular analysis even when the blood and bone marrow appear to be clear of leukaemia. This *minimal residual disease* has prognostic significance and is used in planning therapy.
- If relapse occurs during chemotherapy the outlook is poor but if it happens after years off all treatment the outlook is better. Further chemotherapy and allogeneic SCT should be considered.
- Overall, >85% of children can now expect to be cured. The cure rate in adults drops significantly to less than 5% over the age of 70 years.



Now visit www.wileyessential.com/haematology to test yourself on this chapter.


A vertical strip on the left side of the page showing a microscopic view of blood cells. It includes several large, bright red erythrocytes (red blood cells) and some smaller, pinkish, irregularly shaped cells, possibly leukocytes or platelets, against a dark blue background.

CHAPTER 18

The chronic lymphoid leukaemias

Key topics

■ B-cell diseases	198
■ Chronic lymphocytic leukaemia	198
■ Pathogenesis	198
■ Clinical features	198
■ Laboratory findings	199
■ Treatment	200
■ Hairy cell leukaemia	202
■ T-cell diseases	203



Several disorders are included in this group and are characterized by accumulation in the blood of mature lymphocytes of either B- or T-cell type (Table 18.1). In general, these diseases are incurable but tend to run a chronic and fluctuating course.

Diagnosis

This group is characterized by a chronic persistent lymphocytosis. Subtypes are distinguished by morphology, immunophenotype and genetic analysis. There is some overlap with the non-Hodgkin lymphomas as lymphoma cells may be found in the blood and the distinction between chronic leukaemia and lymphoma can be somewhat arbitrary, depending on the relative proportion of the disease in soft tissue masses compared to blood and bone marrow.

B-cell diseases

Chronic lymphocytic leukaemia

Pathogenesis

Chronic lymphocytic leukaemia (CLL) is the most common of the chronic lymphoid leukaemias and has a peak incidence between 60 and 80 years of age. The aetiology is unknown but there are geographical variations in incidence. It is the most common form of leukaemia within Europe and the USA but less frequent elsewhere. There is a sevenfold increased risk of CLL in the close relatives of patients, which indicates a genetic predisposition to the disease, although the genes that carry this risk are largely unknown.

The CLL tumour cell is a mature B cell with weak surface expression of immunoglobulin (IgM or IgD). CLL cells typically exhibit impaired apoptosis and a prolonged lifespan, and this is reflected in their accumulation in the blood, bone marrow, liver, spleen and lymph nodes. Small lymphocytic lymphoma (SLL) (see Chapter 20) is the tissue equivalent of CLL and lymphoma cells have the same immunophenotype and cytogenetics as CLL. The difference is that in SLL the tumour cells accumulate almost exclusively in the lymph node and there are fewer than $5 \times 10^9/\text{L}$ circulating monoclonal B cells.

Table 18.1 Classification of the chronic lymphoid leukaemias.

B-cell	T-cell
Chronic lymphocytic leukaemia (CLL)	Large granular lymphocytic leukaemia
Prolymphocytic leukaemia (PLL)	T-cell prolymphocytic leukaemia (T-PLL)
Hairy cell leukaemia (HCL)	Adult T-cell leukaemia/lymphoma
Plasma cell leukaemia	
	Sézary syndrome (see Chapter 20)

Source: WHO (2008) classification (see Appendix).

The cytogenetic and molecular genetic abnormalities that may be present at diagnosis are listed below. Multiple clones from linear or branching evolution may be present at diagnosis. At later stages, after chemotherapy one or other resistant subclone may come to dominate (see Chapter 11).

Monoclonal B-cell lymphocytosis Clonal B cells with the same phenotype as CLL are found at low levels in the blood of many older patients. Indeed, this monoclonal B-cell lymphocytosis (MBL) has been demonstrated in 3% of patients over the age of 50 years and it is believed that all cases of clinical CLL progress from this state. Similar genetic changes to these found in CLL may be present. If CLL is to be diagnosed there must be a monoclonal B-cell count of $>5 \times 10^9/\text{L}$ or tissue involvement outside the bone marrow.

Clinical features

- 1 The mean age at diagnosis is 72 years, with only 15% of cases before 50 years of age. The male:female ratio is approximately 2:1.
- 2 Over 80% of cases are diagnosed from the results of a routine blood test, usually taken for another reason.
- 3 Enlargement of cervical, axillary or inguinal lymph nodes is the most frequent clinical sign (Fig. 18.1). The nodes are usually discrete and non-tender.
- 4 Features of anaemia may be present and patients with thrombocytopenia may show bruising or purpura.
- 5 Splenomegaly and, less commonly, hepatomegaly are common in later stages.
- 6 Immunosuppression is often a significant problem resulting from hypogammaglobulinaemia and cellular immune dysfunction. Early in the disease course bacterial infections, such as sinus and chest infections, predominate but with advanced disease viral infections, especially herpes



Figure 18.1 Chronic lymphocytic leukaemia: bilateral cervical lymphadenopathy in a 67-year-old woman. Haemoglobin 125 g/L; white blood count $150 \times 10^9/\text{L}$ (lymphocytes $146 \times 10^9/\text{L}$); platelets $120 \times 10^9/\text{L}$.




Figure 18.2 Chronic lymphocytic leukaemia: herpes zoster infection in a 68-year-old female.




Figure 18.3 Chronic lymphocytic leukaemia: peripheral blood film showing lymphocytes with thin rims of cytoplasm, coarse condensed nuclear chromatin and rare nucleoli. Typical smudge cells are present.

zoster (Fig. 18.2), and fungal infections are also seen. Patients should be vaccinated with a conjugated pneumococcal vaccine.

Laboratory findings

- 1 Lymphocytosis. The absolute clonal B cell lymphocyte count is $>5 \times 10^9/L$ and may be up to $300 \times 10^9/L$ or more. Between 70 and 99% of white cells in the blood film appear as small lymphocytes. 'Smudge' or 'smear' cells are also present (Fig. 18.3).
- 2 Immunophenotyping of the lymphocytes shows them to be B cells (surface $CD19^+$) with low levels of surface immunoglobulin and expression of only one light chain (known as 'light chain restriction'; see Figure 18.4). Characteristi-

cally, the cells are also surface $CD5^+$ and $CD23^+$ but are $CD79b^-$ and $FMC7^-$.

- 3 Two surface proteins that can be detected by flow cytometry and have prognostic significance are $CD38$, a marker of differentiation, and $ZAP70$, a protein kinase involved in signalling (Table 18.2).
- 4 Normochromic normocytic anaemia is present in later stages as a result of marrow infiltration or hypersplenism. Autoimmune haemolysis may also occur (see below). Thrombocytopenia occurs in many patients and may also have an autoimmune basis.
- 5 Bone marrow aspiration shows up to 95% lymphocytic replacement of normal marrow elements. Trephine biopsy reveals nodular, diffuse or interstitial involvement by lymphocytes (Fig. 18.4).




Figure 18.4 Chronic lymphocytic leukaemia: trephine biopsies showing: (a) a marked diffuse increase in marrow lymphocytes (closely packed cells with small dense nuclei); (b) a nodular pattern of lymphocyte accumulation (in a different patient); and (c) interstitial infiltration.

Table 18.2 Immunophenotype of the chronic B-cell leukaemias/lymphomas (all cases CD19+).

	CLL	HCL	FL	MCL
SIg	Weak	++	++	+
CD5	+	–	–	+
CD22/FMC7	–	+	+	++
CD23	+	–	–	–
CD79b	–	–/+	++	++
CD103*	–/+	+	–	–

CLL, chronic lymphocytic leukaemia; FL, follicular lymphoma; HCL, hairy cell leukaemia; MCL, mantle cell lymphoma SIg, surface immunoglobulin.

* CD103 is positive only in HCL.

- 6 Reduced concentrations of serum immunoglobulins are found and this becomes more marked with advanced disease. Rarely, a paraprotein is present.
- 7 Autoimmunity directed against cells of the haemopoietic system is common. Autoimmune haemolytic anaemia is most frequent but immune thrombocytopenia, neutropenia and red cell aplasia are also seen.

Molecular tests

Genetics and molecular genetics

The most common chromosomal abnormalities are deletion of 13q14, trisomy 12, deletion at 11q23 and 17p deletion. These abnormalities carry prognostic significance for some treatment regimens but are becoming less important as treatments become more powerful (Table 18.2). The 13q14 deletion leads to loss of microRNAs (see p. 130), which normally control expression of proteins that regulate B-cell survival. The most common genetic point mutations at presentation are found in *ATM*, *NOTCH1* and *SF3B1* (all at around 10% prevalence), *BIRC3* (4%) and *p53* (<5%) which also have prognostic value (Table 18.4).

Somatic hypermutation of the immunoglobulin genes

When B cells recognize antigen in the germinal centre of secondary lymphoid tissues they undergo a process called **somatic hypermutation** in which random mutations occur in the immunoglobulin heavy-chain gene (see Chapter 9). In CLL the *IGHV* gene shows evidence of this hypermutation in approximately 50% of cases whereas in the other cases the *VH* genes are unmutated. CLL with unmutated immunoglobulin genes has an unfavourable prognosis (Table 18.4).

Staging

It is useful to stage patients at presentation both for prognosis and for deciding on therapy. The **Rai and Binet staging systems** are shown in Table 18.3. Typical survival ranged from 12 years for Rai stage 0 to less than 4 years for stage IV, but

there is considerable variation between patients, and with current therapies survival rates are improving. Many patients in stage 0 have a normal life expectancy.

Treatment

It is very difficult to cure CLL and so the approach to therapy is generally conservative, aiming for symptom control rather than a normal blood count. Indeed, chemotherapy given too early in the disease can shorten rather than prolong life expectancy. Another important fact is that many patients never need treatment. Treatment is given for troublesome enlarged lymph nodes or spleen, and constitutional symptoms such as weight loss or bone marrow suppression. The lymphocyte count alone is not a good guide to treatment but if it doubles in <6 months, treatment will usually be needed soon. As a general guide, patients in Binet stage C will need treatment as will some in stage B.

Chemotherapy

For many years the treatment for CLL has based on the combination of cytotoxic drugs (such as fludarabine, chlorambucil and cyclophosphamide or bendamustine) together with a monoclonal antibody against CD20, such as rituximab. However, there are now several effective new drugs which may replace these regimens. These are discussed below in the section *New agents in the treatment of B-CLL*.

In 2015, the most commonly used first-line *treatment for younger patients* is R-FC, which combines the antibody rituximab (anti-CD20; see p. 221) with fludarabine and cyclophosphamide. These agents are given together every 4 weeks and are able to control the white cell count and reduce organ swelling in most cases. Four to six courses are usually given and treatment can be stopped after a satisfactory response has been achieved. The average 'time to disease progression' after treatment with R-FC is around 4.5 years. This regimen has a number of potential side-effects, including myelosuppression and immunosuppression. R-B, using bendamustine with rituximab is an alternative approach which is less immunosuppressive.

Table 18.3 Staging of chronic lymphocytic leukaemia (CLL).**(a) Rai classification***Stage*

0	Absolute lymphocytosis $>5 \times 10^9/L^+$
I	As stage 0 + enlarged lymph nodes (adenopathy)
II	As stage 0 + enlarged liver and/or spleen $\pm$ adenopathy
III	As stage 0 + anaemia (Hb $<100 g/L$) ⁺ $\pm$ adenopathy $\pm$ organomegaly
IV	As stage 0 + thrombocytopenia (platelets $<100 \times 10^9/L$) ⁺ $\pm$ adenopathy $\pm$ organomegaly

(b) International Working Party classification (Binet)

<i>Stage</i>	<i>Organ enlargement*</i>	<i>Haemoglobin[†] (g/L)</i>	<i>Platelets[†] ($\times 10^9/L$)</i>
A (50–60%)	0, 1 or 2 areas	≥ 100	≥ 100
B (30%)	3, 4 or 5 areas		
C (<20%)	Not considered	<100	and/or <100

* One area = lymph nodes >1 cm in neck, axillae, groins or spleen, or liver enlargement.

[†] Secondary causes of anaemia (e.g. iron deficiency) or autoimmune haemolytic anaemia or autoimmune thrombocytopenia must be treated before staging.

Source: (b) Adapted from Binet J.L. *et al.* (1981) *Cancer* 48: 198.

In older or less fit patients a less intensive treatment plan is used, which still involves chemotherapy and an anti-CD20 antibody. Chlorambucil is an oral alkylating agent and is often used in this setting, although bendamustine is also possible.

Alemtuzumab is an anti-CD52 monoclonal antibody that is effective at killing B and T lymphocytes by complement fixation but leads to serious infectious complications. It has been used in resistant and relapsed disease but is being replaced by the newer agents.

New Agents in the treatment of B-CLL

In recent years several new and highly effective therapies have emerged in the treatment of lymphoid disorders. They are used for treating relapsed, resistant patients but trials of their use as initial therapy are in progress.

1 Drugs which suppress signalling through the B cell receptor (see Chapter 9). The surface immunoglobulin on a B cell acts as the B cell receptor (BCR) for antigen and tumour cells need to receive stimulatory signals through the BCR in order to remain alive. Two drugs have been synthesized that block two different signalling proteins that act 'downstream' of the BCR.

- **Inactivation of the BTK protein** BTK deficiency is a rare inherited disorder of childhood characterised by susceptibility to chest infections due to a B cell lymphopenia and is due to inheritance of inactivating mutations within the *BTK* gene. This reveals how important BTK must be for B cell survival. **Ibrutinib** is an oral drug which inactivates BTK and leads to B cell apoptosis. It is highly effective in the treatment of CLL, either as a single agent or in combination with monoclonal antibodies. Remarkably it is effective for cases in which the tumour carries a 17p deletion or p53 mutation. Increased susceptibility to bleeding is a potential side-effect.

- **Inactivation of the PI3KD protein** Phosphoinositide 3-kinase (PI3K) is also a critical enzyme in BCR signalling (see Figure 9.4) and expression of the delta subunit is relatively specific for lymphoid cells. **Idelalisib** is an oral drug which blocks PI3K Δ activity which is also very effective in CLL. It is well tolerated although colitis can occur. A feature of both drugs is that they initially, like steroids, cause an increase in the lymphocyte count in peripheral blood. This reflects a redistribution from lymph nodes and marrow. For this reason anti-CD20 monoclonal antibody is often given initially with these drugs.

2 Drugs which suppress the activity of BCL-2. BCL-2 is expressed at a high level in most CLL cells and has potent anti-apoptotic effects. ABT-199 is a direct inhibitor of BCL-2 and shows highly potent activity in early phase trials.

3 New effective monoclonal antibodies include ofatumumab and obinutuzumab, which are CD20-specific.

Other forms of treatment

- **Corticosteroids** Prednisolone is given in autoimmune haemolytic anaemia, thrombocytopenia and red cell aplasia. High-dose steroid therapy alone with alemtuzumab was used in patients with 17q deletion, resistant to chemotherapy. Benefit was temporary and serious infections frequent.
- **Radiotherapy** This is valuable in reducing the size of bulky lymph node groups that are unresponsive to chemotherapy. Radiotherapy to the spleen may be valuable in late-stage disease.
- **Lenalidomide** is a thalidomide derivative and has therapeutic activity in CLL. Initial treatment is sometimes associated with a disease 'flare' at affected tissue sites and the mechanism of action is uncertain.
- **Ciclosporin** Red cell aplasia may respond to ciclosporin.

- **Immunoglobulin replacement** Immunoglobulin (e.g. 400 mg/kg/month by intravenous infusion) is useful for patients with hypogammaglobulinaemia and recurrent infections, especially during winter months.
- **Stem cell transplantation** This is currently an experimental approach in younger patients. Allogeneic stem cell transplantation (SCT) may be curative but has a significant mortality rate.

Course of disease

Many patients in Binet stage A or Rai stage 0 or I never need therapy and this is particularly likely for those with favourable prognostic markers (Table 18.4). For those who do need treatment a typical pattern is that of response to several courses of chemotherapy before the gradual onset of extensive bone marrow infiltration, bulky disease and recurrent infection. Molecular and cytogenetic tests show that initially small subclones with, e.g. 17p deletion or P53 mutations, now form the bulk of this resistant disease. The new oral therapies are proving effective even at these late stages. The disease may also transform into a high-grade lymphoma (Richter's transformation) which then requires therapy as for other high-grade B-cell lymphomas.

B-cell prolymphocytic leukaemia

Although prolymphocytic leukaemia (PLL) may initially appear similar to CLL, the diagnosis is made by the appearance of a majority of prolymphocytes in the blood. The prolymphocyte is around twice the size of a CLL lymphocyte and has a large central nucleolus (Fig. 18.5). PLL typically presents with splenomegaly without lymphadenopathy and with a high and rapidly rising lymphocyte count. Anaemia is a poor prognostic feature. Treatment is difficult but splenectomy, chemotherapy and anti-CD20 may help.




Figure 18.5 Prolymphocytic leukaemia: blood film showing prolymphocytes that have prominent central nucleoli and an abundance of pale cytoplasm.

Hairy cell leukaemia

Hairy cell leukaemia (HCL) is an uncommon B-cell lymphoproliferative disease with a male:female ratio of 4:1 and a peak incidence at 40–60 years. Patients typically present with infections, anaemia or splenomegaly. Lymphadenopathy is uncommon. Pancytopenia is usual at presentation and the lymphocyte count is rarely over $20 \times 10^9/L$. Monocytopenia is a distinctive feature. The blood film reveals a variable number of unusual large lymphocytes with villous cytoplasmic projections (Fig. 18.6). Immunophenotyping shows CD11c, CD19, CD25, CD103 and CD123 positivity in most cases (see Table 20.3a). **A mutation in exon 15 of the gene for the protein kinase *BRAF* underlies the disease.** The bone marrow trephine shows a characteristic appearance of mild fibrosis and a diffuse cellular infiltrate (Fig. 18.6).

Table 18.4 Prognostic factors in chronic lymphocytic leukaemia.

	Good	Bad
Stage	Binet A (Rai 0–I)	Binet B, C (Rai II–IV)
Lymphocyte doubling time	Slow	Rapid
Bone marrow biopsy appearance	Nodular	Diffuse
Chromosomes	Deletion 13q14	Deletion 17p
Genetic mutations		<i>NOTCH</i> , <i>SF3B1</i> , <i>p53</i>
VH immunoglobulin genes	Hypermutated	Unmutated Use of VH3.21
ZAP expression	Low	High
CD38 expression	Negative	Positive
LDH	Normal	Raised
LDH, lactate dehydrogenase.		




Figure 18.6 Hairy cell leukaemia: **(a)** peripheral blood film showing typical 'hairy' cells with oval nuclei and finely mottled pale grey-blue cytoplasm with an irregular edge; **(b)** bone marrow trephine.

There are several effective treatments for HCL and the median relapse-free survival after chemotherapy is 16 years. The treatment of choice is 2-chlorodeoxyadenosine (CDA) or deoxycoformycin (DCF) and both agents achieve complete responses in over 80% of cases. In two-thirds of cases a long-term remission is achieved. α -interferon is also effective and is sometimes used first for those with severe cytopenias to improve these before the more effective drugs, which initially tend to lower the neutrophil and platelet counts, are given. Rituximab can be combined with CDA or DCF for relapsed cases. BRAF inhibitors are potentially useful in rare cases of refractory disease.

Lymphocytosis in non-Hodgkin lymphomas

Some cases of splenic marginal zone lymphoma show circulating monoclonal B lymphocytes with a villous cell outline and were previously termed 'splenic lymphoma with villous lymphocytes' (see p. 222). Lymphocytosis may also be seen

in other types of non-Hodgkin lymphoma (e.g. follicular, mantle cell, diffuse large B cell) and are discussed further in Chapter 20.

T-cell diseases

T-cell prolymphocytic leukaemia

This presents as B-PLL with a high white cell count, but lymphadenopathy is more marked and skin lesions and serous effusions are common. Most tumours express CD4⁺. Treatment is with alemtuzumab followed by SCT in appropriate patients.

Large granular lymphocytic leukaemia

Large granular lymphocytic leukaemia (LGL-L) is characterized by the presence of circulating lymphocytes with abundant cytoplasm and large azurophilic granules (Fig. 18.7a). Such cells may be either T or natural killer (NK) cells and show variable




Figure 18.7 **(a)** Large granular lymphocytes in the peripheral blood. **(b)** Adult T-cell leukaemia/lymphoma. Typical convoluted lymphoid cells in peripheral blood.

expression of CD16, CD56 and CD57. *STAT3* mutations are present in 50% of cases. Cytopenia, especially neutropenia, is the main clinical problem, although anaemia, splenomegaly and arthropathy with positive serology for rheumatoid arthritis are also common. The mean age is 50 years. Treatment may not be needed, but, if required, steroids, cyclophosphamide, ciclosporin or methotrexate may relieve the cytopenia. Granulocyte colony-stimulating factor (G-CSF) has been used in cases associated with neutropenia.

Adult T-cell leukaemia/lymphoma

Adult T-cell leukaemia/lymphoma (ATLL) was the first malignancy to be associated with a human retrovirus, human T-cell leukaemia/lymphoma virus type 1 (HTLV-1). The

virus is endemic in parts of Japan and the Caribbean and the disease is very rare in people who have not lived in these areas. ATLL lymphocytes have a bizarre morphology with a convoluted 'clover-leaf' nucleus and a consistent CD4⁺ phenotype (Fig. 18.7b).


Most subjects infected with the virus do not develop the disease. The clinical presentation is often acute and dominated by hypercalcaemia, skin lesions, hepatosplenomegaly and lymphadenopathy. Diagnosis is by morphology and serology. Zidovudine, an anti-retroviral drug, and alpha interferon are first-line therapy if leukaemia is dominant but combination chemotherapy is used if the presentation is more like a lymphoma. The prognosis is poor.

SUMMARY

- Chronic lymphocytic leukaemias are characterized by the accumulation of mature B or T lymphocytes in the blood.
- Individual subtypes are distinguished on the basis of morphology, immunophenotype and cytogenetics.
- Chronic lymphocytic leukaemia (CLL, B cell) represents 90% of cases and has a peak incidence between 60 and 80 years of age. There is genetic predisposition to development of the disease.
- Most cases are identified when a routine blood test is performed. As the disease progresses the patient can develop enlarged lymph nodes, splenomegaly and hepatomegaly.
- Immunosuppression is a significant problem because of hypogammaglobulinaemia and cellular immune dysfunction.
- Anaemia may also develop because of autoimmune haemolysis and bone marrow infiltration.
- Diagnosis is usually performed by immunophenotypic analysis of peripheral blood which reveals a clonal population of CD5⁺ CD23⁺ B cells.
- The best guide to prognosis is the stage of the disease. CLL that has acquired somatic mutations in the immunoglobulin genes has a relatively good prognosis compared to unmutated cases. Cytogenetics also provide prognostic information.
- Treatment is usually given only when clinical symptoms start to develop. Use of chemotherapy and an anti-CD20 monoclonal antibody establishes disease remission but is not curative.
- Oral drugs which block the activity of BTK (ibrutinib) or PI3KΔ (idelalisib) are new highly effective treatments even in otherwise resistant patients.
- Less common subtypes of chronic lymphoid leukaemias include prolymphocytic leukaemia, hairy cell leukaemia and T-cell disorders.



Now visit www.wileyessential.com/haematology to test yourself on this chapter.


A vertical strip on the left side of the page showing a microscopic view of blood cells. It features large, bright red erythrocytes and smaller, pinkish leukocytes against a dark blue background.

CHAPTER 19

Hodgkin lymphoma

Key topics

■ History and pathogenesis	206
■ Clinical features	206
■ Haematological and biochemical findings	206
■ Diagnosis and histological classification	207
■ Clinical staging	208
■ Treatment	210
■ Prognosis	212
■ The late effects of Hodgkin lymphoma and its treatment	212



Lymphomas are a group of diseases caused by malignant lymphocytes that accumulate in lymph nodes and other lymphoid tissue and cause the characteristic clinical feature of lymphadenopathy. Occasionally, they may spill over into blood ('leukaemic phase') or infiltrate organs outside the lymphoid tissue.

The major subdivision of lymphomas is into **Hodgkin lymphoma (HL) and non-Hodgkin lymphoma (NHL)** and this is based on the histological presence of **Reed–Sternberg (RS) cells in Hodgkin lymphoma**.

History and pathogenesis

Thomas Hodgkin was curator of the Anatomy Museum at Guy's Hospital in London and described the disease in 1832. Dorothy Reed and Carl Sternberg were pathologists who identified the abnormal cell that defines this subtype of lymphoma in 1898. The characteristic RS cells, and the associated abnormal mononuclear cells, are neoplastic, whereas the infiltrating inflammatory cells are reactive. Immunoglobulin gene rearrangement studies suggest that the RS cell is of B-lymphoid lineage and that it is often derived from a B cell with a 'crippled' immunoglobulin gene caused by the acquisition of mutations that prevent synthesis of full-length immunoglobulin. HLA class I expression is usually lost on the tumour cells and mutation of the $\beta 2$ -microglobulin gene is frequent. The Epstein–Barr virus (EBV) genome has been detected in over 50% of cases in Hodgkin tissue but its exact role in the pathogenesis is unclear.

Clinical features

The disease can present at any age but is rare in children and has a peak incidence in young adults. There is an almost 2:1 male predominance. The following symptoms are common.

- 1 Most patients present with painless, asymmetrical, firm and discrete enlargement of superficial lymph nodes (Fig. 19.1).** The cervical nodes are involved in 60–70% of patients, axillary nodes in approximately 10–15% and inguinal nodes in 6–12%. In some cases the size of the nodes decreases and increases spontaneously and they may become matted. Typically the disease is localized initially to a single peripheral lymph node region and its subsequent progression is by contiguity within the lymphatic system. Retroperitoneal nodes are also often involved but usually only diagnosed by computed tomography (CT) scan.
- 2 Modest splenomegaly occurs during the course of the disease in 50% of patients.** The liver may also be enlarged because of liver involvement.
- 3 Mediastinal involvement is found in up to 10% of patients at presentation.** This is a feature of the nodular sclerosing type, particularly in young women. There may be associated pleural effusions or superior vena cava obstruction (Fig. 19.2).





Figure 19.1 Cervical lymphadenopathy in a patient with Hodgkin lymphoma.


- 4 Cutaneous Hodgkin lymphoma occurs as a late complication in approximately 10% of patients.** Other organs may also be involved, even at presentation, but this is unusual.
- 5 Constitutional symptoms are prominent in patients with widespread disease.** The following may be seen:
 - (a) fever** occurs in approximately 30% of patients and is continuous or cyclic;
 - (b) pruritus**, which is often severe, occurs in approximately 25% of cases;
 - (c) alcohol-induced pain** in the areas where disease is present occurs in some patients;
 - (d) other constitutional symptoms** include weight loss, profuse sweating (especially at night), weakness, fatigue, anorexia and cachexia. Haematological and infectious complications are discussed below.

Haematological and biochemical findings

- 1 Normochromic normocytic anaemia is most common.** Bone marrow involvement is unusual in early disease, but if it occurs bone marrow failure may develop with a leucoerythroblastic anaemia.
- 2 One-third of patients have a neutrophilia;** eosinophilia is frequent.
- 3 Advanced disease is associated with lymphopenia and loss of cell-mediated immunity.**



(a)



(b)

Figure 19.2 (a) Chest X-ray in Hodgkin lymphoma showing enlargement of left hilar lymph nodes, abnormal soft tissue projected over the upper left lung and a large left pleural effusion. (b) Axial CT scan with intravenous contrast in the same patient. There is a large anterior mediastinal mass (yellow circle) with left hilar nodes enlarged (red circle) and left pleural effusion. Source: Courtesy of Dr Peter Wylie and Dr N. Nir.

- 4 The platelet count is normal or increased during early disease, and reduced in later stages.
- 5 The erythrocyte sedimentation rate (ESR) and C-reactive protein are usually raised. The ESR is useful in monitoring disease progress.

6 Serum lactate dehydrogenase (LDH) is raised initially in 30–40% of cases.

7 The HIV status should be determined.

Diagnosis and histological classification

The diagnosis is made by histological examination of an excised lymph node. The distinctive multinucleate polyploid RS cell is central to the diagnosis of the four classic types (Figs 19.3 and 19.4) and mononuclear Hodgkin cells are also part of the malignant clone. These cells stain with CD30 and CD15 but are usually negative for B-cell antigen expression. Inflammatory components consist of lymphocytes, neutrophils, eosinophils, plasma cells and variable fibrosis. CD68 detects infiltrating macrophages and if this is strongly positive it is an unfavourable feature.

Histological classification is into four classical types and nodular lymphocyte predominant disease (Table 19.1). There is no difference in the prognosis or management of the different sub-types of classical HL. Nodular sclerosis is the most frequent in Europe and the USA, whereas lymphocyte depleted is more common in developing countries and has a particularly strong association with EBV infection. Nodular lymphocyte predominant does not show RS cells and has many features of non-Hodgkin lymphoma and may be treated as such.




Figure 19.3 Diagrammatic representation of the different cells seen histologically in Hodgkin lymphoma.




Figure 19.4 Hodgkin lymphoma: **(a)** high power view of a lymph node biopsy showing two typical multinucleate Reed–Sternberg cells, one with a characteristic owl eye appearance, surrounded by lymphocytes, histiocytes and an eosinophil; **(b)** mixed cellularity; and **(c)** nodular sclerosing Hodgkin lymphoma.

Clinical staging

The disease may be suspected on the basis of clinical presentation, chest X-ray and CT scan (Fig. 19.2).

The selection of appropriate treatment depends on accurate staging of the extent of disease (Table 19.2). Figure

19.5 shows the scheme that is used. Staging is performed by clinical examination together with **combined positron emission tomography (PET) and CT scans**. CT scan alone can be used if PET is not available (Figs 19.5 and 19.6). Magnetic resonance imaging (MRI) scanning may be needed for particular sites (Table 19.2). Bone marrow trephine is

Table 19.1 WHO (2008) classification of Hodgkin lymphoma.	
Classical Hodgkin lymphoma (95% of cases)	
Nodular sclerosis	Collagen bands extend from the node capsule to encircle nodules of abnormal tissue. A characteristic lacunar cell variant of the Reed–Sternberg cell is often found. The cellular infiltrate may be of the lymphocyte-predominant, mixed cellularity or lymphocyte-depleted type; eosinophilia is frequent
Lymphocyte rich	Scanty Reed–Sternberg cells; multiple small lymphocytes with few eosinophils and plasma cells; nodular and diffuse types
Mixed cellularity	The Reed–Sternberg cells are numerous and lymphocyte numbers are intermediate
Lymphocyte depleted	There is either a reticular pattern with dominance of Reed–Sternberg cells and sparse numbers of lymphocytes or a diffuse fibrosis pattern where the lymph node is replaced by disordered connective tissue containing few lymphocytes. Reed–Sternberg cells may also be infrequent in this latter subtype
Nodular lymphocyte-predominant (5% of cases)	
Reed–Sternberg cells are absent; lymphocyte predominant (LP) tumour B cells are present	

Table 19.2 Techniques for staging of lymphoma.

Laboratory	Full blood count ESR Bone marrow aspirate and trephine (not routine) Liver function LDH C-reactive protein
Radiology	Chest X-ray PET/CT of thorax, abdomen, chest and pelvis MRI Bone scan (rare)

CT, computed tomography; ESR, erythrocyte sedimentation rate; LDH, lactate dehydrogenase; MRI, magnetic resonance imaging; PET, positron emission tomography.

sometimes carried out and liver biopsy may also be needed in difficult cases. **PET/CT scanning is useful in monitoring response to treatment and for detection of small foci of residual disease (Fig. 19.7).**

Patients are also classified as A or B according to whether or not constitutional features (fever or weight loss) are present (Fig. 19.5).

Positron emission tomography (PET)

^{18}F -fluorodeoxyglucose positron emission tomography (FDG-PET) is now used widely in the management of lymphoma and other haematological malignancies. It utilizes the fact that rapidly dividing malignant cells readily take up glucose from their environment. Radiolabelled glucose is infused into the




Figure 19.5 Staging of Hodgkin lymphoma. Stage I indicates node involvement in one lymph node area. Stage II indicates disease involving two or more lymph nodal areas confined to one side of the diaphragm. Stage III indicates disease involving lymph nodes above and below the diaphragm. Splenic disease is included in stage III but this has special significance (see below). Stage IV indicates involvement outside the lymph node areas and refers to diffuse or disseminated disease in the bone marrow, liver and other extranodal sites. NB. The stage number in all cases is followed by the letter A or B indicating the absence (A) or presence (B) of one or more of the following: unexplained fever above 38°C ; night sweats; or loss of more than 10% of body weight within 6 months. Localized extranodal extension from a mass of nodes does not advance the stage but is indicated by the subscript E. Thus, mediastinal disease with contiguous spread to the lung or spinal theca would be classified as I_{E} . As involvement of the spleen is often a prelude to widespread haematogenous spread of the disease, patients with lymph node and splenic involvement are staged as III_{S} . Bulky disease (widening of the mediastinum by more than one-third, or the presence of a nodal mass $>10\text{ cm}$ in diameter) is relevant to therapy at any stage.




Figure 19.6 Hodgkin lymphoma. Staging PET/CT: 35-year-old female who had disease above and below the diaphragm at presentation. **(a)** Coronal PET image shows multiple foci of uptake above and below the diaphragm. **(b)** Coronal fused PET/CT scan image shows multiple foci of uptake above and below the diaphragm corresponding to nodes, spleen and lung nodules. PET stage IV. Source: Courtesy Dr Thomas Wagner and the Department of Nuclear Medicine, Royal Free Hospital, London.

patient and the tissues that have taken up the label can then be visualized by the PET scanner.

As well as detecting the presence of active disease at the time of diagnosis, PET/CT scans can also be used to assess the response to treatment and to potentially guide the treatment course. These ‘interim PET/CT’ scans are reported according to the **Deauville 5 point criteria**, which uses the uptake in the mediastinum and liver as an internal control, from which to assess the activity of the tumour. Scores 1 and 2 are generally considered ‘negative’, whereas 4 and 5 are ‘positive’ and a score of 3 must be interpreted according to the clinical context.

- Score 1 no uptake
- Score 2 uptake ≤ mediastinum
- Score 3 uptake > mediastinum but ≤ liver
- Score 4 moderately increased uptake > liver
- Score 5 markedly increased uptake > liver

Treatment

Treatment is with chemotherapy alone or a combination of chemotherapy with radiotherapy. The choice depends primarily on the stage, clinical state A or B (Fig. 19.5) and prognostic factors (Table 19.3). Semen storage, if appropriate, should

be carried out before therapy is begun. For females it is advisable that fertility advice is sought from a specialist. If blood

Table 19.3 Prognosis of stage 1–2 disease (EORTC criteria).

Favourable	Unfavourable
	Any one of:
No large mediastinal lymphadenopathy*	Large mediastinal lymphadenopathy
ESR <50 without B symptoms	ESR > or = 50 without B symptoms
ESR<30 with B symptoms	ESR <or =30 with B symptoms
Age < or = 50 years	Age >50 years
1-3 lymph node sites involved	4 or more lymph node sites involved

*Large is defined as mediastinal thoracic ratio >0.35 at the level of T5/6. EORTC, European Organization for the Research and Treatment of Cancer.




Figure 19.7 Example of the value of imaging in the management of Hodgkin lymphoma. (a) Axial PET, (b) fused PET/CT and (c) CT images at diagnosis demonstrate intense [18 F]FDG uptake in an anterior mediastinal mass. Following two cycles of ABVD chemotherapy (d) the axial PET, (e) fused PET/CT and (f) CT images demonstrate no significant [18 F]FDG uptake in the residual mediastinal mass, in keeping with a complete metabolic response. Source: Courtesy of Dr V.S. Warbey and Professor G.J.R. Cook.

transfusion is needed this must be irradiated to avoid graft-versus-host disease due to the infusion of live lymphocytes which can engraft due to the impaired cellular immunity of the HL patient.

Early stage disease

The outcome for early stage disease is excellent and an important aim is to avoid over-treatment and the risk of late complications. Two broad options are chemotherapy alone or 'combined modality treatment' (CMT) using chemotherapy and radiotherapy. CMT achieves better short-term disease control but in the longer term there is no clear increase in overall survival. Individual treatment decisions will depend on local regimens and patient choice. As examples, favourable prognosis disease could be treated with two courses of A (Adriamycin = Doxorubicin), B (Bleomycin), V (Vinblastine), D (Dacarbazine) (ABVD) chemotherapy followed by 20 Gy radiotherapy. If lymph nodes are not bulky radiotherapy can be omitted but at least three courses of ABVD given.

In contrast, unfavourable disease (1B or 2B) could be treated with four to six courses of ABVD followed by 30 Gy radiotherapy for bulky disease. Alternatively the first two cycles of ABVD can be replaced by more intensive chemotherapy, such as escalated BEACOPP (Bleomycin, Etoposide, Adriamycin, Cyclophosphamide, Vincristine = Oncovin, Procarbazine and Prednisolone).

Advanced stage disease

Cyclical chemotherapy is used for stage III and IV disease. Six to eight cycles of ABVD are most widely used. Six cycles of escalated BEACOPP may be given to achieve higher complete

remission rates at the expense of greater toxicity. Subsequent radiotherapy is given if residual nodes are more than 1.5 cm diameter, or smaller but remain PET positive, or to sites of originally bulky disease.

Assessment of response to treatment

Clinical examination and imaging (PET/CT scans) are used to assess response to treatment. Regular pulmonary function assessment is needed, in older patients and those receiving bleomycin. Patients with HL often show residual masses following treatment which may be because of the large degree of fibrosis present within lymph nodes. PET/CT scanning reveals the areas of active disease (Fig. 19.7). Clinical trials are assessing if PET/CT can be used to define the management of individual patients. For example, it may be used after the first two cycles of ABVD and if there is residual active disease, treatment might be switched to more intensive chemotherapy. Alternatively, if the PET scan is negative, bleomycin might be omitted from subsequent chemotherapy and there is generally no need to repeat the PET/CT scan at the end of therapy. For patients PET/CT-positive at the end of therapy, repeat biopsy and close clinical and imaging assessment is needed. Inflammatory tissue may cause a false positive PET scan.

Relapsed cases

Approximately 25% of patients suffer from disease relapse or are refractory to initial therapy. Treatment is generally given as an alternative combination chemotherapy to the initial regimen and, if necessary, with radiotherapy to sites of bulky disease. Brentuximab vedotin is an anti-CD30 antibody linked to a microtubule-disrupting agent and may produce favourable responses. If




Figure 19.8 Potential mechanism whereby Hodgkin lymphoma is controlled after treatment with antibodies to block PD-1. PD-1 and PD-L1 are natural molecules that limit the attack of normal tissues by cytotoxic T cells. Hodgkin lymphoma overexpresses PD-L1 due to gene amplification or effects of EBV infection. This delivers a strong negative signal to the T cells around the tumour. If antibody-mediated blockade of PD-1 is used, the T cells can then recognise and kill the tumour.

the disease remains chemosensitive, high-dose chemotherapy and autologous stem cell transplantation improve the probability of cure and are given for most patients below the age of 65 years. Allogeneic transplantation may also be curative in a minority of patients who fail other therapies. A new therapy is the use of antibodies which block the inhibitory molecule PD-1 on T cells (Fig. 19.8). Hodgkin lymphoma often expresses high levels of the PD-1 ligand PD-L1 and this acts as a mechanism for evading the T cell immune response. PD-1 blockade is proving to be highly effective in management of relapsed HL and is now likely to be investigated earlier in the treatment pathway.

Prognosis

The prognosis depends on age, stage and histology. Overall approximately 85% of patients are cured.

The late effects of Hodgkin lymphoma and its treatment


Long-term follow-up of patients has revealed a considerable burden of late disease following treatment. Secondary cancers, such as lung cancer and breast cancer, appear to be related to radiotherapy, whereas myelodysplasia or acute myeloid leukaemia are more associated with the use of alkylating agents. Non-Hodgkin lymphoma and other cancers also occur with greater frequency than in controls. Non-malignant complications include sterility, intestinal complications, coronary artery disease and pulmonary complications of the mediastinal radiation or bleomycin chemotherapy. Vinblastine may cause a permanent neuropathy. These features are the main reason why less intensive treatment regimens are now being explored for this disease.

SUMMARY

- Lymphomas are a group of diseases caused by malignant lymphocytes that accumulate in lymph nodes and cause lymphadenopathy.
- The major subdivision of lymphomas is into Hodgkin lymphoma and non-Hodgkin lymphoma and this is based on the presence of Reed–Sternberg cells in Hodgkin lymphoma.
- Reed–Sternberg cells are neoplastic B cells but most cells in the lymph node are reactive inflammatory cells.
- The usual clinical presentation is with painless asymmetrical lymphadenopathy – most commonly in the neck.
- Constitutional symptoms of fever, weight loss and sweating are prominent in patients with widespread disease.
- Blood tests may show anaemia, neutrophilia and raised erythrocyte sedimentation rate (ESR) or lactate dehydrogenase (LDH).
- Diagnosis is made by histological examination of an excised lymph node and there are four subtypes of disease.
- Staging of the disease is important for determining treatment and prognosis. History, examination, blood tests, CT and PET scan are typically used.
- Treatment is with radiotherapy, chemotherapy or a combination of both. The choice depends on the stage and grade of the disease.
- The response to treatment can be monitored by CT and PET scans. Disease relapse can be treated with chemotherapy, sometimes with stem cell transplantation.
- The prognosis is excellent and over 85% of patients can expect to be cured. Late side-effects of treatment are a concern.



Now visit www.wileyessential.com/haematology to test yourself on this chapter.




CHAPTER 20

Non-Hodgkin lymphoma

Key topics

■ Introduction to non-Hodgkin lymphoma	214
■ Clinical features of non-Hodgkin lymphoma	216
■ Laboratory investigation	216
■ Specific subtypes of non-Hodgkin lymphoma (NHL)	220
■ Lymphoplasmacytoid NHL	221
■ Marginal zone NHL	221
■ Follicular NHL	222
■ Mantle cell NHL	223
■ Diffuse large B-cell NHL	223
■ Burkitt lymphoma	224
■ T-cell lymphomas	225



Introduction to non-Hodgkin lymphoma

These are a large group of clonal lymphoid tumours, about 85% of B cell and 15% of T or NK (natural killer) cell origin (Table 20.1). Their clinical presentation and natural history are much more variable than Hodgkin lymphoma. They are characterized by an irregular pattern of spread and a significant proportion of patients develop disease outside the lymph nodes. Their frequency has increased markedly over the last 50 years and, with an incidence of approximately 17 in 100 000, they now represent the fifth most common malignancy in some developed countries (see Fig 11.2).

Classification

The lymphomas are classified within a group of **mature B-cell and T-cell neoplasms**, which also includes some chronic leukaemias and myeloma, which are described in Chapters 18 and 21, respectively (Table 20.1). The World Health Organization (WHO) classification recognizes age (paediatric or elderly) and site of involvement (e.g. skin, central nervous system (CNS), intestine, spleen, mediastinal) as well as the disease histology, immunophenotype and genotype important in disease classification.

In this chapter we consider the more common lymphoma subtypes within this classification (Fig. 20.1).

Cell of origin

The normal B-cell development stages are illustrated in Fig. 9.11. B-cell lymphomas tend to mimic normal B cells at different stages of development (Fig. 20.2). This is shown by

their phenotypic patterns on immune histology or flow cytometry (Fig. 20.3a). They can be divided into those resembling precursor B cells found in the bone marrow, and those which resemble germinal centre (GC) cells or post-GC cells from lymph nodes. T-cell lymphomas resemble precursor T cells in the bone marrow and thymus, or peripheral mature T cells.

Low- and high-grade non-Hodgkin lymphoma

The non-Hodgkin lymphomas (NHL) are a diverse group of diseases and vary from highly proliferative and potentially rapidly fatal diseases to some very indolent and well-tolerated malignancies. For many years clinicians have subdivided lymphomas into ‘low-grade’ and ‘high-grade’ disease. This approach is valuable as, in general terms, the low-grade disorders are relatively indolent, respond well to chemotherapy but are very difficult to cure, whereas high-grade lymphomas are aggressive and need urgent treatment but are more often curable.

Leukaemias and lymphomas

The difference between *lymphomas*, in which lymph nodes, spleen or other solid organs are involved, and *leukaemias*, with predominant bone marrow and circulating tumour cells, may be blurred. Chronic lymphocytic leukaemia and small lymphocytic lymphoma are identical lymphoproliferative diseases but show leukaemic and lymph node distribution respectively. Acute lymphoblastic leukaemia and acute lymphoblastic lymphoma are also similar and have comparable treatment regimens. Small numbers of malignant cells circulate in the blood in many forms of NHL.

Table 20.1 The World Health Organization (WHO) classification of mature B-cell and T-cell neoplasms (modified), which includes the non-Hodgkin lymphomas. B-cell disorders comprise 85% of cases. T cell and NK cell together comprise 15% of cases. A few rare subtypes have been omitted (see Appendix).

Mature B-cell neoplasms	Mature T-cell and NK-cell neoplasms
Chronic lymphocytic leukaemia/small lymphocytic lymphoma	T-cell prolymphocytic leukaemia
B-cell prolymphocytic leukaemia	T-cell large granular lymphocytic leukaemia
Splenic marginal zone lymphoma	Adult T-cell lymphoma/leukaemia
Hairy cell leukaemia	Extranodal NK/T-cell lymphoma, nasal type
Lymphoplasmacytic lymphoma – Waldenström	Enteropathy-associated T-cell lymphoma
macroglobulinaemia	Mycosis fungoides
Heavy chain diseases	Sézary syndrome
Plasma cell myeloma	Peripheral T-cell lymphoma
Plasmacytoma	Angioimmunoblastic T-cell lymphoma
Extranodal marginal zone lymphoma of mucosa-associated lymphoid tissue (MALT lymphoma)	Anaplastic large cell lymphoma, ALK positive
Follicular lymphoma	
Mantle cell lymphoma	
Diffuse large B-cell lymphoma	
Burkitt lymphoma	

ALK, anaplastic lymphoma kinase, the gene on chromosome 2 which is overexpressed; NK, natural killer.




Figure 20.1 The relative frequencies of B-cell non-Hodgkin lymphomas. CLL, chronic lymphocytic lymphoma; DLBCL, diffuse large B-cell lymphoma; MALT, mucosa-associated lymphoid tissue; MCL, mantle cell lymphoma; NOS, not otherwise specified; PMLBCL, primary mediastinal large B-cell lymphoma; SLL, small lymphocytic lymphoma.




Figure 20.2 Proposed cellular origin of B-lymphoid malignancies. Normal B cells migrate from the bone marrow and enter secondary lymphoid tissue. When they encounter antigen a germinal centre is formed and B cells undergo somatic hypermutation of the immunoglobulin genes. Finally, B cells exit the lymph node as memory B cells or plasma cells. The cellular origin of the different lymphoid malignancies can be inferred from immunoglobulin gene rearrangement status and membrane phenotype. Mantle cell lymphoma and a proportion of B-cell chronic lymphocytic lymphoma (B-CLL) cases have unmutated immunoglobulin genes, whereas marginal zone lymphoma, diffuse large cell lymphoma, follicle cell lymphoma, lymphoplasmacytoid lymphoma and some B-CLL cases have mutated immunoglobulin genes.




Figure 20.3 Non-Hodgkin lymphoma: histological sections of lymph nodes showing: **(a)** a diffuse pattern of involvement in lymphocytic lymphoma with the normal architecture totally replaced by neoplastic lymphocytic cells; **(b)** a follicular or nodular pattern in follicular lymphoma – the ‘follicles’ or ‘nodules’ of neoplastic cells compress surrounding tissue and lack a mantle of small lymphocytes.

Pathogenesis

The aetiology of the majority of cases of NHL is unknown although infectious agents are an important cause in particular subtypes (Table 20.2). There are also considerable geographical variations (Table 20.2). Cytogenetic abnormalities are frequent, often involving the immunoglobulin genes in the B-cell neoplasms. Translocations of oncogenes to these loci on chromosomes 2, 14 and 22 may result in overexpression of the gene leading to alteration of the cell cycle, failure of apoptosis

Table 20.2 Infections associated with haemopoietic malignancies.

Infection	Tumour
<i>Virus</i>	
HTLV-1	Adult T-cell leukaemia/lymphoma
Epstein–Barr virus	Burkitt and Hodgkin lymphomas; PTLD
HHV-8	Primary effusion lymphoma; multicentric Castleman's disease
HIV-1	High-grade B-cell lymphoma, primary CNS lymphoma, Hodgkin lymphoma
Hepatitis C	Marginal zone lymphoma
<i>Bacteria</i>	
<i>Helicobacter pylori</i>	Gastric lymphoma (MALT)
<i>Protozoa</i>	Burkitt lymphoma
Malaria	

HHV-8, human herpes virus 8; HIV, human immunodeficiency virus; HTLV-1, human T-lymphotropic virus type 1; MALT, mucosa-associated lymphoid tissue; PTLD, post-transplant lymphoproliferative disease.

or aberrant expression (see Chapter 11). Specific signalling pathways may be affected and next generation sequencing has revealed point mutations in genes involved in, for example, chromatin modelling, the NFkappa B pathway of B-cell activation and splicing. As many as 80 somatic mutations may be present in NHL at presentation (see Fig. 11.3) and further mutations may appear as the disease progresses.

Clinical features of non-Hodgkin lymphoma

- 1 Superficial lymphadenopathy** The majority of patients present with asymmetric painless enlargement of lymph nodes in one or more peripheral lymph node regions.
- 2 Constitutional symptoms** Fever, night sweats and weight loss occur less frequently than in Hodgkin lymphoma. Their presence is usually associated with disseminated disease.
- 3 Oropharyngeal involvement** In 5–10% of patients there is disease of the oropharyngeal lymphoid structures (Waldeyer's ring) which may cause complaints of a 'sore throat' or noisy or obstructed breathing.
- 4 Symptoms due to anaemia, infections** due to neutropenia or *purpura* with thrombocytopenia may be presenting features in patients with diffuse bone marrow disease. Cytopenias may also be autoimmune in origin or due to sequestration in the spleen.
- 5 Abdominal disease** The liver and spleen are often enlarged and involvement of retroperitoneal or mesenteric nodes is frequent. The gastrointestinal tract is the most commonly involved extranodal site after the bone marrow, and patients may present with acute abdominal symptoms.
- 6 Other organs** Involvement of the skin, brain, testis or thyroid is not infrequent. The skin is also primarily involved in two closely related T-cell lymphomas, mycosis fungoides and Sézary syndrome.

Investigations

Histology

Whole lymph node biopsy, or more usually trucut biopsy of lymph node or of other involved tissue (e.g. bone marrow or extranodal tissue), is the definitive investigation (Fig. 20.3). Fine needle aspiration of lymph node or involved tissue is rarely sufficient to establish a definitive diagnosis of lymphoma and for this reason is unreliable and a biopsy should be obtained.

Morphological examination of the biopsy is assisted by immunophenotypic and, in some cases, genetic analysis (Table 20.3). For B-cell lymphomas, expression of either κ or λ light chains confirms clonality and distinguishes the disease from a reactive node (Fig. 20.4).

Laboratory investigations

- 1** In advanced disease with marrow involvement there may be anaemia, neutropenia or thrombocytopenia.




Figure 20.4 Non-Hodgkin lymphoma: lymph node stained by immunoperoxidase shows (a) brown ring staining for κ in the malignant lymphoid nodule, and (b) no labelling for λ confirming the monoclonal origin of the lymphoma.

- 2 Lymphoma cells (e.g. mantle zone cells, 'cleaved follicular lymphoma' or 'blast' cells) may be found in the peripheral blood in some patients (Fig. 20.5).
- 3 Trepine biopsy of marrow is valuable (Fig. 20.6).
- 4 The serum lactate dehydrogenase (LDH) level is raised in rapidly proliferating and extensive disease and is used as a prognostic marker (Table 20.4). Elevation of serum uric acid may occur.
- 5 Immunoglobulin electrophoresis may reveal a paraprotein.
- 6 HIV status should be tested as indicated.

Cytogenetic and genetic analysis

The subtypes of NHL are associated with characteristic chromosomal translocations and genetic mutations which are of diagnostic and prognostic value (Table 20.3). Particularly characteristic translocations are t(14;18) in follicular lymphoma, t(11;14) in mantle cell lymphoma, t(8;14) in Burkitt lymphoma and t(2;5) in anaplastic large cell lymphoma. Genetic

Table 20.3 (a) Characteristic immunophenotype of common B-cell lymphomas.

	SIg	CD20	CD5	CD10	CD23	BCL6	MUM1
Small lymphocytic lymphoma/CLL	weak	+	+	–	+	–	–
Hairy cell leukaemia	+	+	–	–	–	–	–
Lymphoplasmacytic lymphoma	+	+	–	–	–	–	+
MALT lymphoma	+	+	–	–	+/-	–	+/-
Follicular lymphoma	+	+	–	+	+/-	+	–
Mantle cell lymphoma	+	+	+	–	–	–	–
Diffuse large B-cell lymphoma GCB	+/-	+	–	–		+	–
ABC	+/-	+	–	–		–	+
Burkitt lymphoma	+	+	–	+	–	+	–

CLL, chronic lymphocytic leukaemia; MALT, mucosa-associated lymphoid tissue; GCB, germinal centre B-cell type; ABC, activated B-cell type; MUM1 is a lymphocytic-specific transcription factor; SIg, surface immunoglobulin.

Table 20.3 (b) Examples of cytogenetic abnormalities and gene mutations found in cases of lymphoid malignancies.

	Cytogenetics	Gene mutations
Chronic lymphocytic leukaemia	13p, 11q and 17p deletions. Trisomy 12	<i>TP53</i> , <i>ATM</i> , <i>NOTCH</i> , <i>SF3B1</i>
Hairy cell leukaemia	Non-specific	<i>BRAF</i> (>99%)
Lymphoplasmacytic lymphoma	Non-specific	<i>MYD88</i> (>90%)
MALT lymphoma	t(11;18), t(1;14)	Activation of NF κ B pathway
Follicular lymphoma	t(14;18)	Mutations in genes such as <i>CREBBP</i> and <i>MLL2</i> which influence chromatin remodelling
Mantle cell lymphoma	t(11;14)	<i>ATM</i> , <i>TP53</i> and genes influencing chromatin modification
Diffuse large B cell lymphoma (germinal centre subtype)	t(14;18)	<i>CREBBP</i> (chromatin remodelling), <i>EZH2</i> and <i>MYC</i>
Diffuse large B cell lymphoma (activated B cell subtype)	t(3;14)	<i>CREBBP</i> (chromatin remodelling) and genes involved in activation of NF- κ B
Burkitt lymphoma	t(8;14), t(2;8), t(8;22)	Various




Figure 20.5 Blood involvement by malignant lymphoma: **(a)** small cleaved lymphoid cells in follicle centre cell lymphoma; **(b)** mantle cell lymphoma; **(c)** large B-cell lymphoma.




Figure 20.6 Iliac crest trephine biopsy in lymphocytic lymphoma. Prominent nodules of lymphoid tissue are seen in the intertrabecular space and paratrabecular areas.

Table 20.4 The National Comprehensive Cancer Network-International prognostic index (NCCN-IPI) for patients with high-grade lymphoma.

	Score
Age, y	
>40 to ≤60	1
>60 to ≤75	2
>75	3
LDH, normalized	
>1 to ≤3	1
>3	2
Ann Arbor stage III–IV	1
Extranodal disease*	1
Performance status ≥2	1

* Disease in bone marrow, CNS, liver/gastrointestinal tract, or lung.

analysis reveals mutation of *MYD88* in virtually all cases of lymphoplasmacytic lymphoma.

In B-cell lymphomas the immunoglobulin genes are clonally rearranged whereas in T-cell lymphomas there is clonal rearrangement of the T-cell receptor genes (see Chapter 11).

Staging

The staging system is the same as that described for Hodgkin lymphoma (see Chapter 19) but is less clearly related to prognosis than the histological subtype. Staging procedures usually include chest X-ray and PET/CT scanning (Fig. 20.7). PET/CT can also be used to follow treatment response (Fig. 20.8). Bone marrow aspiration and trephine are performed.




Figure 20.7 Non-Hodgkin lymphoma. **(a)** Computed tomography (CT) examination performed for weight loss and abdominal pain in a 86-year-old female showing enlarged para-aortic (yellow arrow) and mesenteric lymph nodes (blue circle). Histology revealed diffuse large B-cell non-Hodgkin lymphoma. Source: Courtesy of Dr P. Wylie. **(b)** CT scan of the abdomen: enlarged retroperitoneal and mesenteric nodes from a man causing the 'floating aorta' (arrowed) appearance. Source: Courtesy of Professor A. Dixon and Dr R.E. Marcus. **(c)** Magnetic resonance imaging (MRI) scan of the chest showing large mediastinal lymph nodes (white and arrowed) adjacent to the great vessels (black). **(d)** MRI T₂-weighted midline sagittal image of a lumbosacral spine showing compression of the dural sac by an extradural mass. A, spinal cord; B, extradural mass; C, roots of corda equina. Source: Courtesy of Dr A. Valentine. **(e)** Positron emission tomography (PET) body scan of a 59-year-old woman with high-grade non-Hodgkin lymphoma. (i) The first scan showed no evidence of disease prior to allogeneic transplant. Normal physiological uptake is seen in the brain and bladder. Two months post-transplant the patient relapsed clinically with a mass on the anterior chest wall. (ii) The PET scan showed evidence of widespread relapse in nodal (para-aortic and iliac nodes) and extranodal sites including the lung and bone. The uptake in bone is clearly demonstrated in the left humerus and femur (arrowed). This scan illustrates how well PET can detect both nodal and extranodal disease and allows whole body assessment at a single scanning session. Source: Courtesy of Dr S.F. Barrington.




Figure 20.8 Non-Hodgkin lymphoma. PET/CT scan of male aged 26 years. Red arrow shows the level at which transaxial section is performed. Upper right-hand panel showing FDG uptake in anterior mediastinal mass and right hilar node. Upper left-hand panel shows corresponding CT section, bottom left-hand panel shows the CT and PET fused image of the corresponding section. Bottom right-hand panel, coronal section showing FDG uptake in mediastinal mass and right hilar node. There is also FDG uptake in the thyroid, heart, bowel, kidneys and bladder. Source: Courtesy of the Department of Nuclear Medicine, University College London, London.

General principles of treatment of non-Hodgkin lymphoma

When treatment is started it is usually in the form of a combination of chemotherapy drugs together with a monoclonal antibody directed against the tumour cell. However, various new drugs have been developed which may change the management of disease in future years. These include:

- Oral agents that block activity of the BTK or PI3KD proteins, as used for B-CLL (see p. 201);
- Drugs that inhibit BCL-2 activity;
- Drugs that block activity of kinases, such as ALK which is increased in patients with anaplastic large cell lymphoma.

Monoclonal antibody therapy

Eighty-five per cent of cases of NHL are of B-cell origin and in these cases antibodies against CD20 have proven of great

benefit. Rituximab was the first such agent and can be given by intravenous or subcutaneous administration. Ofatumumab and obinutuzumab are also anti-CD20 specific antibodies. Antibodies against CD30 are often used for anaplastic large cell lymphoma as well as Hodgkin lymphoma (see Chapter 19).

Specific subtypes of non-Hodgkin lymphoma

Low-grade non-Hodgkin lymphoma

Small lymphocytic lymphoma

This term is used for cases with the same morphology and immunophenotype as B-cell chronic lymphocytic lymphoma (B-CLL) but with less than $5 \times 10^9/L$ peripheral blood B cells and no cytopenias due to bone marrow involvement. Treatment is as for patients with B-CLL.

Lymphoplasmacytoid lymphoma (Waldenström's macroglobulinaemia)

This is an uncommon condition, seen most frequently in men over 50 years of age. There is usually a monoclonal IgM paraprotein and the lymphoplasmacytoid lymphoma (LPL) may then be termed Waldenström's macroglobulinaemia. The cell of origin is a post-germinal centre B cell with the characteristics of an IgM-bearing memory B cell. A mutation of the gene *MYD88* underlies over 90% of all cases.

LPL may be diagnosed by chance in symptomless patients (Fig. 21.3). The disease usually presents with an insidious onset, often with fatigue and weight loss. Hyperviscosity syndrome (see Fig. 21.15) is a common complication as the IgM paraprotein increases blood viscosity more than equivalent concentrations of IgG or IgA. Visual upset is frequent and the retina may show a variety of changes, such as engorged veins, haemorrhages, exudates and a blurred disc (see Fig. 21.15). If the macroglobulin is a cryoglobulin, features of cryoprecipitation, such as Raynaud's phenomenon, may be present.

Anaemia is usually a significant problem and a bleeding tendency may result from macroglobulin interference with coagulation factors and platelet function. Neurological symptoms, dyspnoea and heart failure may be presenting symptoms. Moderate lymphadenopathy and enlargement of the liver and spleen are frequently seen.

Diagnosis is made by the finding of a monoclonal serum IgM together with bone marrow or lymph node infiltration with lymphoplasmacytoid cells (Fig. 20.9). **Mutation of *MYD88* is present in nearly all cases.** The erythrocyte sedimentation rate (ESR) is raised and there may be a peripheral blood lymphocytosis.

Treatment

No treatment is required for patients without symptoms, but treatment should be started if there are features such as




Figure 20.9 Lymphoplasmacytoid lymphoma associated with Waldenström's macroglobulinaemia. Bone marrow shows cells with features of lymphocytes and plasma cells.




Figure 20.10 The potential mechanisms of action of rituximab. Rituximab binds to CD20 on the surface of B cells. It can elicit a number of effector mechanisms including: (a) antibody-dependent cell-mediated cytotoxicity; (b) complement-mediated lysis of tumour cells; and (c) direct apoptosis of the target cell.

organomegaly, symptomatic anaemia or hyperviscosity. Combination treatment with an anti-CD20 antibody (such as Rituximab (Fig. 20.10)) and chemotherapy is generally given. Chemotherapy options include cyclophosphamide, fludarabine, bendamustine or bortezomib. Ibrutinib is effective in cases refractory to other drugs and may become initial therapy. Autologous or allogeneic stem cell transplantation (SCT) is considered for advanced disease. Erythropoietin or regular transfusions may be required for chronic anaemia.

Acute hyperviscosity syndrome is treated with repeated plasmapheresis. As IgM is mainly present in the intravascular space, plasmapheresis is more effective than with IgG or IgA paraproteins when much of the protein is extravascular and so rapidly replenishes the plasma compartment.

Marginal zone lymphomas

Marginal zone lymphomas are low-grade lymphomas that arise from the marginal zone of B cell germinal follicles. It is thought that lymphoid hyperplasia initially occurs in response to antigen or inflammation and then cells acquire secondary genetic damage that leads to lymphoma. Cytogenetic analysis may reveal translocations involving the immunoglobulin loci and molecular tests show point mutations particularly involving the NF- κ B pathway. They are classified according to the anatomical site at which they arise, such as the **spleen**, **mucosa (MALT)** or **lymph node (nodal)**. Mucosa-associated lymphoid tissue (MALT) lymphomas usually arise in the stomach (Fig. 20.11), respiratory tract, skin and salivary glands. Gastric MALT lymphoma is the most common form




Figure 20.11 Gastric mucosa-associated lymphoid tissue (MALT) lymphoma: the tumour cells surround reactive follicles and infiltrate the mucosa. The follicle has a 'starry sky' appearance. Source: Courtesy of Professor P. Isaacson.

and is preceded by *Helicobacter pylori* infection. In the early stages it may respond to antibiotic therapy aimed at eliminating *H. pylori*.

Splenic marginal zone lymphoma usually presents as splenomegaly and may be associated with circulating 'villous' lymphocytes. Localized stage Ia disease may be cured by local radiotherapy. If chemotherapy is needed it is usually based on regimens used in other low-grade lymphomas such as follicular lymphoma. Splenectomy may be useful for symptomatic patients.

Follicular lymphoma

This represents around 25% of NHL with a median age of onset of 60 years. It is associated with the t(14;18) translocation in the great majority of cases (Figs 20.12). The translocation leads to constitutive expression of the *BCL-2* gene with increased survival of cells because of reduced apoptosis. Additional molecular abnormalities are usually present (Table 20.3b). The cells are typically CD10, CD19, CD20, BCL2 and BCL6 positive (Table 20.3a).

Patients are likely to be middle-aged or elderly and their disease is often characterized by a benign course for many years. The median survival from diagnosis is approximately 10 years. The histological appearances are graded as I–III according to the relative proportion of centrocytes and centroblasts. Grade IIb patients have the worst prognosis and are generally treated according to diffuse large B-cell lymphoma (DLBCL) guidelines. Bone marrow involvement is frequent.

Presentation is usually with painless lymphadenopathy, often widespread, and the majority of patients will have stage III or IV disease. However, sudden transformation may occur at a rate of about 3% a year to aggressive diffuse tumours.

Around 10% of patients have initially localized (Stage I) disease and may achieve cure with radiotherapy alone. Those with disseminated (Stage II–IV) disease are generally not treated in the absence of symptoms ('watch and wait') but treatment is introduced when complications occur. An international prognostic score prognosis depends on age, presence or not of anaemia, serum LDH and disease burden as measured by size and extent of nodes and bone marrow involvement. At the current time, chemotherapy is not a curative option. Therapy is




Figure 20.12 Follicular lymphoma: immunostain. (a) The neoplastic cells are diffusely positive for B-cell markers (CD20). (b) Immunostain: the neoplastic cells are diffusely positive for CD10, a germinal centre marker, and are located in the follicular and interfollicular areas. (c) The neoplastic cells are positive for BCL-6, a germinal centre marker. (d) Immunostain: the neoplastic cells are positive for BCL-2.

usually based on monthly courses of rituximab, in combination with cyclophosphamide, vincristine and prednisolone (R-CVP) with an anthracycline added for more aggressive cases, or rituximab with bendamustine or chlorambucil. These regimens can provide clinical responses in up to 90% of patients and usually achieve a remission of several years. Rituximab infusions are given as maintenance therapy, usually every 2–3 months.

Disease relapse for stage II–IV disease is almost inevitable and is usually treated initially with similar chemotherapy regimens followed by rituximab maintenance. Over time the disease becomes increasingly difficult to control with, for example, P53 mutations, and more intensive chemotherapy or radiolabelled antibody therapy (anti-CD20) may be considered. Autologous SCT may be valuable in patients with a history of at least one relapse and allogeneic SCT using reduced intensity protocols offers the prospect of cure for some patients. Idelalisib, a PI3K inhibitor, and ibrutinib are promising new oral agents (see Fig. 9.4).

Mantle cell lymphoma

Mantle cell lymphoma is derived from pre-germinal centre cells localized in the primary follicles or in the mantle region of secondary follicles. The cells usually show angular nuclei in histological sections (Fig. 20.13) and often circulate in the blood (Fig. 20.5). It has a characteristic phenotype of CD19⁺ and CD5⁺ (like CLL) but, in contrast to CLL, is CD22⁺ and CD23⁻. A specific t(11;14) translocation juxtaposes the cyclinD1 gene to the immunoglobulin heavy-chain gene, and leads to **increased expression of cyclin D1** (Fig. 20.13b). Presence of this translocation is required for diagnosis. Other mutations are usually present (Table 20.3b). Clinical presentation is typically with lymphadenopathy and often there is bone marrow infiltration.

Current treatment regimens include:

- 1 Chemotherapy, such as R-CHOP (see DLBCL), bendamustine combinations or more intensive therapy, which may include bortezomib and lenalidomide.

2 Ibrutinib is very effective in relapsed and early disease.

3 Autologous or allogeneic stem cell transplantation.

The prognosis is generally quite poor but may improve with the new agents. Fifteen per cent of patients show an indolent course similar to that for CLL.

High-grade non-Hodgkin lymphoma

Diffuse large B-cell lymphomas (DLBCL)

DLBCL are a heterogeneous group of disorders representing the classic 'high-grade' lymphomas. Histologically the biopsy shows large tumour cells with prominent nucleoli. They are subdivided into 'germinal centre' (GCB) and 'activated B cell' (ACB) subtypes which stain with antibodies to BCL6 and MUM1 respectively (Table 20.3a, Fig. 20.14).

The clinical presentation is usually with rapidly progressive lymphadenopathy, which may also involve the bone marrow, gastrointestinal tract, brain (Fig. 20.15), spinal cord, kidneys or other organs.

A variety of clinical and laboratory findings are relevant to the outcome of therapy. According to the **International Prognostic Index (IPI)** these include age, performance status, stage, number of extranodal sites and serum LDH (Table 20.4). Bulky disease (major mass >5 cm diameter), prior history of low-grade disease or HIV infection, and ABC compared to GCB type are also associated with a poorer prognosis. There is a variety of histological patterns including centroblastic, immunoblastic and anaplastic. The most common cytogenetic changes involve the IGH locus on chromosome 14 and the *BCL-6* gene at chromosome 3q27 and translocation of the *BCL-2* gene occurs in 20%.

The mainstay of treatment is rituximab in combination with the CHOP chemotherapy regimen (cyclophosphamide, hydroxycarbonyl, vincristine (Oncovin) and prednisolone) given in 3-weekly cycles, typically for six to eight courses. Granulocyte colony-stimulating factor (G-CSF) injections are often used to support the neutrophil count. For localized disease, combined radiotherapy and chemotherapy (e.g. three




Figure 20.13 (a) Mantle cell lymphoma: showing characteristic deformed pattern of small lymphocytes with angular nuclei ('centrocytes'). (b) Mantle cell lymphoma: expression of cyclin D1 shown by immunohistochemistry. Source: Campo E. & Pileri S. A. In Hoffbrand A.V. *et al.* (eds) (2016) *Postgraduate Haematology* 7e. Reproduced with permission of John Wiley & Sons.




Figure 20.14 Diffuse large B-cell lymphoma **(a)** which stains positive for CD10 **(b)** and therefore suggests a germinal centre cell origin. Source: Campo E. & Pileri S. A. In Hoffbrand A.V. *et al.* (eds) (2016) *Postgraduate Haematology* 7e. Reproduced with permission of John Wiley & Sons.




Figure 20.15 Cerebral lymphoma in HIV infection; MRI. **(a)** T2-weighted magnetic resonance brain scan showing heterogeneous mass and adjacent oedema in right inferior frontoparietal region. There is compression of the right lateral ventricle and displacement of midline structures. Biopsy showed diffuse large B-cell lymphoma. **(b)** The mass enhances after intravenous gadolinium injection. **(c)** Enhanced image after chemotherapy showing regression of the mass. Source: Courtesy of the Department of Radiology, Royal Free Hospital, London.

courses of R-CHOP) may be optimal. Prophylactic therapy to prevent CNS disease, such as intrathecal or high-dose systemic methotrexate, should be considered for patients with high-risk disease, particularly those with bone marrow involvement. The response to treatment is monitored by repeat CT or PET/CT scans midway through chemotherapy and then following completion.

For patients who relapse, high-dose chemotherapy with drug regimens such as R-ICE (rituximab, ifosfamide, carboplatin and etoposide) can be effective. In those patients who

respond to these treatments, autologous SCT is often used. Reduced-intensity allogeneic SCT has also been shown to be effective. However, for those with primary refractory or chemoresistant disease the outlook is poor. Overall, long-term survival is approximately 65%.

Burkitt lymphoma

Burkitt lymphoma occurs in endemic or sporadic forms. Endemic (African) Burkitt lymphoma is seen in areas with



Figure 20.16 Burkitt lymphoma: characteristic facial swelling caused by extensive tumour involvement of the mandible and surrounding soft tissues.

chronic malaria exposure and is associated with Epstein–Barr virus (EBV) infection. In virtually all cases the *MYC* oncogene is overexpressed because it is translocated to an immunoglobulin gene, usually the heavy-chain locus t(8;14) (see Fig. 11.11). As a result, the gene is expressed in parts of the cell cycle during which it should normally be switched off.

Typically the patient, usually a child, presents with massive lymphadenopathy, often of the jaw (Fig. 20.16), which is initially very responsive to chemotherapy although long-term cure is uncommon. Sporadic Burkitt lymphoma may occur anywhere in the world and EBV infection is seen in 20% of cases. There is an increased incidence in HIV infection. The histological picture is distinctive with a very high proliferative index of over 95% (Fig. 20.17). The prognosis is excellent using chemotherapy regimens which include high-doses of methotrexate, cytosine arabinoside and cyclophosphamide, e.g. CODOX-M/IVAC (which also includes doxorubicin, ifosfamide and etoposide). Intrathecal chemotherapy is also given.




Figure 20.17 Burkitt lymphoma: histological section of lymph node showing sheets of lymphoblasts and 'starry sky' tingible body macrophages.

Primary central nervous system lymphoma

These are rare tumours, more common in older patients and those with HIV infection. Patients are treated with high-dose methotrexate and cytosine arabinoside. Whole brain radiotherapy is also considered. Long-term cognitive dysfunction is an issue.

Lymphoblastic lymphomas

Lymphoblastic lymphomas (B or T cell) occur mainly in children and young adults and these conditions merge clinically and morphologically with acute lymphoblastic leukaemia (ALL). The cells like those in ALL are terminal transferase positive (see Chapter 17), whereas this test is negative in all the other B- and T-cell lymphomas. They are treated using similar protocols.

T-cell lymphomas

Peripheral T-cell lymphomas that present with lymphadenopathy rather than extranodal disease are a heterogeneous group of rare tumours and are usually of CD4⁺ phenotype. Several variants of T-cell lymphomas are recognized.

Peripheral T-cell non-Hodgkin lymphoma, unspecified

These derive from T cells at various stages of differentiation. They are treated with combined chemotherapy (e.g. CHOP). The prognosis is poor. Autografting for patients with chemosensitive disease is often performed.

Angioimmunoblastic lymphadenopathy

Angioimmunoblastic lymphadenopathy usually occurs in elderly patients with lymphadenopathy, hepatosplenomegaly, skin rashes and a polyclonal increase in serum IgG. Patients are treated with chemotherapy or histone deacetylase inhibitors.

Mycosis fungoides

Mycosis fungoides is a chronic cutaneous T-cell lymphoma that presents with severe pruritus and psoriasis-like lesions (Fig. 20.18). Ultimately, deeper organs are affected, particularly lymph nodes, spleen, liver and bone marrow. Treatment is with phototherapy or chemotherapy.

Sézary syndrome

In Sézary syndrome there is dermatitis, erythroderma, lymphadenopathy and circulating T-lymphoma cells. The cells are usually CD4⁺ and have a folded or cerebriform nuclear chromatin. Initial treatment is by local irradiation, topical chemotherapy or photochemotherapy with psoralen and ultraviolet light (PUVA). Chemotherapy (e.g. CHOP) may be needed but is rarely effective for long.

Adult T-cell leukaemia/lymphoma

This is associated with human T-cell leukaemia/lymphoma virus type 1 (HTLV-1) infection (see Chapter 18).

Enteropathy-associated T-cell lymphomas

Enteropathy-associated T-cell lymphomas are associated with coeliac disease and have a very poor response to treatment. Trials using high-dose methotrexate and autografts are in progress.

Anaplastic large cell lymphoma

Anaplastic large cell lymphoma is particularly common in children and is usually of T-cell phenotype. The disease is CD30⁺ and usually associated with the t(2;5) (p23;q35) translocation. This leads to overexpression of anaplastic lymphoma kinase (ALK). It has an aggressive course, characterized by systemic symptoms and extranodal involvement. ALK-negative cases occur and have a worse prognosis. Crizotinib is a specific inhibitor of ALK activity and is a valuable treatment.



Figure 20.18 Mycosis fungoides.


Histiocytic and dendritic cell neoplasms

These are rare tumours including dendritic and macrophage-derived sarcomas which may be localized or disseminated. They usually present as tumours at extranodal sites, especially the intestinal tract, skin and soft tissues. Systemic symptoms are present. The outlook is poor except for those with small localized tumours who may do well. Langerhans' cell histiocytosis, a clonal disease of histiocytes, is discussed on p. 98.

- Non-Hodgkin lymphomas are a large group of clonal lymphoid tumours. Approximately 85% are of B-cell origin and 15% derive from T or NK cells.
- Their clinical presentation and natural history are more variable than Hodgkin lymphoma and can vary from very indolent disease to rapidly progressive subtypes that need urgent treatment.
- The NHL are divided into low-grade and high-grade disease. Low-grade disorders are typically slowly progressive, respond well to chemotherapy but are difficult to cure, whereas high-grade lymphomas are aggressive and need urgent treatment but are more often curable.
- Investigation is with lymph node biopsy, blood tests and imaging usually by PET/CT. Immunohistochemistry of the lymph node is essential and cytogenetic or gene mutation analysis is helpful in many cases.
- Clinical staging is performed as for Hodgkin lymphoma.
- Some of the more common subtypes include:
 - Small lymphocytic lymphoma* is the lymphoma equivalent of chronic lymphocytic leukaemia.
 - Lymphoplasmacytic lymphoma* usually produces an IgM paraprotein, when it is also known as Waldenström's macroglobulinaemia, and often leads to anaemia and hyperviscosity.
 - Marginal zone lymphomas* arise from marginal zone B cells of lymphoid follicles and can occur as mucosa associated (MALT) lymphoma, most frequent in the stomach.
- Follicular lymphoma* represents 25% of all NHL and is associated with the t(14;18) translocation. Treatment usually achieves disease remission but the only curative option is allogeneic stem cell transplantation.
- Mantle cell lymphoma* is associated with increased expression of the cyclin D1 gene and has clinical features of an 'intermediate grade' lymphoma.
- Diffuse large B-cell lymphoma* is a common subtype and is an aggressive disease which needs urgent treatment. There is a wide variety of sub-types. Over 50% of cases are cured.
- Burkitt lymphoma* is one of the most highly proliferative subtypes of tumour. Endemic cases in Africa are associated with EBV infection. Treatment is with aggressive chemotherapy regimens.
- T-cell lymphomas are less common and include *mycosis fungoides*, *peripheral T-cell lymphomas* and *anaplastic large cell lymphoma*.
- Treatments for NHL are based on a variety of chemotherapy regimens. Anti-CD20 antibodies are used in most cases of B-cell lymphomas and have markedly improved the prognosis.



Now visit www.wileyessential.com/haematology to test yourself on this chapter.


A vertical strip on the left side of the page showing a microscopic view of blood cells. It features several red blood cells (erythrocytes) in shades of red and pink, and a few white blood cells (leukocytes) with distinct nuclei. The background is a deep blue.

CHAPTER 21

Multiple myeloma and related disorders

Key topics

■ Paraproteinaemia	229
■ Multiple myeloma	229
■ Other plasma cell tumours	237
■ Monoclonal gammopathy of undetermined significance	237
■ Amyloidosis	238
■ Hyperviscosity syndrome	240



Paraproteinaemia

This is the presence of a monoclonal immunoglobulin band in the serum (Fig. 21.1). Normally, serum immunoglobulins are polyclonal and represent the combined output from millions of different plasma cells. A monoclonal band (M-protein), or **paraprotein**, reflects the synthesis of immunoglobulin from a single clone of plasma cells. This may occur as a primary neoplastic disease or secondary to an underlying benign or neoplastic disease affecting the immune system (Table 21.1).

Multiple myeloma

Multiple myeloma is a neoplastic disease characterized by **plasma cell accumulation in the bone marrow (Fig. 21.2), the presence of monoclonal protein in the serum and/or urine and, in symptomatic patients, related tissue damage** (Table 21.3, Fig. 21.3). Other plasma cell neoplasms are listed in Table 21.2.

Ninety-eight per cent of cases of myeloma occur over the age of 40 years with a peak incidence between 65 and 70 years. The disease is twice as common in black individuals compared to those of white or Asian origin.

Pathogenesis

The myeloma cell is a post-germinal centre plasma cell that has undergone immunoglobulin class switching and somatic hypermutation and secretes the paraprotein that is present in serum. The tumour cell typically contains an average of

Table 21.1 Diseases associated with monoclonal immunoglobulins.

Neoplastic
Multiple myeloma
Solitary plasmacytoma
Monoclonal gammopathy of undetermined significance (MGUS)
Waldenström's macroglobulinaemia
Non-Hodgkin lymphoma
Chronic lymphocytic leukaemia
Primary amyloidosis
Heavy-chain disease
Benign
Chronic cold haemagglutinin disease
Transient (e.g. with infections)
HIV infection
Gaucher's disease

35 somatic mutations at the time of diagnosis. Immunoglobulin heavy and light chain genes are clonally rearranged, with translocations involving the heavy chain on chromosome 14q being the most frequent finding. Normal plasma cells are located within the bone marrow and this feature is retained by the tumour cell. Tumour cells accumulate complex genetic changes with aneuploidy present in almost all cases. Dysregulated or increased expression of the cyclin D1, D2 or D3 genes, either directly through translocations or indirectly via other mutations, is an early unifying event. Later events include secondary translocations (e.g. of *MYC*), point mutations (e.g. *RAS*), deletions (e.g. *TP53*) or epigenetic abnormalities.

Analysis of stored serum samples has shown that almost all cases of myeloma develop from a pre-existing monoclonal gammopathy of undetermined significance (MGUS) (Table 21.3, Fig. 21.3). Most of the early genetic changes are present at the MGUS stage but the size of the clone is considerably smaller (Fig. 21.3).

Myeloma cells adhere to bone marrow stromal cells and extracellular matrix through a variety of adhesion molecules. This inhibits their apoptosis and stimulates a cascade of cytokine release. Osteolytic lesions are caused by osteoclast activation resulting from high serum levels of RANKL (receptor activator of nuclear factor- κ B (NF- κ B) ligand), produced by plasma cells and bone marrow stroma, which binds to activating RANK receptors on the osteoclast surface.

Smouldering myeloma

The term **asymptomatic** or **smouldering myeloma** is used for cases with similar laboratory findings but no organ or tissue damage causing clinical features (Table 21.3). There is about 10% chance each year of these cases becoming symptomatic and requiring therapy. The risk is greater if there are more than




Figure 21.1 Serum protein electrophoresis in multiple myeloma showing an abnormal paraprotein in the γ -globulin region with reduced levels of background β - and γ -globulins.




Figure 21.2 (a) The bone marrow in multiple myeloma showing large numbers of plasma cells, with many abnormal forms. (b) Low-power view showing sheets of plasma cells replacing normal haemopoietic tissue. (c) Immunohistochemical staining of the bone marrow in myeloma with antibody to CD138 revealing extensive numbers of plasma cells.




Figure 21.3 The degree of involvement of bone marrow by clonal tumour cells in: (a) MGUS, smouldering and symptomatic myeloma and plasma cell leukaemia associated with a serum IgG or IgA paraprotein; (b) MGUS and Waldenström's macroglobulinaemia (lymphoplasmacytic lymphoma) associated with a serum IgM paraprotein.

Table 21.2 Plasma cell neoplasms (WHO, 2008).

Monoclonal gammopathy of undetermined significance (MGUS)

Plasma cell myeloma

Variants:

Asymptomatic (smouldering) myeloma

Non-secretory myeloma

Plasma cell leukaemia

Plasmacytoma

Solitary plasmacytoma of bone

Extraosseous (extramedullary) plasmacytoma

Immunoglobulin deposition diseases

Primary amyloidosis

Systemic light and heavy chain deposition diseases

Osteosclerotic myeloma (POEMS syndrome)

60% plasma cells in the marrow, circulating plasma cells, a greatly unbalanced free light chain ratio (see below) or certain unfavourable cytogenetic abnormalities. Treatment as for symptomatic myeloma is usually begun if there are spinal lesions detected on MRI or more than 60% plasma cells in the marrow.

Diagnosis

Symptomatic myeloma is diagnosed if there is:

- 1 Monoclonal protein in serum and/or urine (Fig. 21.1);
- 2 Increased clonal plasma cells in the bone marrow (Fig. 21.2); and
- 3 Related organ or tissue impairment.

A useful acronym for tissue damage is CRAB (hypercalcaemia, renal impairment, anaemia, bone disease) (Table 21.3). Amyloid, hyperviscosity, recurrent infections, peripheral neuropathy and deep vein thrombosis are other clinical complications which are less frequently presenting features (Fig. 21.4).

Clinical features

- 1 **Bone pain** (especially backache) resulting from vertebral collapse and pathological fractures (Fig. 21.5a,b).
- 2 Features of **anaemia**, such as lethargy or weakness.
- 3 **Recurrent infections**: related to deficient antibody production, abnormal cell-mediated immunity and neutropenia.
- 4 Features of **renal failure** and/or hypercalcaemia: polydipsia, polyuria, anorexia, vomiting, constipation and mental disturbance.
- 5 Abnormal **bleeding** tendency: myeloma protein may interfere with platelet function and coagulation factors; thrombocytopenia occurs in advanced disease.
- 6 **Amyloidosis** occurs in 5% with features such as macroglossia, carpal tunnel syndrome and diarrhoea.
- 7 In approximately 2% of cases there is a **hyperviscosity syndrome** with purpura, haemorrhages, visual failure, central nervous system (CNS) symptoms, neuropathies and heart failure.

Laboratory findings include the following:

- 1 **Presence of a paraprotein** Serum and urine should be screened by immunoglobulin electrophoresis. The paraprotein is immunoglobulin G (IgG) in 60% of cases, IgA in 20% and light chain only in almost all the rest. Less than

Table 21.3 The clinical and laboratory features of monoclonal gammopathy of uncertain significance (MGUS), smouldering myeloma and symptomatic myeloma.

	MGUS	Smouldering myeloma	Symptomatic myeloma
Marrow plasma cells	<10%	≥ 10%	≥ 10%
Paraprotein	<30 g/L	≥30 g/L	≥30 g/L
Normal immunoglobulins	Normal	Reduced	Reduced
Free light chain ratio	Normal or abnormal	Abnormal	Abnormal
Clinical features	None	None	Hypercalcaemia C
			Renal failure R
			Anaemia A
			Bone lesions B*
Progression to symptomatic myeloma	1%/year	10%/year	
*MRI or PET scan needed to exclude bone lesions, especially of the spine.			
Other clinical features include amyloid, hyperviscosity, recurrent infections, peripheral neuropathy and deep vein thrombosis.			




Figure 21.4 The pathogenesis of the clinical features of myeloma.

1% have IgD or IgE paraprotein and a similar number are non-secretory.

- 2 **Elevated serum immunoglobulin-free light chains** Immunoglobulin-free light chains (FLC) are κ or λ light chain proteins, synthesized by plasma cells, that have not been paired with heavy chain (Fig. 21.6). They are normally made in small quantities and filtered from the serum into the kidney but can be measured in serum. Free light chains are produced by almost all malignant plasma cells and so the *serum free light chain* assay is useful in diagnosis and monitoring of myeloma and other forms of malignant paraproteinaemia. Typically in myeloma there is an increase in either the κ or λ serum free light chain value. The normal κ : λ serum free light chain ratio of 0.6 (range 0.26–1.65) is also skewed with an excess of either κ or λ chains. Light chain assays have largely replaced the need for analysis of urine paraproteinaemia.
- 3 Normal serum immunoglobulin levels (IgG, IgA and IgM) are reduced, a feature known as **immunoparesis**. The urine contains free light chains, **Bence-Jones protein**, in two-thirds of cases. Rare cases of myeloma are non-secretory and therefore not associated with a paraprotein or Bence-Jones proteinuria although some will still show a disturbed free light chain ratio in the serum.
- 4 There is usually a normochromic normocytic or macrocytic anaemia. Rouleaux formation is marked in most cases (Fig. 21.7). Neutropenia and thrombocytopenia occur in advanced disease. Abnormal plasma cells appear in the blood film in 15% of patients and can be detected by sensitive flow cytometry in over 50%.
- 5 High erythrocyte sedimentation rate (ESR).
- 6 Increased plasma cells in the bone marrow (usually more than 20%) often with abnormal forms (Fig. 21.2). The

characteristic **immunophenotype** of malignant plasma cells is $CD38^{\text{high}}$, $CD138^{\text{high}}$ and $CD45^{\text{low}}$. Anti-CD138 is used to measure the number of plasma cells in the marrow biopsy (Fig. 21.2).

- 7 Radiological investigation of the skeleton reveals bone lesions such as osteolytic areas without evidence of surrounding osteoblastic reaction or sclerosis in 60% of patients (Fig. 21.8) or generalized osteoporosis in 20% (Fig. 21.5). In addition, pathological fractures or vertebral collapse (Fig. 21.5b) are common. Skeletal survey X-rays do not detect small lesions. **MRI of the spine** is needed in cases considered as smouldering myeloma in order to detect potential early bone lesions. PET scan is also a sensitive imaging technique to detect bone damage (Fig. 21.5).
- 8 Serum calcium elevation occurs in 45% of patients. Typically, the serum alkaline phosphatase is normal, except following pathological fractures.
- 9 The serum creatinine is raised in 20% of cases. Proteinaceous deposits from light chain proteinuria, hypercalcaemia, uric acid, amyloid and pyelonephritis may all contribute to renal failure (Fig. 21.9).
- 10 A low serum albumin occurs with advanced disease.
- 11 Serum β_2 -microglobulin is often raised and is a useful indicator of prognosis.
- 12 The molecular and cytogenetic changes have been described earlier. Cytogenetic analysis shows that aneuploidy (more or less than 46 chromosomes) is almost universal. There is also a high incidence of translocations involving the immunoglobulin heavy-chain gene (*IGH*) on chromosome 14. The D-cyclin genes, D_1 , D_2 or D_3 , are often involved in the translocations. Later stage cases have increased complex cytogenetic findings as well as more point mutations relevant to progression.




Figure 21.5 (a) Multiple myeloma: X-ray of lumbar spine showing severe demineralization with partial collapse of L_3 . (b) Magnetic resonance imaging (MRI) of spine: T_2 -weighted study. There is infiltration and destruction of L_3 and L_5 with bulging of the posterior part of the body of L_3 into the spinal canal compressing the corda equina (arrowed). Radiotherapy has caused a marrow signal change in vertebrae C_2 – D_4 because of replacement of normal red marrow by fat (bright white signal). Source: Courtesy of Dr A. Platts. (c) (i) Computed tomography scan of base of skull showing paravertebral soft tissue extramedullary plasmacytoma (white arrow). (ii) Positron Emission Tomography (PET) scan of the same patient showing extensive medullary and extramedullary involvement. Source: Sher T. *et al.* (2010) *Br J Haematol* 150(4): 418–27. Reproduced with permission of John Wiley & Sons.

Figure 21.6 The value of serum immunoglobulin-free light chain (FLC) measurement in multiple myeloma. **(a)** Serum free light chains are immunoglobulin light chains that are synthesized by plasma cells but not paired with heavy chains before they are released into the blood. Low levels are found in normal individuals and these are increased in patients with myeloma. **(b)** Profile of serum free light chains in healthy controls, patients with renal impairment and those with κ or λ light chain multiple myeloma (LCMM). As light chains are normally filtered by the kidney, their levels rise in patients with renal impairment although the κ : λ ratio remains normal. Source: Adapted from Hutchison C. (2008) *BMC Nephrology* 9, 11–19 and Bradwell A.R. (2003) *Lancet* 361, 489–91.





Figure 21.7 The peripheral blood film in multiple myeloma showing rouleaux formations.




Figure 21.8 Skull X-ray in multiple myeloma showing many 'punched-out' lesions.




Figure 21.9 The kidney in multiple myeloma. **(a)** Myeloma kidney: the renal tubules are distended with hyaline protein (precipitated light chains or Bence-Jones protein). Giant cells are prominent in the surrounding cellular reaction. **(b)** Amyloid deposition: both glomeruli and several of the small blood vessels contain an amorphous pink-staining deposit characteristic of amyloid (Congo red stain). **(c)** Nephrocalcinosis: calcium deposition (dark 'fractured' material) in the renal parenchyma. **(d)** Pyelonephritis: destruction of renal parenchyma and infiltration by acute inflammatory cells.

Treatment

This may be divided into specific and supportive (Fig. 21.10).

Specific

The life expectancy of patients with myeloma has improved markedly in recent years with the introduction of new drugs such as proteasome inhibitors and immunomodulatory agents. The major initial treatment decision is between the use of **intensive combination chemotherapy** (mostly for patients aged less than 70 years) or **non-intensive therapy** for older patients.

Intensive therapy

Intensive therapy involves four to six courses of chemotherapy to reduce the tumour burden, usually followed by stem cell collection and autologous SCT after high-dose chemotherapy. The initial chemotherapy is given as repeated intravenous or oral chemotherapy cycles combining two or three drugs: bortezomib, dexamethasone, thalidomide, lenalidomide or cyclophosphamide.

Some of the typical combination therapies are:

- VTD – bortezomib (velcade), thalidomide and dexamethasone
- VCD – velcade, cyclophosphamide and dexamethasone
- VRD – velcade, lenalidomide (revlimid) and dexamethasone
- CDT – cyclophosphamide, dexamethasone and thalidomide
- RD – lenalidomide (revlimid) and dexamethasone

Trials in initial therapy of new drugs, pomalidomide and carfilzomib, already found to be effective in relapsed patients, are in progress.

After several courses of treatment, when the number of tumour cells has been reduced, the patient usually undergoes an autologous stem cell transplant. Peripheral blood stem cells are usually collected after mobilization using a combination of chemotherapy and granulocyte colony-stimulating factor (G-CSF). High-dose melphalan is the typical conditioning regimen that is given. Two consecutive autologous transplants are used in some centres for selected patients. Trials of post-transplant consolidation with two or three courses of chemotherapy are in progress. Maintenance treatment after autologous SCT is also in clinical study, using drugs such as thalidomide, lenalidomide or bortezomib, examining whether there is increased overall survival and for side-effects. About 10% of patients who have undergone intensive therapy and autologous SCT appear to be cured and the median life expectancy has increased from 2–3 to 8 years or more. As drug treatments improve, it may become preferable to omit SCT from future protocols or perform autologous SCT only after first relapse.

Although *allogeneic transplantation* may cure the disease it carries a high procedure-related mortality. Moreover patients frequently relapse after the procedure. It is only used in highly selected patients.

Non-intensive therapy

Typical options are courses of the oral alkylating agent melphalan in combination with prednisolone, and thalidomide (MPT) or bortezomib (VMP). Alternatively agents similar to those used in the intensive regimes, such as VCD or RD (lenalidomide and dexamethasone) may be used. These protocols are normally given monthly, depending on response, for up to 12–18 months. Typically, paraprotein levels fall after treatment, bone lesions show improvement and blood counts may improve. When treatment is stopped the patient is seen at regular intervals in the outpatient clinic.

Relapsed patients

After a variable period of time the disease is likely to progress with rising paraprotein levels and return of symptoms.




Figure 21.10 Algorithm of potential approaches to the management of multiple myeloma.

Further chemotherapy may be given with the above drugs, although the disease becomes increasingly difficult to control. Local radiotherapy to painful bone lesions can be given where needed. Bendamustine appears to be effective in some relapsed cases and new drugs and monoclonal antibodies are in trial.

Notes on specific drugs used in myeloma

Thalidomide was the first immune-modulator drug to be used in myeloma. It has a number of side-effects such as sedation,

constipation, neuropathy and venous thrombosis (Table 21.3). The addition of dexamethasone increases the response rate but venous thrombosis becomes a major concern and prophylactic anticoagulation with low molecular weight heparin, warfarin or aspirin is needed.

Lenalidomide is an analogue of thalidomide and is highly active in the management of myeloma. It is used widely for first-line therapy and for relapsed disease. It is associated with myelosuppression and increased risk of thrombosis, but causes less neuropathy than thalidomide (Table 21.3).

Pomalidomide is the most recent addition to this class of drugs and shows a high level of activity against relapsed disease. Like lenalidomide it causes less neuropathy than thalidomide.

Bortezomib inhibits the cellular proteasome and NF- κ B activation, and is very valuable in the treatment of myeloma. It is highly active in first-line treatment and relapsed disease. Its main side-effect is a neuropathy. **Carfilzomib** is a newly introduced proteasome inhibitor which is less likely than bortezomib to cause a neuropathy.

Radiotherapy is useful in treating the symptomatic bone disease in myeloma. It is used for areas of bone pain or spinal cord compression.

Supportive care

Renal impairment It is advisable for patients to drink at least 3 L of fluid each day throughout the course of their disease in order to limit the accumulation of paraprotein within the kidney. Some patients present with renal failure and treatment should include rehydration and treatment of contributing factors, such as hypercalcaemia or hyperuricaemia. Dialysis is generally well tolerated if required.

Bone disease and hypercalcaemia Bisphosphonates, such as pamidronate or zoledronic acid, are effective in reducing the progression of bone disease and may also improve overall survival. Acute hypercalcaemia is treated with rehydration using isotonic saline, a diuretic and corticosteroids followed by a bisphosphonate.

Compression paraplegia Surgical decompression laminectomy or irradiation are treatments of choice. Chemotherapy or corticosteroid treatment is usually also given.

Anaemia Transfusion or erythropoietin may be used.

Bleeding Bleeding caused by paraprotein interference with coagulation and hyperviscosity syndrome may be treated by repeated plasmapheresis.

Infections Rapid treatment of any infection is essential. Prophylactic infusions of immunoglobulin concentrates together with oral broad-spectrum antibiotics and antifungal agents may be needed for recurrent infections.

Prognosis

The outlook for patients with myeloma is improving markedly. The overall median survival is now 7–10 years and in younger (less than 50 years) patients it can be over 10 years.

Other plasma cell tumours

Solitary plasmacytoma

These are isolated plasma cell tumours, usually involving bones or soft tissue such as the mucosa of the upper respiratory and gastrointestinal tracts or the skin. The associated

paraprotein disappears following radiotherapy to the primary lesion.

Plasma cell leukaemia

This rare disease is characterized by a high number of circulating malignant plasma cells (Fig. 21.3). The clinical features tend to be a combination of those found in acute leukaemia (pancytopenia and organomegaly) with features of myeloma (hypercalcaemia, renal involvement and bone disease). Treatment is with supportive care and systemic chemotherapy but prognosis is poor.

Osteosclerotic myeloma (POEMS syndrome)

This is a rare condition in which polyneuropathy is associated with a monoclonal plasma cell disorder and osteosclerotic bone lesions. Elevated levels of vascular endothelial growth factor (VEGF) are present and there are often clinical features such as splenomegaly, hepatomegaly or lymphadenopathy, extravascular fluid overload, endocrine abnormalities, skin changes or Castlemans disease.

Monoclonal gammopathy of undetermined significance

Transient or persistent paraproteins can occur in many other conditions as well as in multiple myeloma (Table 21.1). A serum paraprotein may be sometimes be detected without any evidence of myeloma or other underlying disease and is termed **monoclonal gammopathy of undetermined significance (MGUS)**. It is increasingly common with age, being present in 3–4% of persons older than 50 years. There is an increase in incidence of venous and arterial thrombosis, infections, osteoporosis and bone fractures compared to controls. The proportion of plasma cells in the marrow is normal (less than 4%) or slightly raised (less than 10%) (Table 21.3, Fig. 21.3). The concentration of monoclonal immunoglobulin in serum is less than 30 g/L and other serum immunoglobulins are not depressed. The κ or λ light chain is increased in serum in one-third of patients; the greater the imbalance, the more the risk of transformation. No treatment is needed but patients with IgG or IgA MGUS develop overt myeloma at a rate of 1% each year and so are usually followed up regularly in the outpatient clinic. The survival of patients with MGUS is reduced compared with control populations and this effect increases with duration of follow-up and age (Fig. 21.11).

Amyloidosis

The amyloidoses are a heterogeneous group of disorders characterized by the extracellular deposition of protein in an abnormal fibrillar form (Table 21.4). Amyloidosis may be hereditary






Figure 21.11 Survival of patients with monoclonal gammopathy of undetermined significance (MGUS) ($n = 1384$) is reduced compared to the control population; median 8.1 vs. 11.8 years, respectively ($p < 0.001$). Source: Adapted from Kyle R.A. *et al.* (2009) *Haematologica* 94: 1714.



(a)



(b)

Figure 21.12 Amyloidosis. (a) Congo red staining and (b) blue–green birefringence under polarized light.

or acquired and deposits may be focal, localized or systemic in distribution. The amyloid is made from different amyloid fibril precursor proteins in each type of disease. Except for intracerebral amyloid plaques, all amyloid deposits contain a non-fibrillary glycoprotein amyloid P which is derived from a normal serum precursor structurally related to C-reactive protein (CRP). The classic diagnostic histological test is red–green birefringence after staining with Congo red and viewing under polarized light (Fig. 21.12).

Systemic AL amyloidosis

Systemic amyloid light chain (AL) amyloid disease is caused by deposition of monoclonal light chains produced from a clonal plasma cell proliferation. The level of paraprotein may be very low and is not always detectable in serum or urine but the serum free light chain ratio is usually abnormal (Fig. 21.6). It is responsible for 1 in 1500 deaths in the UK. The clinical features are caused by involvement of the heart in more than 50% of patients usually with a restrictive cardiomyopathy, kidneys in 30% (Figure 21.9), tongue (Fig. 21.13), gastrointestinal tract, peripheral nerves and autonomic nervous system (Fig. 21.9). The patient may present with non-specific symptoms such as fatigue, anorexia, weight loss, or with heart failure, renal failure including the nephrotic syndrome, macroglossia, peripheral neuropathy or carpal tunnel syndrome. A serum amyloid P (SAP) scan (Fig. 21.14) is used to determine the extent and severity of disease. Treatment is with chemotherapy similar to that used in myeloma, possibly with autologous SCT, which may improve prognosis. Patients are, however, more at risk of serious side-effects than those with myeloma.




Figure 21.13 Multiple myeloma: the tongue and lips are enlarged because of nodular and waxy deposits of amyloid.

Table 21.4 Classification of amyloidosis: types, structure and organ involvement.

Type	Chemical nature	Organs involved
Systemic AL amyloidosis Associated with myeloma, Waldenström's macroglobulinaemia or MGUS May also occur on its own as primary amyloidosis (associated with an occult plasma cell proliferation) May also occur in localized form with local 'immunocyte' proliferation	Immunoglobulin light chains (AL)	Tongue Skin Heart Nerves Connective tissue Kidneys Liver Spleen
Reactive systemic AA amyloidosis Rheumatoid arthritis, tuberculosis, bronchiectasis, chronic osteomyelitis, inflammatory bowel disease, Hodgkin lymphoma, carcinomas, familial Mediterranean fever	Protein A (AA)	Liver Spleen Kidneys Bone marrow
Familial amyloidosis	e.g. Transthyretin abnormalities	Nerves Heart Eyes
Localized amyloidosis Central nervous system Endocrine Senile	β -Amyloid protein Peptic hormones Various	Alzheimer's disease Endocrine tumours Heart, brain, joints, prostate, etc.

AA, AL, these are defined by their chemical nature as in the table; MGUS, monoclonal gammopathy of undetermined significance.

Figure 21.14 Serial anterior whole body ^{125}I -labelled serum amyloid P component (SAP) scans of a 52-year-old woman who presented with renal failure resulting from systemic AL amyloidosis. **(a)** The initial scan demonstrates a large amyloid load with hepatic, splenic, renal and bone marrow deposits. The underlying plasma cell dyscrasia responded to high-dose melphalan followed by autologous stem cell rescue. **(b)** Follow-up SAP scintigraphy 3 years after chemotherapy showed greatly reduced uptake of tracer indicating substantial regression of her amyloid deposits. Source: Courtesy of Professor P.N. Hawkins, National Amyloidosis Centre, Royal Free Hospital, London.






Figure 21.15 Hyperviscosity syndrome in Waldenström's macroglobulinaemia. **(a)** The retina before plasmapheresis shows distension of retinal vessels, particularly the veins which show bulging and constriction (the 'linked sausage' effect) and areas of haemorrhage; **(b)** following plasmapheresis the vessels have returned to normal and the areas of haemorrhage have cleared.

Hyperviscosity syndrome

The most common cause is polycythaemia (see p. 169). Hyperviscosity may also occur in patients with myeloma or Waldenström's macroglobulinaemia or in patients with chronic myeloid or acute leukaemias associated with very high white cell counts. The clinical features of the hyperviscosity syndrome include visual disturbances, lethargy, confusion, muscle weakness, nervous system symptoms and signs, and congestive heart failure. The retina may show a variety of changes: engorged veins, haemorrhages, exudates and a

blurred disc (Fig. 21.15). Emergency treatment varies with the cause:

- 1 correct dehydration in all patients;
- 2 venesection or isovolaemic exchange with a plasma substitute for red cells in a polycythaemic patient;
- 3 plasmapheresis in myeloma, Waldenström's disease or hyperfibrinogenaemia;
- 4 and leucopheresis or chemotherapy in leukaemias associated with high white cell counts.

The long-term treatment depends on control of the primary disease with specific therapy.

SUMMARY

- The term *paraproteinaemia* refers to the presence of a monoclonal immunoglobulin band in serum and reflects the synthesis of immunoglobulin from a single clone of plasma cells.
- Multiple myeloma is a tumour of plasma cells that accumulate in the bone marrow, release a paraprotein and cause tissue damage. The disease has a peak incidence in the seventh decade.
- Almost all cases of myeloma develop from a pre-existing *monoclonal gammopathy of undetermined significance* (MGUS) in which there is low level paraprotein and no evidence of tissue damage. Approximately 1% of cases progress to myeloma each year.
- A useful reminder for the spectrum of tissue damage in symptomatic myeloma is **CRAB** – hypercalcaemia, renal impairment, anaemia, bone disease.
- Smouldering myeloma describes patients with bone marrow and protein findings of myeloma but without CRAB or other clinical symptoms or findings.
- In patients younger than 70 years, symptomatic myeloma is usually treated by intensive chemotherapy with a three-drug combination followed by an autologous stem cell transplant, using stem cells harvested from the patient.
- In older patients chemotherapy alone is used. Immunomodulatory drugs (thalidomide, lenalidomide, pomalidomide) and proteasome inhibitory drugs

(bortezomib, carfilzomib), usually used in combination with dexamethasone, are improving the outlook for patients and median survival is now 7–10 years.


- A *plasmacytoma* is a localized mass of malignant plasma cells and is usually treated with radiotherapy. Many cases progress to myeloma.
- *Amyloidoses* are caused by extracellular deposition of protein in an abnormal fibrillar form. Systemic AL amyloid disease is caused by monoclonal light chains

produced from a clonal plasma cell proliferation and may cause heart failure, macroglossia, peripheral neuropathy or renal failure.

- *Hyperviscosity syndrome* may occur in paraproteinaemia or in patients with very high red or white cell counts. Clinical features include visual disturbances, confusion and heart failure. Venesection, plasma exchange or chemotherapy may be required.



Now visit www.wileyessential.com/haematology to test yourself on this chapter.


A vertical strip on the left side of the page showing a microscopic view of various blood cells, including red blood cells and white blood cells, in shades of red, pink, and yellow against a dark blue background.

CHAPTER 22

Aplastic anaemia and bone marrow failure

Key Topics

■ Pancytopenia	243
■ Aplastic anaemia	243
■ Paroxysmal nocturnal haemoglobinuria	247
■ Red cell aplasia	247
■ Schwachman–Diamond syndrome	249
■ Congenital dyserythropoietic anaemia	249
■ Osteopetrosis	249



Pancytopenia

Pancytopenia is a reduction in the blood count of all the major cell lines – red cells, white cells and platelets. It has several causes (Table 22.1), which can be broadly divided into decreased bone marrow production or increased peripheral destruction.

Aplastic anaemia

Aplastic (hypoplastic) anaemia is defined as pancytopenia resulting from hypoplasia of the bone marrow (Fig 22.1). It is classified into primary (congenital or acquired) or secondary types (Table 22.2).


Table 22.1 Causes of pancytopenia.

Decreased bone marrow function
Aplasia (reduction of haemopoietic stem cells)
Acute leukaemia, myelodysplasia, myeloma
Infiltration with lymphoma, solid tumours, tuberculosis
Megaloblastic anaemia
Paroxysmal nocturnal haemoglobinuria
Myelofibrosis
Haemophagocytic syndrome
Increased peripheral destruction
Splenomegaly


Table 22.2 Causes of aplastic anaemia.

Primary	Secondary
Congenital (Fanconi and non-Fanconi types)	Ionizing radiation: accidental exposure (radiotherapy, radioactive isotopes,)
Idiopathic acquired	Chemicals: benzene, organophosphates and other organic solvents, DDT and other pesticides, recreational drugs (ecstasy) Drugs: Those that regularly cause marrow depression (e.g. busulfan, melphalan, cyclophosphamide, anthracyclines, nitrosoureas) Those that occasionally or rarely cause marrow depression (e.g. chloramphenicol, sulphonamides, gold, anti-inflammatory, antithyroid, psychotropic, anticonvulsant/ antidepressant drugs) Viruses: viral hepatitis (non-A, non-B, non-C, non-G in most cases), EBV Autoimmune diseases: systemic lupus erythematosus Transfusion associated GVHD (see p. 342) Thymoma (this is more usually associated with red cell aplasia)

EBV, Epstein–Barr virus.



(a)



(b)

Figure 22.1 Aplastic anaemia: low power views of bone marrow show severe reduction of haemopoietic cells with an increase in fat spaces. (a) Aspirated fragment. (b) Trephine biopsy.

Pathogenesis

The underlying defect in all cases appears to be a substantial reduction in the number of haemopoietic pluripotential stem cells, and a fault in the remaining stem cells or an immune reaction against them, which makes them unable to divide and differentiate sufficiently to populate the bone marrow.

Congenital: Fanconi anaemia (FA)

The Fanconi type has an autosomal recessive pattern of inheritance and is often associated with **growth retardation and congenital defects of the skeleton** (e.g. microcephaly, absent radii or thumbs), **of the renal tract** (e.g. pelvic or horseshoe kidney) (Fig. 22.2) **or skin** (areas of hyper- and hypopigmentation); sometimes there is learning disability. The syndrome is genetically heterogeneous with 16 different genes involved: *FANCA–Q*. *FANCD1* is identical to *BRCA2*, the breast cancer susceptibility gene. The proteins coded for by these genes cooperate in a common cellular pathway which results in ubiquitination of *FANCD2*, which protects cells against genetic damage. Cells from FA patients show an abnormally high frequency of spontaneous chromosomal breakage and the diagnostic test is elevated breakage after incubation of peripheral blood lymphocytes with the DNA cross-linking agent diepoxybutane (DEB test).

Dyskeratosis congenita (DC) is a rare sex-linked disorder with nail and skin atrophy and a high risk of pulmonary fibrosis, cirrhosis, osteoporosis and cancer. It is associated with mutations in the *DKC1* (dyskerin) or *TERC* (telomerase reverse transcriptase RNA template) involved in the maintenance of telomere length.

The usual age of presentation of FA is 3–14 years. Approximately 10% of patients develop acute myeloid leukaemia. Treatment is usually with androgens and/or SCT. The blood count usually improves with androgens but side-effects, especially in children, are distressing (virilization and liver abnormalities); remission rarely lasts more than 2 years. SCT may cure the patient. Because of the sensitivity of the patient's cells to DNA damage, conditioning regimes are mild and irradiation avoided.

Other inherited bone marrow failure syndromes include Diamond–Blackfan anaemia (DBA), (p. 247), Schwachman–Diamond syndrome (SDS), (p. 249), severe congenital neutropenia (p. 95), amegakaryocytic thrombocytopenia (p. 281) and thrombocytopenia with absent radii (p. 281). In DC, DBA and SDS there are genetic defects in ribosomal biosynthesis and function (Fig. 22.3).

Idiopathic acquired aplastic anaemia

This is the most common type of aplastic anaemia, accounting for at least two-thirds of acquired case. In most cases haemopoietic tissue is the target of an autoimmune process dominated by oligoclonal expression of cytotoxic CD8⁺ T cells. Clonal haemopoiesis with somatic mutations of genes such as *PIGA*, *ASXL1* and *DNMT3A*, presumably arising by selection in a failed marrow, occur in about 50% of cases. The disease must be distinguished from a late onset of a congenital form of aplastic anaemia and from hypoplastic myelodysplasia. Mutations of the telomere repair complex and short telomeres may be present, apparently as acquired abnormalities. The favourable responses to antilymphocyte globulin (ALG) and ciclosporin support the concept of an autoimmune disorder.




Figure 22.2 (a) X-rays showing absent thumbs in a patient with Fanconi anaemia (FA). (b) Intravenous pyelogram in a patient with FA showing a normal right kidney but a left kidney abnormally placed in the pelvis.




Figure 22.3 Ribosomal RNA processing and the sites of disruption in bone marrow failure syndromes. The different RNA species have roles in cell stress response, proliferation and apoptosis. Source: Courtesy of Professor I. Dokal. DBA, Diamond-Blackfan anaemia; SDS, Schwachman-Diamond syndrome.

Secondary causes

Aplastic anaemia may be caused by direct damage to the haemopoietic marrow by radiation or cytotoxic drugs. The antimetabolite drugs (e.g. methotrexate) and mitotic inhibitors (e.g. daunorubicin) cause only temporary aplasia but the alkylating agents, particularly busulfan, may cause chronic aplasia closely resembling the chronic idiopathic disease. Some individuals develop aplastic anaemia as a rare idiosyncratic side-effect of drugs such as chloramphenicol or gold (Table 22.2). They may also develop the disease during or within a few months of viral hepatitis (most frequently negative for all known hepatitis viruses). Because the incidence of marrow toxicity is particularly high for chloramphenicol, this drug should be reserved for treatment of infections that are life-threatening and for which it is the optimum antibiotic (e.g. typhoid). Chemicals such as benzene may be implicated and, rarely, aplastic anaemia may be the presenting feature of acute lymphoblastic or myeloid leukaemia, especially in childhood. Myelodysplasia (see Chapter 16) may also present with a hypoplastic marrow.

Clinical features

The onset is at any age with peak incidences around 10–25 and over 60 years. It is more frequent in Asia, e.g. China, than Europe. It can be insidious or acute with symptoms and signs resulting from anaemia, neutropenia or thrombocytopenia. Bruising, bleeding gums, epistaxes and menorrhagia are the most frequent haemorrhagic manifestations and the usual presenting features (Fig 22.4), often with symptoms of anaemia. Retinal haemorrhage may impair vision. Infections, particularly of the mouth and throat, are common and generalized infections are frequently life threatening. The lymph nodes, liver and spleen are not enlarged. A careful history and



Figure 22.4 Aplastic anaemia: spontaneous mucosal haemorrhages in a 10-year-old boy with severe Fanconi anaemia. Platelet count $<5 \times 10^9/L$. Source: Hoffbrand A.V., Pettit J.E. & Vyas P. (2010) *Color Atlas of Clinical Hematology*, 4th edn. Reproduced with permission of John Wiley & Sons.

examination, e.g. for bone deformities, are needed at all ages to exclude inherited forms.

Laboratory findings

In aplastic anaemia, there must be at least two of the following:

- 1 Anaemia (haemoglobin <100 g/L). This is normochromic, normocytic or macrocytic (mean cell volume (MCV) often 95–110 fL). The reticulocyte count is usually extremely low in relation to the degree of anaemia.
- 2 Neutrophil count $<1.5 \times 10^9$ /L.
- 3 Platelet count $<50 \times 10^9$ /L.
- 4 Severe cases show neutrophils $<0.5 \times 10^9$ /L and platelets $<20 \times 10^9$ /L, reticulocytes $<20 \times 10^9$ /L and marrow cellularity $<25\%$. Very severe cases show neutrophils $<0.2 \times 10^9$ /L.
- 5 There are no abnormal cells in the peripheral blood.
- 6 The bone marrow shows hypoplasia, with loss of haemopoietic tissue and replacement by fat which comprises over 75% of the marrow. Trephine biopsy may show patchy cellular areas in a hypocellular background. The main cells present are lymphocytes and plasma cells; megakaryocytes in particular are severely reduced or absent.
- 7 Cytogenetic and, more recently, molecular analysis is performed to exclude inherited forms and myelodysplasia.

Diagnosis

The disease must be distinguished from other causes of pancytopenia (Table 22.1). The assessment of disease severity is also important in treatment decisions and prognosis.

Paroxysmal nocturnal haemoglobinuria (PNH) must be excluded by flow cytometry testing of red cells for CD55 and CD59. In older patients, hypoplastic myelodysplasia may show similar appearances. Abnormalities of the blood cells and clonal cytogenetic or molecular changes suggest myelodysplasia. Some patients diagnosed as having aplastic anaemia develop PNH, myelodysplasia or acute myeloid leukaemia in subsequent years. This may occur even in patients who have responded well to immunosuppressive therapy. Large granular lymphocytic leukaemia (Chapter 18) may also be associated with pancytopenia and a hypoplastic marrow.

Treatment

This is best carried out in a specialized centre.

General

If a primary cause such as a drug has been identified, this is removed. Initial management consists largely of supportive care with blood transfusions, platelet concentrates, and treatment and prevention of infection. All blood products should be leucodepleted, to reduce the risk of alloimmunization, and irradiated, to prevent grafting of live donor lymphocytes. An antifibrinolytic agent (e.g. tranexamic acid) may be used to reduce haemorrhage in patients with severe prolonged thrombocytopenia. Granulocyte transfusions are rarely used, but may be given to patients with severe bacterial or fungal infections not responding to antibiotics. Oral antibacterial and antifungal drugs may be used to reduce infections.

Specific

This must be tailored to the severity of the illness as well as the age of the patient and availability of stem cell donors. Severe cases have a high mortality in the first 6–12 months unless they respond to specific therapy. Less severe cases may have an acute transient course or a chronic course with ultimate recovery, although the platelet count often remains subnormal for many years. Relapses, sometimes severe and occasionally fatal, may also occur and rarely the disease transforms into myelodysplasia, acute leukaemia or PNH (see below).

The following 'specific' treatments are used with varying success.

- 1 **Antithymocyte globulin** (ATG) This is prepared by immunizing animals (horse or rabbit) with human thymocytes and alone is of benefit in approximately 50–60% of acquired cases. It is usually given with ciclosporin to improve the rate of response. Corticosteroids are given short term to reduce the immediate allergic effects and the incidence and severity of serum sickness (fever, rash and joint pains) which may occur approximately 7 days after administration. The platelet count should be maintained $>10 \times 10^9$ /L and if possible >20 – 30×10^9 /L. If there is no response to ATG after 4 months a second course may be tried, prepared from the same or another species. Overall, up to 80% of patients respond to combined ALG and ciclosporin.
- 2 **Ciclosporin** This is an effective agent which appears particularly valuable in combination with ALG. In older subjects it is sometimes used alone.
- 3 **Alemtuzumab** (anti-CD 52 antibody). This has proved effective in about 50% of patients in small studies and is usually only used after ATG has failed.
- 4 **Eltrombopag** This is a thrombomimetic (see Chapter 25) which stimulates platelet production but may also result in a durable improvement in red cell and neutrophil counts.
- 5 **Androgens** These are beneficial in some patients with FA and acquired aplastic anaemia although an overall improved survival in acquired aplastic anaemia has not been proven. Side-effects are marked, including virilization, salt retention and liver damage with cholestatic jaundice or rarely hepatocellular carcinoma. If there is no response in 4–6 months, androgens should be stopped. If there is a response the drug should be withdrawn gradually.
- 6 **Stem cell transplantation** (SCT) Allogeneic SCT offers the chance of permanent cure in selected patients. Conditioning is with cyclophosphamide without irradiation and ciclosporin is used to reduce the risks of graft failure and graft-versus-host disease. The relative role of SCT versus immunosuppressive therapy in individuals with acquired aplastic anaemia is under constant review. In general, SCT is favoured in patients less than 35 years old with severe aplastic anaemia and a human leucocyte antigen (HLA) matching sibling donor. Cure rates of up to 80% are obtained. However, non-myeloablative transplants (p. 253) are used in selected patients over the age of 40 years. SCT using umbilical cord blood, unrelated volunteer

donors and mismatched family members are used in carefully selected patients. In older subjects and those with less severe disease, immunosuppression is tried first.

7 Haemopoietic growth factors Granulocyte colony-stimulating factor (G-CSF) may produce minor responses but does not lead to sustained improvement. Other growth factors have not proved helpful.

8 Iron chelation therapy This may be needed in patients who require regular transfusion.

Paroxysmal nocturnal haemoglobinuria (PNH)

PNH is a rare, acquired, clonal disorder of marrow stem cells in which there is **deficient synthesis of the glycosylphosphatidylinositol (GPI) anchor**, a structure that attaches several surface proteins to the cell membrane. There is a clinical triad of chronic intravascular haemolysis, venous thrombosis and bone marrow failure. It results from acquired mutations in the X chromosome gene coding for phosphatidylinositol glycan protein class A (PIG-A), which is essential for the formation of the GPI anchor. The net result is that GPI-linked proteins (such as CD55 and CD59) are absent from the cell surface of all the cells derived from the abnormal stem cell (Fig. 22.5). CD55 and CD59 are also normally present on white cells and platelets. The lack of the surface molecules decay-activating factor (DAF, CD55) and membrane inhibitor of reactive lysis (MIRL, CD59) render red cells sensitive to lysis by complement and the result is chronic intravascular haemolysis.




Figure 22.5 Schematic representation of the phosphatidylinositol glycan which anchors many different proteins to the cell membrane, e.g. CD59 (MIRL, membrane inhibitor of reactive lysis).

Haemosiderinuria is a constant feature and can give rise to iron deficiency which may exacerbate the anaemia. Haptoglobins are absent; free haemoglobin may damage the kidney and it removes nitric oxide from smooth muscle causing dysphagia and pulmonary hypertension. The other main clinical problem in PNH is of venous thrombosis. Patients may develop recurrent thromboses of large vessels, including the portal, hepatic and mesenteric veins. Arterial thrombosis, such as strokes or myocardial infarction, can also occur. Intermittent abdominal pain due to mesenteric vein thrombosis is a common feature.

PNH is almost invariably associated with some form of bone marrow hypoplasia and there may even be complete aplastic anaemia. It appears that the PNH clone may expand as a result of a selective pressure, possibly immunologically mediated, against cells that have normal GPI-linked membrane proteins.

PNH is diagnosed by flow cytometry which shows loss of expression of the GPI-linked proteins CD55 and CD59. This has replaced the demonstration of red cell lysis in serum at low pH – the Ham test.

Eculizumab, a humanized antibody against complement C5, inhibits the activation of terminal components of complement and reduces haemolysis, transfusion requirements and the incidence of thrombosis. Iron therapy is used for iron deficiency and long-term anticoagulation with warfarin may be needed. Immunosuppression can be useful and allogeneic stem cell transplantation is a definitive treatment. The disease occasionally remits spontaneously. The median survival is over 10 years. As for aplastic anaemia, transformation to MDS or AML may occur.

Red cell aplasia

Chronic form

This is a rare syndrome characterized by anaemia with normal leucocytes and platelets and grossly reduced or absent erythroblasts in the marrow (Fig. 22.6). The congenital form is known as **Diamond–Blackfan syndrome** (Table 22.3) and is




Figure 22.6 The bone marrow in primary red cell aplasia. There is selective loss of erythropoiesis.

Table 22.3 Classification of pure red cell aplasia.

Acute, transient	Chronic congenital	Chronic acquired
Parvovirus infection Infancy and childhood Drugs (e.g. azathioprine, co-trimoxazole)	Diamond–Blackfan syndrome	Idiopathic Associated with thymoma, systemic lupus erythematosus, rheumatoid arthritis, lymphoma, chronic lymphocytic leukaemia, T-cell large granular lymphocytosis, myelodysplasia, viral infection, drugs

inherited as a recessive condition. It is associated with a varying number of somatic abnormalities (e.g. of the face or heart). Mutation of a gene on chromosome 19 or other genes that encode ribosomal proteins underlies most cases (Fig. 22.3). Corticosteroids are the first line of treatment and SCT may be curative. Androgens may also produce improvement but side-effects on growth can be severe. Iron chelation is needed after multiple transfusions.

The acquired chronic form can occur without any obvious associated disease or precipitating factor (idiopathic), or may be seen with autoimmune diseases (especially systemic lupus erythematosus), with a thymoma, lymphoma or chronic lymphocytic leukaemia. In some cases, immunosuppression with corticosteroids, rituximab, ciclosporin, azathioprine or ATG is helpful. Monoclonal antibodies, such as rituximab

(anti-CD20), are being used increasingly in treatment of refractory acquired red cell aplasia and other autoimmune cytopenias.

If regular blood transfusions are needed, iron chelation therapy will also be necessary. SCT has been carried out in some severe cases.

Transient form

Parvovirus B19 infects red cell precursors via the P antigen and causes a transient (5–10 days) red cell aplasia. This can result in rapid onset of severe anaemia in patients with pre-existing shortened red cell survival, such as those with sickle cell disease or hereditary spherocytosis (Fig. 22.7). Transient red cell aplasia with anaemia may also occur in association with drug therapy




Figure 22.7 Parvovirus infection: flow chart showing transient fall in haemoglobin and reticulocytes in a patient with hereditary spherocytosis.

(Table 22.3), and in normal infants or children, often with a history of a viral infection in the preceding 3 months.

Schwachman–Diamond syndrome

This is a rare autosomal recessive syndrome characterized by varying degrees of cytopenia, especially neutropenia with a propensity to transform to myelodysplasia or acute myeloid leukaemia. Exocrine pancreatic dysfunction is an invariable feature, while skeletal abnormalities, hepatic impairment and short stature are frequent. The disorder results from inherited mutations in the gene *SD*, involved in ribosome synthesis (Fig. 22.3).

Congenital dyserythropoietic anaemia

Congenital dyserythropoietic anaemias (CDAs) are a group of hereditary refractory anaemias characterized by ineffective erythropoiesis and erythroblast multinuclearity. The patient may be jaundiced with bone marrow expansion. The white cell and platelet counts are normal. The reticulocyte count is low for the degree of anaemia, despite increased marrow cellularity. The anaemia is of variable severity and is usually first noted in infancy or childhood. Iron overload may develop and

splenomegaly is common. The CDAs have been classified into four types based on the degree to which megaloblastic changes, giant erythroblasts and dyserythropoietic changes are present. CDA1 is associated with somatic abnormalities. It is due to mutation of the gene *CDANI*, active during the S phase of the cell cycle. Type II is known as HEMPAS (hereditary erythroblast multinuclearity with a positive acidified serum lysis test). The test is positive with some sera but not with the patient's serum. The basic lesion in HEMPAS is a defect in a gene *SEC23B* coding for a protein involved in synthesis of endoplasmic reticulum derived vesicles destined for the Golgi component. α -Interferon has induced remission in some cases.

Osteopetrosis


This is a rare heterogeneous group of disorders due to failure of bone resorption by osteoclasts. Inheritance may be recessive or dominant. The bones are dense but brittle and fractures are common. The marrow space is reduced and a leucoerythroblastic anaemia occurs. The liver and spleen are enlarged. Early death from the consequences of bone marrow failure is usual. SCT offers a chance of cure.

SUMMARY

- Aplastic anaemia presents as pancytopenia (subnormal haemoglobin, neutrophils and platelets) associated with a hypoplastic bone marrow.
- Aplastic anaemia may be congenital (Fanconi) or acquired (idiopathic or due to drugs, viral infection or toxins).
- Fanconi anaemia is autosomal recessive, associated with congenital skeletal, skin or renal abnormalities. It is caused by inherited mutations of genes involved in DNA repair.
- Acquired aplastic anaemia is treated with immunosuppressive drugs (e.g. antithymocyte globulin, ciclosporin) or by stem cell transplantation.
- Paroxysmal nocturnal haemoglobinuria is an acquired clonal haemolytic anaemia associated with pancytopenia arising in a hypoplastic marrow. There is defective synthesis of the glycosylphosphatidylinositol anchor for many membrane proteins.
- Red cell aplasia causes anaemia with normal white cell and platelet counts. It may be transient, usually caused by parvovirus infection, or chronic. This may be congenital (Diamond–Blackfan) or acquired, associated with, e.g. systemic lupus erythematosus, lymphoma or chronic lymphocytic leukaemia.
- Congenital dyserythropoietic anaemias are a group of rare inherited disorders of erythropoiesis.



Now visit www.wileyessential.com/haematology to test yourself on this chapter.


A vertical strip on the left side of the page showing a microscopic view of various blood cells, including red blood cells and white blood cells, in shades of red, pink, and yellow against a dark blue background.

CHAPTER 23

Stem cell transplantation

Key topics

■ Principles of stem cell transplantation	251
■ Peripheral blood stem cell collection	251
■ Bone marrow stem cell collection	251
■ Myeloablative and non-myeloablative conditioning	253
■ Autologous stem cell transplantation	254
■ Allogeneic stem cell transplantation	255
■ Human leucocyte antigen (HLA) system	255
■ Complications	256
■ Graft-versus-leukaemia and donor leucocyte infusions	261



Principles of stem cell transplantation

Stem cell transplantation (SCT) involves eliminating a patient's haemopoietic and immune system by chemotherapy and/or radiotherapy and replacing it with stem cells either from another individual or with a previously harvested portion of the patient's own haemopoietic stem cells (Fig. 23.1). Depending on the source of the stem cells the term includes **bone marrow transplantation** (BMT), **peripheral blood stem cell** (PBSC) **transplantation** and umbilical cord transplantation. Depending on the type of donor, SCT may be **syngeneic**, **allogeneic** or **autologous** (Table 23.1).

The principal diseases for which SCT is performed are listed in Table 23.2. However, the exact role of SCT in the management of each disease is complex and depends on factors such as disease severity and subtype, remission status, age and, for allogeneic transplantation, availability of a donor.

Collection of stem cells

Stem cells can be collected from the peripheral blood, bone marrow or umbilical cord blood.

Peripheral blood stem cell collection

This is now the preferred source of stem cells for both autologous and allogeneic transplantation. Peripheral blood stem cells (PBSCs) are taken using a cell-separator machine connected to the patient or donor via peripheral cannulae (Fig. 23.2). Blood is taken through one cannula and pumped around the machine where mononuclear cells are collected by centrifugation before the red cells are returned to the patient. This continuous process may take a few hours before enough mononuclear cells are collected.

Peripheral blood normally contains too few haemopoietic stem cells to allow collection of sufficient numbers for transplantation. Growth factors can increase the number by around 10–100 times. Granulocyte colony-stimulating factor (G-CSF) is given to patients or donors as a course of injections (typically 10 µg/kg/day for 4–6 days) until the white cell count starts to rise. Plerixafor, an inhibitor of stem cell adhesion in the bone marrow, is also given if mobilisation is likely to be inadequate. PBSC collections are then taken and, depending on the efficiency of stem cell mobilization, repeated collections may be needed for up to 3 days. A typical out-patient protocol is G-CSF on days 1–4 and harvests on days 5 and 6. The adequacy of the collection is assessed by CD34⁺ cell count. Generally greater than $2.0 \times 10^6/\text{kg}$ CD34⁺ cells are needed for transplantation.

Bone marrow collection

The donor is given a general anaesthetic and 500–1200 mL of marrow is harvested from the pelvis. The marrow is anticoagulated and a mononuclear cell count is taken to assess the yield, which should be approximately $2\text{--}4 \times 10^8$ nucleated cells/kg body weight of the recipient.

Table 23.1 Stem cell transplantation: potential donors.

HLA-matching sibling	} Allogeneic
Unrelated HLA-matching volunteer	
Umbilical cord blood	
Identical twin	Syngeneic
Self	Autologous
HLA, human leucocyte antigen.	

Table 23.2 Stem cell transplantation: indications.

Allogeneic (or syngeneic)	Autologous
Acute lymphoblastic or myeloid leukaemia. Other malignant disorders of the marrow (e.g. chronic myeloid leukaemia, myelodysplasia, multiple myeloma, lymphoma, severe aplastic anaemia)	Hodgkin and non-Hodgkin lymphoma, multiple myeloma, primary amyloidosis
Inherited disorders: thalassaemia major, sickle cell anaemia, immune deficiencies, inborn errors of metabolism in the haemopoietic and mesenchymal system (e.g. osteopetrosis) Other acquired severe marrow diseases (e.g. paroxysmal nocturnal haemoglobinuria, red cell aplasia, myelofibrosis)	Amyloidosis For 'gene therapy' of genetic disease (e.g. adenosine deaminase deficiency)

Umbilical cord blood

Fetal blood is a rich source of haemopoietic stem cells which may be collected from cord blood. Because of the relatively small numbers of stem cells collected from a single cord, they are most useful for children who do not have a fully matching sibling or unrelated donor. Less stringent HLA-matching is needed. Double cord donations may be needed to obtain sufficient stem cells for adult recipients. Immune reconstitution is slower after cord blood transplantation.

Stem cell processing

After collection, the stem cell harvest can be processed with removal of red cells and concentration of the mononuclear cells. In some protocols antibodies to remove T lymphocytes are used to reduce the risk of graft-versus-host disease (GVHD).




Figure 23.1 Procedures for **(a)** allogeneic, and **(b)** autologous stem cell transplantation. G-CSF, granulocyte colony-stimulating factor; GVHD, graft-versus-host disease.



Figure 23.2 Peripheral blood stem cell (PBSC) collection: a donor undergoing collection of PBSCs on a cell separator.

Autologous collections may be ‘purged’ by chemotherapy or antibodies in an attempt to remove residual malignant cells. CD34⁺ stem cells may be selected from both types of harvest (Fig. 23.3).

Conditioning

Prior to infusion of haemopoietic stem cells, patients receive chemotherapy, sometimes in combination with total body irradiation (TBI; Fig. 23.1) in a procedure called **conditioning**. This is designed to eradicate the patient’s haemopoietic and immune system and, if present, malignancy. In addition, in the setting of allogeneic SCT, by suppressing the host immune system, it helps to prevent rejection of the ‘foreign’ stem cells. An important development that has occurred in SCT is a




Figure 23.3 Peripheral blood stem cell collection: enriched CD34⁺ cells stained by May–Grünwald–Giemsa stain. The cells have the appearance of small and medium-sized lymphocytes.

major shift from **myeloablative** regimes to **non-myeloablative** conditioning.

Myeloablative conditioning regimens irreversibly destroy the haemopoietic function of the bone marrow with high doses of chemotherapy or radiotherapy. TBI is usually used in patients with malignant disease and is usually administered over several days (*fractionated*). The most commonly used chemotherapy drug is cyclophosphamide but busulfan (preferably intravenously), melphalan, or other drugs are given in some protocols. Treosulfan is a newer alkylating agent in trials. Before autologous transplantation, high-dose melphalan is used for myeloma and a combination regimen, e.g. BEAM (carmustine, etoposide, cytarabine and melphalan) for lymphoma. For benign disorders TBI is avoided and cyclophosphamide, busulfan and fludarabine with alemtuzumab or ATG *in vivo* are employed in different protocols.

At least 36 hours are allowed for the elimination of the drugs from the circulation following the last dose of chemotherapy before donor stem cells are infused. Conditioning therapy is often complicated by mucositis and patients sometimes need parenteral nutrition. T cell depletion of the donor stem cells *in vitro* reduces the risk of GVHD but increases the risk of non-engraftment, infections and relapse. Alternatively, T cell depletion of the host *in vivo* by antithymocyte globulin (ATG) or alemtuzumab (anti-CD52) may be employed to reduce GVHD risk, but may increase risk of infection and relapse.

Non-myeloablative conditioning regimens have been developed to reduce the morbidity and mortality of allogeneic transplantation. These do not completely destroy the host bone marrow. These include agents such as fludarabine, low doses of busulfan or cyclophosphamide, low-dose irradiation, ATG or other antibodies that delete T cells. The aim in these mini or low-intensity transplants is to use enough immunosuppression to allow donor stem cells to engraft without completely eradicating host marrow stem cells. Donor leucocyte infusions (DLI) are commonly used at a later stage in order to encourage complete donor engraftment and to enhance the graft-versus-leukaemia or lymphoma effect. Such regimes extend the age range and increase the treatment indications for allogeneic transplantation.

Post-transplant engraftment and immunity

After a period of typically 1–3 weeks of severe pancytopenia, the first signs of successful engraftment are monocytes and neutrophils in the blood with a subsequent increase in platelet count (Fig. 23.4). A reticulocytosis also begins. G-CSF may be used to reduce the period of neutropenia. Engraftment is usually quicker following PBSC transplantation compared with BMT.

The marrow cellularity gradually returns to normal but the marrow reserve remains impaired for 1–2 years. There is profound immunodeficiency for 3–12 months with a low level of CD4 helper cells. Immune recovery is quicker after




Figure 23.4 Typical haematological chart of a patient undergoing allogeneic marrow transplantation for aplastic anaemia. WBC, white blood cells.

autologous and syngeneic SCT than following allogeneic SCT. The patient's blood group changes to that of the donor and antigen-specific immunity becomes that of the donor after approximately 60 days.

Autologous stem cell transplantation

This allows the delivery of a high dose of chemotherapy, with or without radiotherapy, which otherwise would result in prolonged bone marrow aplasia. Stem cells are harvested and

stored before the treatment is given and are then reinfused to rescue the patient from the myeloablative effects of the treatment (Fig. 23.1). A limitation of the procedure is that tumour cells contaminating the stem cell harvest may be reintroduced into the patient. Nevertheless, autografting has a major role in the treatment of haematological diseases such as lymphoma and myeloma. The major problem associated with autografting is recurrence of the original disease (Fig. 23.5). GVHD is not an issue. Procedure-related mortality is generally well below 5%.





Figure 23.5 The causes of death following autologous, HLA matched sibling and unrelated allogeneic transplantation. IPn, interstitial pneumonitis. Source: Craddock C. & Chakraverty R. In Hoffbrand A.V. *et al.* (eds) (2016) *Postgraduate Haematology 7e*. Reproduced with permission of John Wiley & Sons.

Allogeneic stem cell transplantation


In this procedure, stem cells harvested from another person are infused into the patient. The procedure has a significant morbidity and mortality and one of the major reasons is the immunological incompatibility between donor and patient despite matching of the human leucocyte antigens (HLA). This may manifest as immunodeficiency, GVHD or graft failure (Fig. 23.5). Paradoxically, there is also a graft-versus-leukaemia (GVL) effect which probably underlies much of the success of the procedure.

The human leucocyte antigen (HLA) system

Allografting would be impossible without the ability to perform HLA typing. The short arm of chromosome 6 contains a cluster of genes known as the major histocompatibility complex (MHC) or the HLA region (Fig. 23.6a). Genes in this



(a)



(b)

Figure 23.6 (a) The human leucocyte antigen (HLA) complex. (b) HLA class I and II molecules showing protein domains and bound peptide.

Table 23.3 The human leucocyte antigens.

	Class I	Class II
Antigens	HLA-A, -B, -C	HLA-DR, -DP, -DQ
Distribution	All nucleated cells, platelets	B lymphocytes Monocytes Macrophages Activated T cells
Structure	Large polypeptide chain) and a β_2 -microglobulin	Two polypeptide chains (α and β)
Interacts with	CD8 lymphocytes	CD4 lymphocytes

region encode the HLA antigens and many other molecules, including complement components, tumour necrosis factor (TNF) and proteins associated with antigen processing. HLA proteins are divided into two types: class I and II (Table 23.3). Their role is to bind intracellular peptides and 'present' these to T lymphocytes for antigen recognition (see Chapter 9). Class I molecules (HLA-A, -B and -C) present antigen to CD8⁺ T cells and class II molecules (HLA-DR, -DQ and -DP) present to CD4⁺ T cells (Fig. 23.6b).

Class I antigens are present on most nucleated cells and on the cell surface they are associated with β_2 -microglobulin. **Class II antigens** have a more restricted tissue distribution and comprise α and β chains, both encoded by genes in the HLA region. The inheritance of the four loci (HLA-A, -B, -C and -DR) is closely linked, one set of loci is inherited from each parent so that there is approximately a one in four chance of two siblings having identical HLA antigens (Fig. 23.7a). Crossing-over of genes during meiosis accounts for occasional unexpected disparities. The inheritance is independent of sex or blood group.

Human leucocyte antigen and transplantation

The natural role of HLA molecules is in directing T-lymphocyte responses and the greater the HLA mismatch the more severe is the immune response between transplanted cells. HLA typing is critical in donor selection for allogeneic SCT.

Minor histocompatibility antigens, e.g. HA-1, HA-2, Hy, are peptides that are presented by HLA molecules and are able to act as antigens in SCT either because they are polymorphic in the population or because they are encoded on the Y chromosome and therefore represent novel antigens when a female immune system engrafts in a male. They are likely to be important antigens in GVHD and the GVL reaction (see below).

HLA typing may be carried out by serological or molecular techniques.

The nomenclature for *HLA* alleles is standardized. A single antigenic specificity defined by serological typing (e.g. HLA-A2) can be divided into different alleles by DNA sequencing. Each allele is given in numerical designation. The gene name is followed by an asterisk. The first field of digits indicates the allele group. The second field of digits lists subtypes, which differ

in the protein sequence. Subsequent digits indicate minor differences in non-coding regions. As an example, alleles at the *HLA-A* loci are written as *HLA-A*01:01* to *HLA-A*80:01*. The nomenclature for the class II genes is similar but complicated by the fact that there may be more than one *HLA-DRB* gene on each chromosome (Fig. 23.7b).

When searching for an unrelated donor the aim is to match HLA-A, -B and -DR between recipient and donor and this is then called a 10/10 match based on A, B, C, DRB1 and DQB1 matching. An exception is when a donor with only a single HLA haplotype match, usually a parent or sibling, is used in a **haploidentical SCT**. Such transplants usually require a stem cell graft that is heavily depleted of T cells in order to limit the development of GVHD. There are many millions of volunteer donors registered on International Registries and the chance of identifying a matched unrelated donor for a patient lacking an HLA identical sibling (depending on the ethnic group) is usually greater than 50%.

Chimaerism analysis

Following allogeneic SCT, the recipient's blood shows the presence of both donor and recipient cells (chimaerism). This can be detected by FISH analysis (see Chapter 11) of the proportion of Y chromosome containing cells if there is a sex mismatch or by DNA analysis techniques.

Complications (Table 23.4)

The causes of death following SCT are shown in Figure 23.5. The overall rate from the procedure itself is lowest (less than 5%) for autologous SCT and highest in unrelated and haploidentical SCT.

Graft-versus-host disease (GVHD)

This is caused by donor-derived immune cells, particularly T lymphocytes, reacting against recipient tissues. Its incidence is increased with increasing age of donor and recipient and if there is any degree of HLA mismatch between them. Donor alloimmunization, e.g. a female who has had multiple pregnancies, and, in the recipient, viral infection (e.g. CMV), liver, inflammatory bowel or rheumatological disease, also increase




Figure 23.7 (a) An example of the possible pattern of inheritance of the A, B and DR (*DRB1*) series alleles of the human leucocyte antigen (HLA) complex. (b) Molecular genetics of the HLA class II gene complex. There are four major haplotypes of MHC class II genes in the population and each individual may have up to two (one on each chromosome). The *DRA1* gene codes for the DRα protein and the *DRB1*, *DRB3*, *DRB4* and *DRB5* genes encode DRβ chains. Expression from the *DRB1* gene is higher than from the other genes. The number of alleles at each gene is shown underneath the gene in italics. Alleles at each locus have a standard nomenclature (e.g. the alleles at the *DRB1* gene are termed *DRB1*0101* to *DRB1*1608*). It is now known that the DR51, DR52 and DR53 antigens, which are defined by serological testing, are encoded from the *DRB5*, *DRB4* and *DRB3* genes, respectively.

the risk of acute GVHD. GVHD prophylaxis is usually given as ciclosporin (or tacrolimus), with methotrexate (or sirolimus or mycophenolate mofetil).

In **acute pattern GVHD**, usually occurring in the first 100 days, the skin, gastrointestinal tract or liver are affected (Table 23.5). The skin rash typically affects the face, palms,

soles and ears but may, in severe cases, affect the whole body (Fig. 23.8). Diarrhoea may lead to fluid and electrolyte depletion. Typically, bilirubin and alkaline phosphatase are raised but the other hepatic enzymes are relatively normal. Acute GVHD is usually treated by high doses of corticosteroids which are effective in the majority of cases.

Table 23.4 Complications of stem cell transplantation.

Early (usually <100 days)	Late (usually >100 days)
Infections, especially bacterial, fungal, herpes simplex virus, CMV	Infections, especially varicella-zoster, capsulate bacteria
Haemorrhage	Chronic pattern GVHD (arthritis, malabsorption, hepatitis, scleroderma, sicca syndrome, lichen planus, pulmonary disease, serous effusions)
Acute pattern GVHD (skin, liver, gut)	Chronic pulmonary disease
Graft failure	Autoimmune disorders
Haemorrhagic cystitis	Cataract
Interstitial pneumonitis	Infertility
Others: veno-occlusive disease, cardiac failure	Second malignancies

CMV, cytomegalovirus; GVHD, graft-versus-host disease.

Table 23.5 Acute pattern graft-versus-host disease: clinical staging (Seattle system).

Stage	Skin	Liver (bilirubin, $\mu\text{mol/L}$)	Gut (diarrhoea, L/day)
I	Rash <25%	20–35	0.5–1.0
II	Rash 25–50%	35–80	1.0–1.5
III	Erythroderma	80–150	1.5–2.5
IV	Bullae, desquamation	>150	>2.5; severe pain, ileus

In **chronic pattern GVHD**, which usually occurs after 100 days and may evolve from acute GVHD, these tissues are involved, but also the joints and other serosal surfaces, the oral mucosa and lacrimal glands. Features of autoimmune disease

with scleroderma, Sjögren's syndrome and lichen planus may develop. The immune system is impaired (including hyposplenism) with risk of infection. Malabsorption and pulmonary abnormalities are frequent. Corticosteroids are tried with second-line drugs, including ciclosporin, rituximab, sirolimus, mycophenolate mofetil and extracorporeal photopheresis. The response may be poor.

Infections

In the early post-transplant period, bacterial or fungal infections are frequent (Fig. 23.9). These may be reduced by reverse barrier nursing with laminar or positive-pressure air flow and the use of skin and mouth antiseptics. Prophylactic therapy with aciclovir, antifungal agents and oral antibiotics is often given. If a fever or other evidence of an infection occurs, broad-spectrum intravenous antibiotics are commenced immediately after blood cultures and other appropriate microbiological specimens have been taken. Failure of response to antibacterial agents is usually an indication to commence systemic antifungal therapy with amphotericin B, caspofungin or voriconazole. Fungal infections, especially *Candida* and *Aspergillus* species (see Chapter 12) are a particular problem because of the prolonged neutropenia.



Figure 23.8 Widespread erythematous skin rash in acute graft-versus-host disease following allogeneic stem cell transplantation.

Viral infections, particularly with the herpes group of viruses, are frequent with herpes simplex, cytomegalovirus (CMV) and varicella zoster virus (VZV) occurring at different peak intervals (Fig. 23.9). CMV presents a particular threat and is associated with a potentially fatal interstitial pneumonitis (Fig. 23.10) as well as with hepatitis and falling blood counts. The infection may be caused by reactivation of CMV in the recipient or a new infection transmitted by the donor. In CMV-seronegative patients with CMV-seronegative donors, CMV-negative, leucodepleted blood products must be given. Aciclovir may be useful in prophylaxis. Most centres screen patients regularly for evidence of CMV reactivation following allogeneic transplantation. If these tests become positive, ganciclovir may suppress the virus before disease occurs. Ganciclovir, foscarnet, cidofovir (a newer powerful but nephrotoxic agent) and CMV immunoglobulin may be tried for established CMV infection. VZV infection is also frequent post-SCT but occurs later with a median onset at 4–5 months. Rarely, disseminated VZV infection occurs. Intravenous aciclovir is indicated. Epstein–Barr

virus (EBV) infections and EBV-associated lymphoproliferative disease (see Fig 23.13) are less frequent after SCT than after solid organ transplants.

Pneumocystis jirovecii (Fig 23.10) is cause of pneumonitis that may be prevented by prophylactic co-trimoxazole.

Interstitial pneumonitis

This is one of the most frequent causes of death post-SCT (Fig. 23.10). CMV is a frequent agent but other herpes viruses and *P. carinii* account for other cases; in most cases, no cause other than the previous radiation and chemotherapy can be implicated. Bronchoalveolar lavage or open lung biopsy may be needed to establish the diagnosis.

Blood product support

Platelet concentrates are given to maintain a count of $10 \times 10^9/L$ or more. Platelets and blood transfusions given in the post-transplant period must be irradiated prior to administration in order to kill any lymphocytes that might cause GVHD.




Figure 23.9 Time sequence for development of different types of infection following allogeneic stem cell transplantation. CMV, cytomegalovirus; Gr+, Gr–, Gram-positive or -negative; GVHD, graft-versus-host disease; HSV, herpes simplex virus.




Figure 23.10 (a) Chest radiograph showing interstitial pneumonitis following bone marrow transplantation. Widespread diffuse mottling can be seen. The patient had received total body irradiation and had grade III graft-versus-host disease. No infective cause of the pneumonitis was identified. Possible causes include pneumocystis, cytomegalovirus, herpes zoster, fungal infection or a combination of these. (b) Sputum cytology: intranuclear CMV inclusion body in a pulmonary cell. Papanicolaou stain. (c) *Pneumocystis jirovecii* in bronchial washings, Gram–Weigert stain.

Other complications of allogeneic transplantation

Graft failure

The risk of graft failure is increased if the patient has aplastic anaemia or if T-cell depletion of donor marrow is used as GVHD prophylaxis. This suggests that donor T cells are needed to overcome host resistance to engraftment of stem cells.

Haemorrhagic cystitis

This is caused by the cyclophosphamide metabolite acrolein. Mesna is given in an attempt to prevent this. Certain viruses (e.g. adenovirus or polyomavirus) may also cause this complication.

Other complications

These include veno-occlusive disease of the liver (manifest as jaundice, hepatomegaly and ascites or weight gain) and cardiac failure as a result of the conditioning regime (especially high doses of cyclophosphamide) and previous

chemotherapy on the heart. Haemolysis because of ABO incompatibility between donor and recipient may cause problems in the first weeks. Microangiopathic haemolytic anaemia may also occur.

Late complications

Relapse of the original disease (e.g. acute or chronic leukaemia) may occur (Fig. 23.5). Bacterial infections are frequent, especially with encapsulated organisms affecting the respiratory tract. Oral penicillin is given prophylactically to reduce this risk. VZV and fungal infections are also frequent. The use of prophylactic co-trimoxazole and oral aciclovir for 3–6 months reduces the risk of *Pneumocystis* and herpes infections, respectively.

Delayed pulmonary complications include restrictive pneumonitis and bronchiolitis obliterans. Endocrine complications include hypothyroidism, growth failure in children, impaired sexual development and infertility. These endocrine problems

are more marked if TBI has been used. Clinically apparent autoimmune disorders are infrequent and include myasthenia, rheumatoid arthritis, anaemia, thrombocytopenia or neutropenia. Autoantibodies are frequently detected in the absence of symptoms. Second malignancies (especially non-Hodgkin lymphoma) occur with a six- or seven-fold incidence compared with controls. CNS complications include neuropathies and eye problems caused by chronic GVHD (sicca syndrome) or cataracts. Iron overload may be present due to repeated red cell transfusions before and after transplantation. This can be treated by iron chelation or venesections when there is adequate red cell recovery. It is not necessary to 'de-iron' before the transplant procedure.

Graft-versus-leukaemia effect and donor leucocyte infusions

After allogeneic transplantation the donor immune system helps to eradicate the patient's leukaemia, a phenomenon known as the **graft-versus-leukaemia** (GVL) effect. Evidence includes the decreased relapse rate in patients with GVHD, the increased relapse rate in identical twins and, most convincingly,

the ability of **donor leucocyte infusions** (DLI) to cure relapsed leukaemia in some patients. Graft-versus-lymphoma and -myeloma effects also exist. The principle of DLI is that peripheral blood mononuclear cells are collected from the original allograft donor and directly infused into the patient at the time of leukaemia relapse (Fig. 23.11).

There is a large difference in the outcome of different diseases treated by DLI. Chronic myeloid leukaemia (CML) is most sensitive whereas acute lymphoblastic leukaemia rarely responds. In CML the response to DLI is better in cases of early relapse. PCR is used to monitor serial blood samples for evidence of recurrence of the *BCR-ABL1* transcript before karyotypic or clinical relapse occurs (Fig. 23.11). DLI can then be used in cases of molecular relapse. The response to DLI may take several weeks but usually results in a permanent cure. The mechanism is unclear but a T-cell-mediated alloreactive immune response is likely to be a major component.

Positron emission tomography (PET) scans can be used to detect residual disease in cases of lymphoma and to guide the requirement for DLI and determining the disease response (Fig. 23.12).




Figure 23.11 Donor leucocyte infusions. Example of donor leucocyte infusion (DLI) in the treatment of chronic myeloid leukaemia (CML) which relapsed following allogeneic stem cell transplantation (SCT). Polymerase chain reaction (PCR) analysis of the blood for the *BCR-ABL1* transcript shows that there was transient loss of the transcript but molecular and cytogenetic relapse occurred at 10 months. One infusion of donor leucocytes led to re-establishment of a durable complete remission.




Figure 23.12 Example of disease control following administration of a donor leucocyte infusion (DLI) after stem cell transplantation. **(a)** A PET scan revealed residual disease activity in a patient at 6 months following allogeneic transplantation for non-Hodgkin lymphoma. The bright signals reflect the metabolic activity of malignant cells in the spleen and axillary lymph nodes. Donor leucocyte infusion was then given and after 3 months a repeat PET scan **(b)** revealed no evidence of residual disease. Source: Courtesy of Professor Nigel Russell.

Post-transplant lymphoproliferative diseases

These are polyclonal or monoclonal lymphoid proliferation that occurs in recipients of stem cell or more frequently solid organ allografts, as a result of the intensive immunosuppression. There may be Epstein–Barr (EB) virus driven lymphocytosis or

lymphoma, usually B cell. There is often involvement of bowel, lung or bone marrow (Fig. 23.13). Treatment is by withdrawing immunosuppression (if feasible) anti-CD20 (rituximab) and if appropriate chemotherapy or cytotoxic T cells engineered to kill the EBV-positive tumour cells.





Figure 23.13 Post transplantation lymphoproliferative diseases: 17-year-old male 5 months after renal transplantation had small bowel perforation caused by diffuse large B-cell lymphoma. **(a)** Low-power view of lymphoid mass invading small bowel. **(b)** High-power view of lymphoid mass. **(c)** Immunostaining for CD20. **(d)** EBV-ISH (in situ hybridization) stain showing the tumour cells are positive for EBV. Source: Courtesy of Professor P. Amrolia and Dr. N. Sebire.

- Stem cell transplantation (SCT) involves replacing the haemopoietic and immune systems by stem cells from either the same subject (autologous) or another individual (allogeneic). The donor stem cells can be harvested from bone marrow, peripheral blood or umbilical cord.
- Autologous SCT is most frequently performed for lymphomas or myeloma.
- For allogeneic SCT the recipient's own haemopoietic and immune systems are eliminated by chemotherapy, radiotherapy and monoclonal antibodies.
- Allogeneic SCT requires a tissue (HLA) matching sibling or unrelated donor. The human leucocyte antigens (HLA) are coded for by genes on chromosome 6. They are extremely polymorphic and are involved in presentation of antigens to T lymphocytes.
- Allogeneic SCT is indicated in selected cases of acute leukaemia, other malignant bone marrow diseases, and severe acquired or genetic marrow diseases (e.g. aplastic anaemia, thalassaemia major). Reduced-intensity conditioning SCT is preferred in older subjects.
- Donor leucocyte infusions may be given to treat relapse of leukaemia post-allogeneic SCT, by a 'graft-versus-leukaemia' effect.
- Early complications of allogeneic SCT include acute graft-versus-host disease, infections, graft failure and veno-occlusive disease. Long-term complications include chronic graft-versus-host disease, damage to many different organs (e.g. skin, heart, lungs and liver) and post-transplant lymphoproliferative disease.



Now visit www.wileyessential.com/haematology to test yourself on this chapter.


A vertical strip on the left side of the page showing a microscopic view of blood cells. It includes several large, red, biconcave disc-shaped red blood cells, smaller yellowish spherical cells with granules (leukocytes), and very small, dark purple specks (platelets).

CHAPTER 24

Platelets, blood coagulation and haemostasis

Key topics

■ Platelets	265
■ Blood coagulation	270
■ Endothelial cells	273
■ Haemostatic response	273
■ Fibrinolysis	275
■ Tests of haemostatic function	276



The normal haemostatic response to vascular damage depends on a closely linked interaction between the blood vessel wall, circulating platelets and blood coagulation factors (Fig. 24.1).

An efficient and rapid mechanism for stopping bleeding from sites of blood vessel injury is clearly essential for survival. Nevertheless, such a response needs to be tightly controlled to prevent extensive clots developing and to be able to break down such clots once damage is repaired. The haemostatic system thus represents a delicate balance between procoagulant and anticoagulant mechanisms allied to a process for fibrinolysis. The five major components involved are platelets, coagulation factors, coagulation inhibitors, fibrinolysis and blood vessels.

Platelets

Platelet production

Platelets are produced in the bone marrow by fragmentation of the cytoplasm of megakaryocytes, one of the largest cells in the body. The precursor of the megakaryocyte – the megakaryoblast – arises by a process of differentiation from the haemopoietic stem cell (see Fig. 1.2). The megakaryocyte matures by endomitotic synchronous replication (i.e. DNA replication in

the absence of nuclear or cytoplasmic division) enlarging the cytoplasmic volume as the number of nuclear lobes increases in multiples of two (Fig. 24.2). Early on invaginations of plasma membrane are seen, called the demarcation membrane, which evolves through the development of the megakaryocyte into a highly branched network. At a variable stage in development the cytoplasm becomes granular. Mature megakaryocytes are extremely large, with an eccentrically placed single lobulated nucleus and a low nuclear:cytoplasmic ratio (Fig. 24.3). Platelets form by fragmentation from the tips of cytoplasmic extensions of megakaryocyte cytoplasm, each megakaryocyte giving rise approximately to 1000–5000 platelets (Fig. 24.3c). The platelets are released through the endothelium of the vascular niches of the marrow where megakaryocytes reside. The time interval from differentiation of the human stem cell to the production of platelets averages 10 days.

Thrombopoietin (TPO) is the major regulator of platelet formation and 95% is produced by the liver. Approximately 50% is produced constitutively, the plasma level depending on its removal from plasma by binding to c-MPL receptors on platelets and megakaryocytes (Fig 24.4a). Therefore, levels are high in thrombocytopenia as a result of marrow aplasia but low in patients with raised platelet counts. The other 50% is




Figure 24.1 The involvement of blood vessels, platelets and blood coagulation in haemostasis. ADP, adenosine diphosphate.




Figure 24.2 Simplified diagram to illustrate platelet production from megakaryocytes.

regulated in response to platelet destruction. As platelets age they lose surface sialic acid. This exposes galactose residues that attach to the Ashwell–Morell receptor in the liver. This attachment signals for production of new TPO (Fig. 24.4b). TPO increases the number and rate of maturation of megakaryocytes via c-MPL receptor. Platelet levels start to rise 6 days after the start of therapy. Although TPO itself is not available for clinical use, thrombomimetic agents which bind to c-MPL are now used clinically to increase the platelet count (see p. 283).

The normal platelet count is approximately $250 \times 10^9/L$ (range $150\text{--}400 \times 10^9/L$) and the normal platelet lifespan is 10 days. This is determined by the ratio of the apoptotic BAX and anti-apoptotic BCL-2 proteins in the cell. Up to one-third of the marrow output of platelets may be trapped at any one time in the normal spleen but this rises to 90% in cases of massive splenomegaly (see Fig. 25.9).

Platelet structure

Platelets are extremely small and discoid, $3.0 \times 0.5 \mu\text{m}$ in diameter. The ultrastructure of platelets is represented in Figure 24.5. The glycoproteins of the surface coat are particularly important in the platelet reactions of adhesion and aggregation, which are the initial events leading to platelet plug formation during haemostasis. Adhesion to collagen is facilitated by glycoprotein Ia (GPIa) (Fig. 24.6). Glycoproteins Ib (defective in Bernard–Soulier syndrome) and IIb/IIIa (also called αIIb and β_3 , defective in Glanzmann's thrombasthenia) are important in the attachment of platelets to von Willebrand factor (VWF) and hence to vascular

subendothelium (Fig. 24.6). The binding site for IIb/IIIa is also the receptor for fibrinogen which, like VWF, is important in platelet–platelet aggregation.

The plasma membrane invaginates into the platelet interior to form an open membrane (canalicular) system which provides a large reactive surface to which the plasma coagulation proteins may be selectively absorbed. The membrane phospholipids (previously known as platelet factor 3) are of particular importance in the conversion of coagulation factor X to Xa and prothrombin (factor II) to thrombin (factor IIa) (see Fig. 24.8).

The platelet contains three types of storage granules: dense, α and lysosomes (Fig. 24.5). The more frequent specific α granules contain clotting factors, VWF, platelet-derived growth factor (PDGF) and other proteins. Dense granules are less common and contain adenosine diphosphate (ADP), adenosine triphosphate (ATP), serotonin and calcium. Lysosomes contain hydrolytic enzymes. Platelets are also rich in signalling and cytoskeletal proteins, which support the rapid switch from quiescence to activation that follows vessel damage. During the release reaction described below, the contents of the granules are discharged into the open canalicular system.

Platelet antigens

Several platelet surface proteins have been found to be important antigens in platelet-specific autoimmunity and they have been termed human platelet antigens (HPA). In most cases, two different alleles exist, termed a or b alleles (e.g. HPA-1a). Platelets also express ABO and human leucocyte antigen (HLA) class I but not class II antigens.




Figure 24.3 Megakaryocytes: **(a)** immature form with basophilic cytoplasm; **(b)** mature form with many nuclear lobes and pronounced granulation of the cytoplasm. **(c)** Megakaryocyte in culture, stained for α -tubulin (green). Proplatelets can be seen budding from the tips of megakaryocyte cytoplasm. Source: From Pecci A. et al. (2009) *Thrombosis and Haemostasis* 109: 90–96.

Platelet function

The main function of platelets is the formation of mechanical plugs during the haemostatic response to vascular injury. In the absence of platelets, spontaneous leakage of blood through small vessels may occur. There are three major platelet functions: **adhesion**, **aggregation** and **release reactions and amplification**. The immobilization of platelets at the sites of vascular injury requires specific platelet–vessel wall (adhesion) and platelet–platelet (aggregation) interactions, both partly mediated through VWF (Fig. 24.6).

Von Willebrand factor (VWF)

VWF is involved in shear-dependent platelet adhesion to the vessel wall and to other platelets (aggregation) (Fig. 24.6). It also carries factor VIII. It is a large glycoprotein, with multimers made up on average of 2–50 dimeric subunits. VWF is synthesized both in endothelial cells and megakaryocytes, and stored in Weibel–Palade bodies and platelet α granules, respectively.

Plasma VWF is almost entirely derived from endothelial cells, with two distinct pathways of secretion. The majority is continuously secreted and a minority is stored in Weibel–Palade bodies. The stored VWF can raise the plasma levels when released under the influence of several secretagogues, such as stress, exercise, adrenaline and infusion of desmopressin (1-diamino-8-D-arginine vasopressin; DDAVP). The VWF released from Weibel–Palade bodies is in the form of large and ultra-large multimers, the most adhesive and reactive form of VWF. They are in turn cleaved in plasma to smaller multimers and monomeric VWF by the specific plasma metalloprotease, ADAMTS13 (see Fig. 25.7).

Platelet aggregation

This is characterized by cross-linking of platelets through active GPIIb/IIIa receptors with fibrinogen bridges. A resting platelet has GPIIb/IIIa receptors which do not bind fibrinogen, VWF or other ligands. Stimulation of a platelet leads to an increase in




Figure 24.4 (a) Thrombopoietin (TPO) binds to the MPL receptor on the surface of megakaryocytes and platelets and is not recycled. Plasma levels of TPO, constitutively produced by the liver, are therefore affected by platelet and megakaryocyte MPL mass, the levels being high if this mass is low, as in aplastic anaemia, and low if this mass is high, as in essential thrombocythaemia. **(b)** The regulation of platelet production through hepatic clearance of de-sialylated platelets which attach to the Ashwell–Morell hepatic receptor. This initiates a signal via JAK2 for further thrombopoietin (TPO) and hence platelet production. GAL, galactose.

GPIIb/IIIa molecules, enabling platelet cross-linking via VWF and fibrinogen bridges (Fig. 24.6).

Platelet release reaction and amplification

Primary activation by various agonists induces intracellular signalling, leading to the release of α granule contents. These have an important role in platelet aggregate formation and stabilization and, in addition, the ADP released from dense granules has a major positive feedback role in promoting platelet activation.

Thromboxane A_2 (TXA₂) is important in secondary amplification of platelet activation to form a stable platelet aggregate. It is formed *de novo* upon activation of cytosolic phospholipase A_2 (PL_{A2}) (Fig. 24.7). TXA₂ lowers platelet cyclic adenosine monophosphate (cAMP) levels and initiates the release reaction (Fig. 24.7). TXA₂ not only potentiates platelet aggregation, but also has powerful vasoconstrictive activity. The release

reaction is inhibited by substances that increase the level of platelet cAMP. One such substance is prostacyclin (PGI₂), which is synthesized by vascular endothelial cells. It is a potent inhibitor of platelet aggregation and prevents their deposition on normal vascular endothelium.

Platelet procoagulant activity

After platelet aggregation and release, the exposed membrane phospholipid (platelet factor 3) is available for two reactions in the coagulation cascade. Both phospholipid-mediated reactions are calcium-ion dependent. The first (tenase) involves factors IXa, VIIIa and X in the formation of factor Xa (Fig. 24.8). The second (prothrombinase) results in the formation of thrombin from the interaction of factors Xa, Va and prothrombin (II). The phospholipid surface forms an ideal template for the crucial concentration and orientation of these proteins.




Figure 24.5 The ultrastructure of platelets. ADP, adenosine diphosphate; PDGF, platelet-derived growth factor; PF, platelet factor; VWF, von Willebrand factor.




Figure 24.6 Platelet adhesion to the blood vessel wall and to each other (aggregation). The binding of glycoprotein (GP)Ib to von Willebrand factor leads to platelet adhesion to the subendothelium. It also exposes the GPIIb/IIIa binding sites. The GPIa site permits direct adhesion of platelets to collagen and also further exposes the GPIIb/IIIa binding site. This leads to further binding to von Willebrand factor and so results in further platelet adhesion to the subendothelium. Platelets also attach to each other, platelet aggregation. The GPIb and IIb/IIIa are involved via von Willebrand factor. Fibrinogen is also directly involved via receptors for it on GPIIb/IIIa.




Figure 24.7 The synthesis of prostacyclin and thromboxane A₂. The opposing effects of these agents are mediated by changes in the concentration of cyclic adenosine monophosphate (cAMP) in platelets via stimulation or inhibition of the enzyme adenylate cyclase. cAMP controls the concentration of free calcium ions in the platelet, which are important in the processes that cause adhesion and aggregation. High levels of cAMP lead to low free calcium ion concentrations and prevent aggregation and adhesion. ATP, adenosine triphosphate; Ca, calcium; PG, prostaglandin (G₂ and H₂).

Growth factor

PDGF found in the α granules of platelets stimulates vascular smooth muscle cells to multiply and this may hasten vascular healing following injury.

Natural inhibitors of platelet function

Nitric oxide (NO) is constitutively released from endothelial cells (Fig. 24.9) and also from macrophages and platelets. It inhibits platelet activation and promotes vasodilatation. Prostacyclin synthesized by endothelial cells also inhibits platelet function (Fig. 24.9) and causes vasodilatation by raising cyclic guanosine monophosphate (GMP) levels. An ectonucleotidase (CD39) acts as an ADPase and helps prevent platelet aggregation in the intact vessel wall.

Blood coagulation

The coagulation cascade

Blood coagulation *in vivo* involves a **biological amplification** system in which relatively few initiation substances sequentially activate by proteolysis a **cascade of circulating precursor proteins** (the coagulation factor enzymes) which culminates

in the **generation of thrombin**; this, in turn, converts soluble plasma fibrinogen into fibrin (Fig. 24.8). **Fibrin** enmeshes the platelet aggregates at the sites of vascular injury and converts the unstable primary platelet plugs to firm, definitive and stable haemostatic plugs. A list of the coagulation factors is given in Table 24.1. The operation of this enzyme cascade requires local concentration of circulating coagulation factors at the site of injury.

Surface-mediated reactions occur on exposed collagen, platelet phospholipid and tissue factor. With the exception of fibrinogen, which is the fibrin clot subunit, the coagulation factors are either enzyme precursors or cofactors (Table 24.1). All the enzymes, except factor XIII, are serine proteases (i.e. their ability to hydrolyse peptide bonds depends upon the amino acid serine at their active centre). The scale of amplification achieved in this system is dramatic, (e.g. 1 mol of activated factor XI through sequential activation of factors IX, X and prothrombin may generate up to 2×10^8 mol of fibrin).

Coagulation *in vivo*

The generation of thrombin *in vivo* is a complex network of amplification and negative feedback loops to ensure a localized and limited production. The generation of thrombin is

Table 24.1 The coagulation factors.

Factor number	Descriptive name	Active form
I	Fibrinogen	Fibrin subunit
II	Prothrombin	Serine protease
III	Tissue factor	Receptor/cofactor*
V	Labile factor	Cofactor
VII	Proconvertin	Serine protease
VIII	Antihaemophilic factor	Cofactor
IX	Christmas factor	Serine protease
X	Stuart–Prower factor	Serine protease
XI	Plasma thromboplastin antecedent	Serine protease
XII	Hageman (contact) factor	Serine protease
XIII	Fibrin-stabilizing factor	Transglutaminase
	Prekallikrein (Fletcher factor)	Serine protease
	HMWK (Fitzgerald factor)	Cofactor*

HMWK, high molecular weight kininogen.
* Active without proteolytic modification.

dependent on three enzyme complexes, each consisting of protease, cofactor, phospholipids (PL) and calcium. They are: (i) extrinsic Xase (VIIa, TF, PL, Ca^{2+}) generating FXa; (ii) intrinsic Xase (IXa, VIIIa, PL, Ca^{2+}) also generating FXa; and (iii) prothrombinase complex (Xa, Va, PL, Ca^{2+}) generating thrombin. The generation of thrombin following vascular injury occurs in two waves of very different magnitude. During the initial phase small amounts of thrombin are generated (picomolar concentrations). This thrombin leads to a second million times larger burst of thrombin production (Figs 24.8 and 24.9).

Initiation

Coagulation is initiated after vascular injury by the interaction of the membrane bound tissue factor (TF), exposed and activated by vascular injury, with plasma factor VII.

It is expressed on fibroblasts and small muscles of the vessel wall and in the blood stream on microparticles, and on other non-vascular cells. The factor VIIa–tissue factor (extrinsic factor Xase) complex activates both factor IX and factor X. The factor Xa, in the absence of its cofactor, forms small amounts of thrombin from prothrombin. This is insufficient to initiate significant fibrin polymerization. Amplification is needed.




Figure 24.8 The pathway of blood coagulation in vivo initiated by tissue factor (TF) on the cell surface. When plasma comes into contact with TF, factor VII binds to TF. The complex of TF and activated VII (VIIa) activates X and IX. The generation of thrombin from prothrombin by the action of Xa–Va complex leads to fibrin formation. Importantly, the small amounts of thrombin generated serve to greatly amplify coagulation. Thrombin activates XI, VIII, V and XIII, and also cleaves VIII from its carrier von Willebrand factor (VWF), massively increasing the formation of VIIIa–IXa and hence of Xa–Va.




Figure 24.9 The actions of thrombin in the coagulation pathway. It also activates platelets and inflammation.

Amplification

The initiation pathway or extrinsic Xase is rapidly inactivated by tissue factor pathway inhibitor (TFPI). Thrombin generation is now dependent on the traditional intrinsic pathway (Figs 24.8 and 24.10). In this factor VIII and V are converted

to VIIIa and Va by the small amounts of thrombin generated during initiation. In this amplification phase the intrinsic Xase, formed by IXa and VIIIa on phospholipid surface in the presence of Ca^{2+} , activates sufficient Xa, which then, in combination with Va, PL and Ca^{2+} , forms the prothrombinase complex and results in the explosive generation of thrombin which acts on fibrinogen to form the fibrin clot.

Factor XI does not seem to have a role in the physiological initiation of coagulation. It has a supplementary role in the activation of factor IX (see above) and may be important at major sites of trauma or at operations, in which situations factor XI deficient individuals tend to bleed excessively. It is also involved in the contact pathway (Fig 24.10).

Thrombin hydrolyses fibrinogen, releasing fibrinopeptides A and B to form fibrin monomers (Fig. 24.11). Fibrin monomers link spontaneously by hydrogen bonds to form a loose insoluble fibrin polymer. Factor XIII is also activated by thrombin and stabilizes the fibrin polymers with the formation of covalent bond cross-links.

Fibrinogen consists of two identical subunits, each containing three dissimilar polypeptide chains (α , β and γ) which are linked




Figure 24.10 The intrinsic (contact), extrinsic and common pathways of blood coagulation. The APPT tests the intrinsic and common pathways, the PT the extrinsic and common pathways and the TT tests for thrombin inhibitors and deficiency or abnormality of fibrinogen. HMWK, high molecular weight kininogen; APPT, PT, TT, see Table 24.3.




Figure 24.11 The formation and stabilization of fibrin.

by disulphide bonds. After cleavage by thrombin of small fibrinopeptides A and B from the α and β chains, fibrin monomer consists of three paired α , β and γ chains which rapidly polymerise.

Some of the properties of the coagulation factors are listed in Table 24.2. The activity of factors II, VII, IX and X is dependent upon vitamin K, which is responsible for carboxylation of a number of terminal glutamic acid residues on each of these molecules (see Fig. 26.7).

Although factor VIII and V cofactors are not protease enzymes, they circulate in a precursor form that requires limited cleavage by thrombin for expression of full cofactor activity (Fig. 24.9).

Endothelial cells

The endothelial cell has an active role in the maintenance of vascular integrity. This cell provides the basement membrane that normally separates collagen, elastin and fibronectin of the subendothelial connective tissue from the circulating blood (Fig. 24.12). Loss of or damage to the endothelium results in both haemorrhage and activation of the haemostatic mechanism. The endothelial cell also has a potent inhibitory influence on the haemostatic response, largely through the synthesis of prostaglandin, NO, and the ectonucleotidase CD39, which have vasodilatory properties and inhibit platelet aggregation.

Synthesis of tissue factor that initiates haemostasis only occurs in endothelial cells following activation, when its natural inhibitor, TFPI, is also synthesized. Endothelial synthesis of prostacyclin, VWF, factor VIII, plasminogen activator, antithrombin and thrombomodulin, the surface protein responsible for activation of protein C, provides agents that are vital to both platelet reactions and blood coagulation (Fig. 24.12).

Haemostatic response (Fig. 24.1)

Vasoconstriction

Immediate vasoconstriction of the injured vessel and reflex constriction of adjacent small arteries and arterioles is responsible for an initial slowing of blood flow to the area of injury. When there is widespread damage, this vascular reaction prevents exsanguination. The reduced blood flow allows contact activation of platelets and coagulation factors. The vasoactive

Table 24.2 The coagulation factors.

Factor	Plasma half-life (h)	Plasma concentration (mg/L)	Comments
II	65	100	Prothrombin group: vitamin K needed for synthesis; require Ca^{2+} for activation
VII	5	0.5	
IX	25	5	
X	40	10	
I	90	3000	Thrombin interacts with them; increase in inflammation, pregnancy, oral contraceptives
V	15	10	
VIII	10	0.1	
XI	45	5	
XIII	200	30	




Figure 24.12 The endothelial cell forms a barrier between platelets and plasma clotting factors and the subendothelial connective tissues. Endothelial cells produce substances that can initiate coagulation, cause vasodilatation, inhibit platelet aggregation or haemostasis, or activate fibrinolysis.

amines and TXA₂ liberated from platelets (Fig. 24.6), and the fibrinopeptides liberated during fibrin formation (Fig. 24.11), also have vasoconstrictive activity.

Platelet reactions and primary haemostatic plug formation

Following a break in the endothelial lining, there is an initial adherence of platelets (via GP1a and GP1b receptors) to exposed connective tissue, mediated (GP1b) by VWF. Under conditions of high shear stress (e.g. arterioles) the exposed subendothelial matrix is initially coated with VWF. Collagen exposure and thrombin generated through activation of tissue factor produced at the site of injury cause the adherent platelets to release their granule contents and also activate platelet prostaglandin synthesis, leading to the formation of TXA₂. Released ADP causes platelets to swell and aggregate.

Platelet rolling in the direction of blood flow over exposed VWF with activation of GPIIb/IIIa receptors results in firmer binding. Additional platelets from the circulating blood are drawn to the area of injury. This continuing platelet aggregation promotes the growth of the haemostatic plug, which soon covers the exposed connective tissue. The unstable primary haemostatic plug produced by these platelet reactions in the first minute or so following injury is usually sufficient to provide temporary control of bleeding. The highly localized enhancement of platelet activation by ADP and TXA₂ results in a platelet mass large enough to plug the area of endothelial injury.

Stabilization of the platelet plug by fibrin

Definitive haemostasis is achieved when fibrin, formed by blood coagulation, is added to the platelet mass and by platelet-induced clot retraction/compaction.

Following vascular injury, the formation of extrinsic Xase (VIIa, TF, PL and Ca²⁺) initiates the coagulation cascade. Platelet aggregation and release reactions accelerate the coagulation process by providing abundant membrane phospholipid. The much larger amounts of thrombin generated by the secondary intrinsic Xase at the injury site converts soluble plasma fibrinogen into fibrin, potentiates platelet aggregation and secretion and also activates factor XI and XIII and cofactors V and VIII. The fibrin component of the haemostatic plug increases as the fused platelets completely degranulate and autolyse, and after a few hours the entire haemostatic plug is transformed into a solid mass of cross-linked fibrin (Fig. 24.11). Clot retraction occurs, which is mediated by GPIIb/IIIa receptors which link the cytoplasmic actin filaments to surface-bound fibrin polymers. Nevertheless, because of incorporation of plasminogen and TPA (see p. 275), this plug begins to autodigest during the same time frame.

Physiological limitation of blood coagulation

Unchecked, blood coagulation would lead to dangerous occlusion of blood vessels (thrombosis) if the protective mechanisms of coagulation factor inhibitors, blood flow and fibrinolysis were not in operation.

Coagulation factor inhibitors

It is important that the effect of thrombin is limited to the site of injury. The first inhibitor to act is TFPI, which is synthesized in endothelial cells and is present in plasma and platelets and accumulates at the site of injury caused by local platelet activation. TFPI inhibits Xa and VIIa and tissue factor to limit the main *in vivo* pathway. There is also direct inactivation of thrombin and other serine protease factors by other circulating inhibitors, of which antithrombin is the most potent. It inactivates serine proteases (see Fig. 28.3). Heparin potentiates its action markedly. Another protein, heparin cofactor II, also inhibits thrombin. α_2 -Macroglobulins, α_2 -antiplasmin, C₁ esterase inhibitor and α_1 -antitrypsin also exert inhibitory effects on circulating serine proteases.

Protein C and protein S

These are inhibitors of coagulation cofactors V and VIII. Thrombin binds to an endothelial cell surface receptor, thrombomodulin. The resulting complex activates the vitamin K-dependent serine protease, protein C, which is able to destroy activated factors V and VIII, thus preventing further thrombin generation. The action of protein C is enhanced by another vitamin K-dependent protein, S, which binds protein C to the platelet surface (Fig. 24.13). An endothelial protein C receptor localizes protein C to the endothelial surface, promoting protein C activation by the thrombin–thrombomodulin complex. In addition, activated protein C enhances fibrinolysis (Fig. 24.13).

As with other serine proteases, activated protein C is subject to inactivation by serum protease inactivators (serpins), e.g. antithrombin.

Blood flow

At the periphery of a damaged area of tissue, blood flow rapidly achieves dilution and dispersal of activated factors before fibrin formation has occurred. Activated factors are destroyed by liver parenchymal cells and particulate matter is removed by liver Kupffer cells and other reticuloendothelial cells.

Fibrinolysis

Fibrinolysis (like coagulation) is a normal haemostatic response to vascular injury. Plasminogen, a proenzyme in blood and tissue fluid, is converted to the serine protease plasmin by activators either from the vessel wall (intrinsic activation) or from the tissues (extrinsic activation) (Fig. 24.14). The most important route follows the release of **tissue plasminogen activator (TPA)** from endothelial cells. TPA is a serine protease that binds to fibrin. This enhances its capacity to convert thrombus-bound plasminogen into plasmin. This fibrin dependence of TPA action strongly localizes plasmin generation by TPA to the fibrin clot. Release of TPA occurs after such stimuli as trauma, exercise or emotional stress. Activated protein C stimulates fibrinolysis by destroying plasma inhibitors of TPA (Fig. 24.13).

Plasmin generation at the site of injury limits the extent of the evolving thrombus. The split products of fibrinolysis are also competitive inhibitors of thrombin and fibrin polymerization. Normally, α_2 -antiplasmin inhibits any local free plasmin.

Fibrinolytic agents are widely used in clinical practice (see p. 318). Recombinant TPA, the bacterial agent, streptokinase, and urokinase, initially isolated from human urine, are available.




Figure 24.13 Activation and action of protein C by thrombin which has bound to thrombomodulin on the endothelial cell surface. Protein S is a cofactor that facilitates binding of activated protein C to the platelet surface. The inactivation of factors Va and VIIIa results in the inhibition of blood coagulation. The inactivation of tissue plasminogen activator inhibitor (TPAI) enhances fibrinolysis. EPCR, endothelial protein C receptor.




Figure 24.14 The fibrinolytic system. Thrombin activates a fibrinolysis inhibitor. tPA, tissue plasminogen activator.

Plasmin is capable of digesting fibrinogen, fibrin, factors V and VIII and many other proteins. Cleavage of peptide bonds in fibrin and fibrinogen produces a variety of split (degradation) products (Fig. 24.11). Large amounts of the smallest fragments can be detected in the plasma of patients with disseminated intravascular coagulation (see p. 297).

Inactivation of plasmin

Tissue plasminogen activator is inactivated by plasminogen activator inhibitor (PAI). Circulating plasmin is inactivated by the potent inhibitors α_2 -antiplasmin and α_2 -macroglobulin.

Tests of haemostatic function

Defective haemostasis with abnormal bleeding may result from:

- 1 A vascular disorder;
- 2 Thrombocytopenia or a disorder of platelet function; or
- 3 Defective blood coagulation.

A number of simple tests are employed to assess the platelet, vessel wall and coagulation components of haemostasis.

Blood count and blood film examination

As thrombocytopenia is a common cause of abnormal bleeding, patients with suspected bleeding disorders should initially have a blood count including platelet count and blood film examination. In addition to establishing the presence of thrombocytopenia, the cause may be obvious (e.g. acute leukaemia). Modern counters measure platelet volume but this parameter is not used routinely in clinical practice for diagnosis of platelet disorders. The absolute count of immature platelets (which contain RNA) correlates with increased platelet production, e.g. after haemorrhage, but again is not routinely measured or used clinically.

Screening tests of blood coagulation

Screening tests provide an assessment of the 'extrinsic' and 'intrinsic' systems of blood coagulation and also the central conversion of fibrinogen to fibrin (Fig 24.10, Table 24.3).

Table 24.3 Screening tests used in the diagnosis of coagulation disorders (see also Fig. 24.10)

Screening tests	Abnormalities indicated by prolongation	Most common cause of coagulation disorder
Thrombin time (TT)	Deficiency or abnormality of fibrinogen or inhibition of thrombin by heparin or FDPs	DIC Heparin therapy
Prothrombin time (PT)	Deficiency or inhibition of one or more of the following coagulation factors: VII, X, V, II, fibrinogen	Liver disease Warfarin therapy DIC
Activated partial thromboplastin time (APTT or PTTK)	Deficiency or inhibition of one or more of the following coagulation factors: XII, XI, IX (Christmas disease), VIII (haemophilia), X, V, II, fibrinogen	Haemophilia, Christmas disease (+ conditions above)
Fibrinogen quantitation	Fibrinogen deficiency	DIC, liver disease

DIC, disseminated intravascular coagulation; FDPs, fibrin degradation products.
N.B. Platelet count and the tests of platelet function are also used in screening patients with a bleeding disorder (p. XXX).

The **prothrombin time (PT)** measures factors VII, X, V, prothrombin and fibrinogen. Tissue thromboplastin (a brain extract) or [synthetic] tissue factor with lipids and calcium is added to citrated plasma. The normal time for clotting is 10–14 s. It may be expressed as the international normalized ratio (INR) (p. 315).

The **activated partial thromboplastin time (APTT)** measures factors VIII, IX, XI and XII in addition to factors X, V, prothrombin and fibrinogen. Three substances – phospholipid, a surface activator (e.g. kaolin) and calcium – are added to citrated plasma. The normal time for clotting is approximately 30–40 s.

Prolonged clotting times in the PT and APTT because of factor deficiency are corrected by the addition of normal plasma to the test plasma (50:50 mix). If there is no correction or incomplete correction with normal plasma, the presence of an inhibitor of coagulation is suspected.

The **thrombin (clotting) time (TT)** is sensitive to a deficiency of fibrinogen or inhibition of thrombin. Diluted bovine thrombin is added to citrated plasma at a concentration giving a clotting time of 14–16 s with normal subjects.

Specific assays of coagulation factors

Most factor assays are based on an APTT or PT in which all factors except the one to be measured are present in the substrate plasma. This usually requires a supply of plasma from patients with hereditary deficiency of the factor in question or artificially produced factor-deficient plasma. The corrective effect of the unknown plasma on the prolonged clotting time of the deficient substrate plasma is then compared with the corrective effect of normal plasma. Results are expressed as a percentage of normal activity.

A number of chemical, chromogenic and immunological methods are available for quantification of other proteins such as fibrinogen, VWF, factor Xa and factor VIII.

Bleeding time

The bleeding time is not a reliable assessment of platelet function as it is insensitive and has poor reproducibility. It was used to identify abnormal platelet function, including the diagnosis of VWF deficiency, but is no longer used in routine clinical practice. The test involved the application of pressure to the upper arm with a blood pressure cuff, after which small incisions are made in the flexor surface forearm skin. Bleeding stops normally in 3–8 minutes. It has been replaced by specific platelet aggregation tests, platelet adhesion assays and the platelet function analysis-100 (PFA-100) test (see below). The bleeding time is prolonged in thrombocytopenia but is normal in vascular causes of abnormal bleeding.

Tests of platelet function

Conventional **platelet aggregometry** measures the fall in light absorbance in platelet-rich plasma as platelets aggregate. Initial (primary) aggregation is caused by an external agent; the secondary

response is caused by aggregating agents released from the platelets themselves. The five external aggregating agents most commonly used are ADP, collagen, ristocetin, arachidonic acid and adrenaline. The pattern of response to each agent helps to make the diagnosis (see Fig. 25.10). Flow cytometry is now increasingly used in routine practice to identify platelet glycoprotein defects.

In the **PFA-100 test**, citrated blood is aspirated through a capillary tube on to a membrane coated with collagen/ADP or collagen/adrenaline. Blood flow is maintained. Platelets begin to adhere and aggregate, primarily via VWF interactions with GPIb and GPIIb/IIIa, resulting in occlusion of the aperture. The PFA-100 is prolonged in VW disease and with other defects of platelet function. The analysis may give false-negative results with relatively common platelet defects. Full platelet aggregation tests and VWF screening may be required to exclude abnormal platelet function, even if the PFA-100 test is normal.


Tests of fibrinolysis

Testing for hyperfibrinolysis by traditional tests, such as the euglobulin clot lysis times, is rarely performed. A clinically significant hyperfibrinolytic state, e.g. during liver transplantation, can be detected by viscoelastic measurement of clot stability using **thromboelastography (TEG) or thromboelastometry (ROTEM)** (see Fig. 26.10). Treatment with tranexamic acid reduces hyperfibrinolysis and reduces bleeding.

D-dimer is a measurement of fibrin degradation products and is an indication of sequential thrombin and then plasmin activity. The test can be performed on citrated plasma samples along with simple coagulation tests. There are many causes of a high D-dimer, including infection, cancer and pregnancy, as well as venous thromboembolism. Plasma levels are very high in patients with disseminated intravascular coagulation (DIC).

- Normal haemostasis requires vasoconstriction, platelet aggregation and blood coagulation. The intact endothelial cell separates collagen and other subendothelial connective tissues that would stimulate platelet aggregation from circulating blood. The endothelial cells also produce prostacyclin, nitric oxide and an ectonucleotidase, which inhibit platelet aggregation.
- Platelets are produced from megakaryocytes in the bone marrow stimulated by thrombopoietin. They have surface glycoproteins which facilitate direct adherence to subendothelial tissues and also, via von Willebrand factor, to collagen, to other platelets (aggregation) and to fibrinogen. Platelets contain different types of storage granules which are released after platelet activation.
- Blood coagulation in vivo in response to vascular injury commences with tissue factor binding to clotting factor VII and this initiates a cascade which results in thrombin generation. Thrombin then activates cofactors VIII and V and factor XI which greatly amplify the coagulation pathway resulting in a fibrin clot.
- Coagulation factor inhibitors include antithrombin, protein C and protein S.
- Dissolution of fibrin clots (fibrinolysis) occurs by activation of plasminogen to plasmin.
- Tests of haemostatic function include the thrombin time (TT), prothrombin time (PT), activated partial thromboplastin time (APTT) as well as individual coagulation factor assays and assay of von Willebrand factor. Tests of platelet function include the PFA-100 and platelet aggregation tests.




A vertical strip on the left side of the page showing a microscopic view of blood cells. It features large, bright red disc-shaped cells (erythrocytes) and smaller, yellowish granular cells (leucocytes) against a dark blue background.

CHAPTER 25

Bleeding disorders caused by vascular and platelet abnormalities

Key topics

■ Vascular bleeding disorders	279
■ Thrombocytopenia	281
■ Autoimmune (idiopathic) thrombocytopenic purpura	282
■ Thrombotic thrombocytopenic purpura	285
■ Disorders of platelet function	287
■ Diagnosis of platelet disorders	288
■ Platelet transfusions	289



Abnormal bleeding

This may result from:

- 1 Vascular disorders;
- 2 Thrombocytopenia;
- 3 Defective platelet function; or
- 4 Defective coagulation.

The pattern of bleeding is relatively predictable depending on the aetiology. Vascular and platelet disorders tend to be associated with bleeding from mucous membranes and into the skin, whereas in coagulation disorders the bleeding is often into joints or soft tissue (Table 25.1). Table 25.2 shows the WHO grades of bleeding.

The first three categories are discussed in this chapter and the disorders of blood coagulation follow in Chapter 26.

Vascular bleeding disorders

The vascular disorders are a heterogeneous group of conditions characterized by easy bruising and spontaneous bleeding from the small vessels. The underlying abnormality is

Table 25.1 Clinical differences between diseases of platelets/vessel wall or of coagulation factors.

	Platelets/vessel wall diseases	Coagulation diseases
Mucosal bleeding	Common	Rare
Petechiae	Common	Rare
Deep haematomas	Rare	Characteristic
Bleeding from skin cuts	Persistent	Minimal
Sex of patient	Equal	>80% male

Table 25.2 WHO bleeding grades.

Grade 0	None
Grade 1	Petechiae, ecchymoses, occult blood loss, mild spotting
Grade 2	Gross bleeding, i.e. epistaxis, haematuria, haematemesis not requiring transfusion
Grade 3	Haemorrhage requiring transfusion
Grade 4	Haemorrhage with haemodynamic compromise, retinal haemorrhage with visual impairment, CNS haemorrhage, fatal at any location
CNS, central nervous system	

either in the vessels themselves or in the perivascular connective tissues. Most cases of bleeding caused by vascular defects alone are not severe. Frequently, the bleeding is mainly in the skin causing petechiae, ecchymoses or both. In some disorders there is also bleeding from mucous membranes. In these conditions the standard screening tests are normal. The bleeding time is usually normal and the other tests of haemostasis are also normal. Vascular defects may be inherited or acquired.

Inherited vascular disorders

Hereditary haemorrhagic telangiectasia

This uncommon disease is transmitted as an autosomal dominant trait. Various genetic defects underlie the disease such as mutations of the endothelial protein, endoglin. There are dilated microvascular swellings which appear during childhood and become more numerous in adult life. These telangiectasia develop in the skin, mucous membranes (Fig. 25.1a) and internal organs. Pulmonary, hepatic, splenic and cerebral arteriovenous shunts are seen in a minority of cases and may need local treatment. Recurrent epistaxes are frequent and recurrent gastrointestinal tract haemorrhage may cause chronic iron deficiency anaemia. Treatment is with embolization, laser treatment, oestrogens, tranexamic acid and iron supplementation. Thalidomide, lenalinomide and bevcizumab (anti-vascular endothelial growth factor) have been used to try to reduce gastrointestinal bleeding in severe cases.

Connective tissue disorders

In the Ehlers–Danlos syndromes there are hereditary collagen abnormalities with purpura resulting from defective platelet adhesion, hyperextensibility of joints and hyperelastic friable skin. Pseudoxanthoma elasticum is associated with arterial haemorrhage and thrombosis. Patients may present with superficial bruising and purpura following minor trauma or after the application of a tourniquet. Bleeding and poor wound healing after surgery may be a problem.

Giant cavernous haemangioma

These congenital malformations occasionally cause chronic activation of coagulation leading to laboratory features of disseminated intravascular coagulation (DIC) and in some cases thrombocytopenia.

Acquired vascular defects

- 1 **Simple easy bruising** is a common benign disorder which occurs in otherwise healthy women, especially those of child-bearing age.
- 2 **Senile purpura** caused by atrophy of the supporting tissues of cutaneous blood vessels is seen mainly on dorsal aspects of the forearms and hands (Fig. 25.1b).



Figure 25.1 (a) Hereditary haemorrhagic telangiectasia: the characteristic small vascular lesions are obvious on the lips and tongue. (b) Senile purpura. (c) Characteristic perifollicular petechiae in vitamin C deficiency (scurvy).

- 3 Purpura associated with infections, mainly of bacterial, viral or rickettsial origin, may cause purpura from vascular damage by the organism, from DIC or as a result of immune complex formation (e.g. measles, dengue fever or meningococcal septicaemia).
- 4 The **Henoch–Schönlein syndrome** is usually seen in children and often follows an acute upper respiratory tract infection. It is an IgA-mediated vasculitis. The characteristic purpuric rash accompanied by localized oedema and itching is usually most prominent on the buttocks and extensor surfaces of the lower legs and elbows (Fig. 25.2). Painful joint swelling, haematuria and abdominal pain may also occur. It is usually a self-limiting condition but occasional patients develop renal failure.
- 5 **Scurvy.** In vitamin C deficiency defective collagen may cause perifollicular petechiae, bruising and mucosal haemorrhage (Fig. 25.1c).
- 6 **Steroid purpura.** The purpura, which is associated with long-term steroid therapy or Cushing's syndrome, is caused by defective vascular supportive tissue.

Tranexamic acid and aminocaproic acid are useful antifibrinolytic drugs that may reduce bleeding resulting from vascular disorders or thrombocytopenia, but are relatively

contraindicated in the presence of haematuria because they might lead to clots obstructing the renal tract.




Figure 25.2 Henoch–Schönlein purpura: (a) unusually severe purpura on legs with bulla formation in a 6-year-old child; and (b) early urticarial lesions.

Thrombocytopenia

Abnormal bleeding associated with thrombocytopenia or abnormal platelet function is characterized by spontaneous skin purpura (Fig. 25.3) and mucosal haemorrhage and prolonged bleeding after trauma (Table 25.1). The main causes of thrombocytopenia are listed in Tables 25.3 and 25.4.

Failure of platelet production

This is the most common cause of thrombocytopenia and is usually part of a generalized bone marrow failure (Table 25.3). Selective megakaryocyte depression may result from drug toxicity or viral infection. Rarely it is congenital as a result of mutation of the *c-MPL* thrombopoietin receptor or of the *RBM8A* gene in association with absent radii. It is also seen in the May–Hegglin anomaly with large inclusions




Figure 25.3 (a) Typical purpura; and (b) massive subcutaneous haemorrhage in a patient with drug-induced thrombocytopenia.

Table 25.3 Causes of thrombocytopenia.

Failure of platelet production

Selective megakaryocyte depression
rare congenital defects (see text)
drugs, chemicals, viral infections
Part of general bone marrow failure
cytotoxic drugs
radiotherapy
aplastic anaemia
leukaemia
myelodysplastic syndromes
myelofibrosis
marrow infiltration (e.g. carcinoma, lymphoma, Gaucher's disease)
multiple myeloma
megaloblastic anaemia
HIV infection

Increased consumption of platelets

Immune
autoimmune
idiopathic
associated with systemic lupus erythematosus, chronic lymphocytic leukaemia or lymphoma;
infections: *Helicobacter pylori*, HIV, other viruses, malaria
drug-induced, e.g. heparin
post-transfusional purpura
feto-maternal alloimmune thrombocytopenia
Disseminated intravascular coagulation
Thrombotic thrombocytopenic purpura

Abnormal distribution of platelets

Splenomegaly (e.g. liver disease)

Dilutional loss

Massive transfusion of stored blood to bleeding patients
HIV, human immunodeficiency virus.

in granulocytes, or in Wiskott–Aldrich syndrome (WAS) with eczema and immune deficiency (see Chapter 8). WAS is caused by mutation of the *WASP* gene, the protein being a regulator of signalling in haemopoietic cells. Diagnosis of these causes of thrombocytopenia is made from the clinical history, peripheral blood count, the blood film and bone marrow examination.

Table 25.4 Thrombocytopenia as a result of drugs or toxins.

Bone marrow suppression

Predictable (dose-related)
ionizing radiation, cytotoxic drugs, ethanol
Occasional
chloramphenicol, co-trimoxazole, idoxuridine, penicillamine, organic arsenicals, benzene, etc.

Immune mechanisms (proven or probable)

Analgesics, anti-inflammatory drugs
gold salts
Antimicrobials
penicillins, rifamycin, sulphonamides, trimethoprim, para-aminosalicylate
Sedatives, anticonvulsants
diazepam, sodium valproate, carbamazepine
Diuretics
acetazolamide, chlorothiazides, furosemide
Antidiabetics
chlorpropamide, tolbutamide
Others
digitoxin, heparin, methyl dopa, oxyprenolol, quinine, quinidine

Increased destruction of platelets

Autoimmune (idiopathic) thrombocytopenic purpura

Autoimmune (idiopathic) thrombocytopenic purpura (ITP) may be divided into chronic and acute forms.

Chronic idiopathic thrombocytopenic purpura (ITP)

This is a relatively common disorder. The highest incidence has been considered to be in women aged 15–50 years, although some reports suggest an increasing incidence with age. It is the most common cause of thrombocytopenia without anaemia or neutropenia. It is usually idiopathic but may be seen in association with other diseases such as systemic lupus erythematosus (SLE), human immunodeficiency virus (HIV) infection, viral hepatitis, *Helicobacter pylori* infection, chronic lymphocytic leukaemia (CLL), Hodgkin lymphoma or autoimmune haemolytic anaemia (Table 25.3).

Pathogenesis

Platelet autoantibodies, usually IgG, result in the premature removal of platelets from the circulation by macrophages of the reticuloendothelial system, especially the spleen (Fig. 25.4). In many cases, the antibody is directed against the glycoprotein (GP) IIb/IIIa or Ib complex. The normal lifespan of a platelet is 10 days but in ITP this is reduced to a few hours. Total megakaryocyte mass and platelet turnover are increased in parallel to approximately five times normal.

Clinical features

The onset is often insidious with petechial haemorrhage, easy bruising and, in women, menorrhagia. Mucosal bleeding (e.g. epistaxes or gum bleeding) occurs in severe cases but fortunately intracranial haemorrhage is rare. The severity of bleeding in ITP is usually less than that seen in patients with comparable degrees of thrombocytopenia from bone marrow failure; this is attributed to the circulation of predominantly young,




Figure 25.4 The pathogenesis of thrombocytopenia in autoimmune thrombocytopenic purpura. Platelets coated by antibodies are phagocytosed by macrophages. The actions of thrombopoietin (TPO) and thrombopoietin receptor agonists (TPO-RA) (thrombomimetics) are shown. These are orally active or given by injection and act to increase platelet production.

functionally superior platelets in ITP. Chronic ITP tends to relapse and remit spontaneously so the course may be difficult to predict. Many asymptomatic cases are discovered by a routine blood count. The spleen is not palpable unless there is an associated disease causing splenomegaly.

Diagnosis

- 1 The platelet count is usually $10\text{--}100 \times 10^9/\text{L}$. The haemoglobin concentration and white cell count are typically normal unless there is iron deficiency anaemia because of blood loss.
- 2 The blood film shows reduced numbers of platelets, those present often being large. There are no morphological abnormalities in the other cell lines.
- 3 The bone marrow shows normal or increased numbers of megakaryocytes.
- 4 Sensitive tests are able to demonstrate specific anti-glycoprotein GPIIb/IIIa or GPIb antibodies on the platelet surface or in the serum in most patients. Platelet-associated IgG assays are less specific. These tests are not usually used in clinical practice.

Treatment

As this is a chronic disease the aim of treatment should be to maintain a platelet count above the level at which spontaneous bruising or bleeding occurs with the minimum of intervention. In general, a platelet count above $20 \times 10^9/\text{L}$ without symptoms does not require treatment.

- 1 **Corticosteroids** Eighty per cent of patients remit on high-dose corticosteroid therapy. Prednisolone 1 mg/kg/day is the usual initial therapy in adults and the dosage is gradually reduced after 10–14 days. In poor responders the dosage is reduced more slowly but alternative immunosuppression or splenectomy is considered.
- 2 **High-dose intravenous immunoglobulin therapy** is able to produce a rapid rise in platelet count in the majority of patients. A regimen of 400 mg/kg/day for 5 days or 1 g/kg/day for 2 days is used. It is particularly useful in patients with life-threatening haemorrhage, in steroid-refractory ITP, during pregnancy or prior to surgery. The mechanism of action may be blockage of Fc receptors on macrophages or modification of autoantibody production.
- 3 **Monoclonal antibody** Rituximab (anti-CD20) produces responses in approximately 50%, which are often durable and it is now usually tried before splenectomy.
- 4 **Immunosuppressive drugs** (e.g. vincristine, cyclophosphamide, azathioprine, mycophenolate mofetil or ciclosporin alone or in combination) are usually reserved for those patients who do not respond sufficiently to steroids or rituximab.
- 5 **Thrombopoietin-receptor agonists** Romiplostim (subcutaneously) and eltrombopag (orally) are active non-peptide thrombopoietin-receptor agonists (thrombomimetics)

(Fig. 25.4). They stimulate thrombopoiesis (Fig. 25.5) and are indicated for patients in whom steroids are contraindicated or who are refractory to steroids. Trials of their use as initial therapy in combination with corticosteroids are in progress. Increased reticulin and fibrosis in the bone marrow may occur with prolonged treatment but are reversible on stopping the therapy.

- 6 **Splenectomy** With the increase in number of alternative drugs, splenectomy is now performed less frequently for ITP than previously. Good results occur in most patients but in those with ITP refractory to steroids, immunoglobulin or rituximab there may be little benefit. Splenectomy must be removed otherwise subsequent relapse of ITP can occur.
- 7 **Other treatments** that may elicit a remission include danazol (an androgen which may cause virilization in women) and intravenous anti-D immunoglobulin. It is often necessary to combine two drugs (e.g. danazol and an immunosuppressive agent). *Helicobacter pylori* infection should be treated as there are some reports that this may improve the platelet count, particularly in countries where the incidence of the infection is common. Hepatitis C should also be treated, if present.
- 8 **Platelet transfusions** Platelet concentrates are beneficial in patients with acute life-threatening bleeding but their benefit will only last for a few hours.
- 9 **Stem cell transplantation** has cured some severe cases.

Acute idiopathic thrombocytopenic purpura

This is most common in children. In approximately 75% of patients the episode follows vaccination or an infection such as chickenpox or infectious mononucleosis. Most cases are caused by non-specific immune complex attachments to platelets. Spontaneous remissions are usual but in 5–10% of cases the disease becomes chronic, lasting more than 6 months. Fortunately, morbidity and mortality in acute ITP is very low. The main risk is of cerebral haemorrhage, fortunately rare. Most children do not have any bleeding even with platelet counts $<10 \times 10^9/\text{L}$ but need to avoid trauma such as contact sports.

The diagnosis is one of exclusion. If the platelet count is over $30 \times 10^9/\text{L}$ no treatment is necessary unless the bleeding is severe. Indeed many doctors do not treat even with platelet counts $<10 \times 10^9/\text{L}$ if there is no haemorrhage. The main risk is of cerebral haemorrhage. Treatment is with steroids and/or intravenous immunoglobulin, especially if there is significant bleeding.

Infections

It seems likely that the thrombocytopenia associated with many viral and protozoal infections is immune-mediated. In HIV infection, reduced platelet production is also involved (see p. 328).




Figure 25.5 Response to eltrombopag in chronic immune thrombocytopenic purpura in a female aged 75 after failure of response to prednisolone. Source: Courtesy of Professor A. Newland.

Drug-induced immune thrombocytopenia

An immunological mechanism has been demonstrated as the cause of many drug-induced thrombocytopenias (Fig. 25.6). Quinine (including that in tonic water), quinidine and heparin are particularly common causes (Table 25.4).

The platelet count is often less than $10 \times 10^9/L$, and the bone marrow shows normal or increased numbers of megakaryocytes. Drug-dependent antibodies against platelets may be demonstrated in the sera of some patients. The

immediate treatment is to stop all suspected drugs but platelet concentrates should be given to patients with dangerous bleeding.

Post-transfusion purpura

This occurs as thrombocytopenia occurring approximately 10 days after a blood transfusion, and has been attributed to antibodies in the recipient developing against human platelet antigen-1a (HPA-1a) on the transfused platelets (see p. 342)




Figure 25.6 Usual type of platelet damage caused by drugs in which an antibody–drug–protein complex is deposited on the platelet surface. If complement is attached and the sequence goes to completion, the platelet may be lysed directly. Otherwise it is removed by reticuloendothelial cells because of opsonization with immunoglobulin and/or the C3 component of complement.

Thrombotic thrombocytopenic purpura (TTP) and haemolytic uraemic syndrome (HUS)

Thrombotic thrombocytopenic purpura (TTP) occurs in familial or acquired forms. There is **deficiency of the ADAMTS13 metalloprotease** which breaks down ultra-large von Willebrand factor multimers (ULVWF) (Fig. 25.7). In the familial forms more than 50 ADAMTS13 mutations have been reported whereas the acquired forms follow the development of an inhibitory IgG autoantibody, the presence of which may be stimulated by

infection, autoimmune/connective tissue disease, certain drugs, stem cell transplantation or cardiac surgery. ULVWF multimeric 'strings' secreted from Weibel–Palade bodies are anchored to the endothelial cells, and passing platelets adhere via their GPIb receptors. Increasing platelet aggregation on to the ULVWF multimeric strings has the potential to form large occlusive platelet thrombi. These strings are capable of embolizing to microvessels downstream and contributing to organ ischemia (Fig. 25.8). In the closely related haemolytic uraemic syndrome (HUS), ADAMTS13 levels are normal.






Figure 25.7 Proposed pathogenesis of thrombotic thrombocytopenic purpura (TTP). Von Willebrand factor (VWF) consists of a series of VWF multimers each of molecular weight (MW) 250 kDa which are covalently linked. **(a)** Under physiological circumstances a metalloprotease ADAMTS13 cleaves high molecular weight multimers at a Tyr-842–Met-843 bond and the resulting VWF has an MW of 500–20 000 kDa. **(b)** In non-familial TTP, an antibody develops to the metalloprotease and so blocks cleavage of VWF multimers. **(c)** In congenital forms of TTP, the protease appears to be absent. In both cases, the resultant ultra-large VWF multimers can bind platelets under high shear stress conditions and lead to platelet aggregation.



(a)



(b)

Figure 25.8 Thrombotic thrombocytopenic purpura. (a) Platelet thrombus in a small cardiac vessel with minor endothelial and inflammatory reaction. Source: Courtesy of Dr J.E. McLaughlin. (b) Peripheral blood film showing red cell fragmentation.

TTP has traditionally been described as a pentad of **thrombocytopenia, microangiopathic haemolytic anaemia, neurological abnormalities, renal failure and fever**. The microvascular thrombosis causes variable degrees of tissue ischaemia and infarction and is responsible for the microangiopathic haemolytic anaemia and thrombocytopenia (Fig. 25.8). Thrombocytopenia, schistocytes in the blood film and an impressively elevated serum lactate dehydrogenase (LDH) value are sufficient to suggest the diagnosis. The serum LDH is derived both from ischaemic or necrotic tissue cells and lysed red cells. Coagulation tests are normal in contrast to the findings in DIC (see Fig. 26.9). ADAMTS13 is absent or severely reduced in plasma.

Treatment is with plasma exchange, using fresh frozen plasma (FFP) or cryosupernatant. This removes the large molecular weight VWF multimers and the antibody and provides ADAMTS13. The platelet count and serum LDH are useful for monitoring the response to treatment. Rituximab (anti-CD20) is also effective, used in conjunction with plasma infusions or with plasma exchange, and subsequently for reducing the risk of relapse. In refractory cases and chronic relapsing cases, high-dose corticosteroids, vincristine, intravenous immunoglobulin, rituximab and immunosuppressive therapy with azathioprine or cyclophosphamide have been used. In untreated cases mortality may approach 90%. Relapses are frequent.

HUS in children has many common features but organ damage is limited to the kidneys. There is also usually diarrhoea and epileptic seizures may occur. Many cases are associated with *Escherichia coli* infection with the verotoxin 0157 strain or with other organisms, especially *Shigella*. Supportive renal dialysis and control of hypertension are the mainstays of treatment. Platelet transfusions are contraindicated in HUS and TTP. For atypical HUS, treatment is as for renal failure but eculizumab (see p. 247) may be added to inhibit complement activation.




Figure 25.9 The platelet distribution between the circulation and spleen in normal individuals (left), and in patients with moderate or massive splenomegaly (right).

Disseminated intravascular coagulation

In this disorder thrombocytopenia may result from a high rate of platelet destruction due to increased levels of consumption.

Increased splenic pooling

The major factor responsible for thrombocytopenia in splenomegaly is platelet 'pooling' by the spleen. In splenomegaly, up to 90% of platelets may be sequestered in the spleen, whereas normally this accounts for approximately one-third of the total platelet mass (Fig. 25.9). Platelet lifespan is normal and, in the absence of additional haemostatic defects, the thrombocytopenia of splenomegaly is not usually associated with bleeding.

Massive transfusion syndrome

Platelets are unstable in blood stored at 4°C and the platelet count rapidly falls in blood stored for more than 24 hours. Patients transfused with massive amounts of stored blood, such as more than 10 units over a 24-hour period, frequently show abnormal clotting and thrombocytopenia. These should be corrected by the use of platelet transfusions and FFP.

Disorders of platelet function

Disorders of platelet function are suspected in patients who show skin and mucosal haemorrhage despite a normal platelet count and normal levels of VWF. These disorders may be hereditary or acquired.

Hereditary disorders

Rare inherited disorders may produce defects at each of the different phases of the platelet reactions leading to the formation of the haemostatic platelet plug.

Thrombasthenia (Glanzmann's disease)

This autosomal recessive disorder leads to failure of primary platelet aggregation. It is caused by a variety of different mutations in the genes coding for GPIIb or IIIa (Fig. 24.6). It usually presents in the neonatal period and, characteristically, platelets fail to aggregate *in vitro* to any agonist except ristocetin.

Bernard–Soulier syndrome

In this autosomal recessive disease due to mutations in the GPIb gene, the platelets are larger than normal. There is defective binding to VWF, defective adherence to exposed sub-endothelial connective tissues and platelets do not aggregate with ristocetin. There is a variable degree of thrombocytopenia.

Storage pool diseases

A mutation in a growth factor independent gene (*GFI1B*) underlies the grey-platelet syndrome. The platelets are larger than normal, with virtual absence of alpha granules. In the more common delta-storage pool disease, there is deficiency of dense granules. Platelet-dependent haemostasis is abnormal in the much more common von Willebrand disease because of an inherited defect in VWF (see p. 295).

Acquired disorders

Antiplatelet drugs

Aspirin therapy is the most common cause of defective platelet function. It produces abnormal closure times in the platelet function analysis-100 (PFA-100) test and, although purpura may not be obvious, the defect may contribute to the associated gastrointestinal haemorrhage. The cause of the aspirin defect is inhibition of cyclo-oxygenase with impaired thromboxane A_2 synthesis (see Fig. 28.5). There is consequent impairment of the release reaction and aggregation (Fig. 25.10). After a single dose the defect lasts 10 days (i.e. the life of the platelet).

Dipyridamole inhibits platelet aggregation by blocking reuptake of adenosine and is usually used as an adjunct to aspirin. **Clopidogrel and prasugrel** inhibit binding of ADP to its platelet receptor (Fig. 28.5), shown by impaired aggregation with ADP (Fig. 25.10). They are used mainly for prevention of thrombotic events (e.g. after coronary stenting or angioplasty) in patients with a history of symptomatic atherosclerotic disease. The intravenous agents abciximab, eptifibatide and tirofiban are inhibitors of GPIIb/IIIa receptor sites and may be used in patients undergoing percutaneous coronary intervention, with unstable angina and acute coronary syndromes. There is a risk of transient thrombocytopenia with these agents.

See also Chapter 28.

Hyperglobulinaemia

Hyperglobulinaemia associated with multiple myeloma or Waldenström's disease may cause interference with platelet adherence, release and aggregation.

Myeloproliferative and myelodysplastic disorders

Intrinsic abnormalities of platelet function occur in many patients with essential thrombocythaemia, other myeloproliferative and myelodysplastic diseases and in paroxysmal nocturnal haemoglobinuria.

Uraemia

This is associated with various abnormalities of platelet function.




Figure 25.10 Defective platelet aggregation in patients on aspirin or clopidogrel therapy. With aspirin there is no secondary phase aggregation with adenosine diphosphate (ADP) and reduced responses to arachidonic acid, adrenaline and collagen. With clopidogrel the defect is mainly in ADP induced aggregation.

Diagnosis of platelet disorders

Patients with suspected platelet or blood vessel abnormalities should initially have a blood count and blood film examination (Fig. 25.11). Bone marrow examination is often needed in thrombocytopenic patients to determine whether or not there is a failure of platelet production. The marrow may also reveal one of the conditions associated with defective production

(Table 25.3). In children and young adults with isolated thrombocytopenia, the marrow test is often not performed. In the elderly, marrow examination is needed particularly to exclude myelodysplasia. In patients with thrombocytopenia, normal haemoglobin and white cell counts, a negative drug history, normal or excessive numbers of marrow megakaryocytes and no other marrow abnormality or splenomegaly, ITP is the usual diagnosis. Screening tests for DIC are also




Figure 25.11 Laboratory tests for platelet disorders. N.B. Some intrinsic platelet functional disorders are associated with thrombocytopenia (e.g. Bernard–Soulier syndrome). ADP, adenosine diphosphate; DIC, disseminated intravascular coagulation.

useful, as are tests for an underlying disease such as SLE or HIV infection.

When the blood count, including platelet count and blood film examination, are normal, PFA-100 is used to detect abnormal platelet function. In most patients with abnormal platelet function demonstrated by the PFA-100 test, the defect is acquired and associated either with systemic disease (e.g. uraemia) or with aspirin therapy. The rare hereditary defects of platelet function require more elaborate *in vitro* tests to define the specific abnormality. These include platelet aggregation studies (Fig. 25.10) and measurements of platelet nucleotide levels. If von Willebrand disease is suspected, assay of VWF and coagulation factor VIII are required (see p. 295).

Thrombomimetics

These are drugs that increase platelet production by activating the thrombopoietin receptor on megakaryocytes. Two such drugs are **romiplostim**, given subcutaneously once weekly, and **eltrombopag**, active orally and given daily. They are used in ITP (Fig. 25.5) and are in trial in other conditions (e.g. post-chemotherapy, myelodysplasia and aplastic anaemia). They may

cause disturbed liver function. Their long-term use may cause increased marrow reticulin and fibrosis which is reversible by stopping the drug.

Platelet transfusions

Transfusion of platelet concentrates is indicated in the following circumstances:

- 1 Thrombocytopenia or abnormal platelet function when bleeding or before invasive procedures and where there is no alternative therapy available (e.g. steroids, high-dose immunoglobulin or thrombomimetic agents). The platelet count should be above $50 \times 10^9/L$ before, for example, liver biopsy or lumbar puncture.
- 2 Prophylactically in patients with platelet counts of less than $5\text{--}10 \times 10^9/L$. If there is infection, potential bleeding sites or coagulopathy, the count should be kept above $20 \times 10^9/L$.


The indications for transfusion of platelet concentrates are discussed further on p. 344. These indications may change with the wider use of thrombomimetic drugs.

- Vascular bleeding disorders may be congenital, including hereditary haemorrhagic telangiectasia and the Ehlers–Danlos syndrome.
- Acquired vascular disorders include fragile capillaries in healthy women, senile purpura, purpura associated with infections, Henoch–Schönlein syndrome, scurvy and steroid therapy.
- Thrombocytopenia, if severe, also causes skin and mucous membrane bleeding. It has a wide range of causes including: (i) failure of platelet production from a congenital cause, drugs or viral infection or a general bone marrow failure; (ii) increased consumption of platelets. This may be acute or chronic autoimmune, drug-induced, caused by disseminated intravascular coagulation or thrombotic thrombocytopenic purpura.

- Chronic autoimmune thrombocytopenia is treated by immunosuppression with corticosteroids, rituximab, azathioprine, ciclosporin or by splenectomy.
- The platelet count may be raised by platelet transfusion or by the thrombomimetic drugs eltrombopag or romiplostim.
- Disorders of platelet function may be hereditary, as in von Willebrand disease, Glanzmann's thrombasthenia and Bernard–Soulier syndrome, or acquired, most frequently caused by drugs (e.g. aspirin, clopidogrel and dipyridamole) but also non-steroidal anti-inflammatory drugs.
- Platelet function analysis (PFA-100), platelet aggregation studies and VWF assays may be needed to diagnose platelet functional defects.



Now visit www.wileyessential.com/haematology to test yourself on this chapter.


A vertical strip on the left side of the page showing a microscopic view of blood cells. It features several large, red, biconcave disc-shaped red blood cells, some smaller white blood cells with visible nuclei, and a few yellowish platelets. The background is a dark blue.

CHAPTER 26

Coagulation disorders

Key topics

■ Hereditary coagulation disorders	291
■ Haemophilia A	291
■ Factor IX deficiency	295
■ Von Willebrand disease	295
■ Hereditary deficiency of other coagulation factors	296
■ Acquired coagulation disorders	296
■ Disseminated intravascular coagulation	297
■ Massive transfusion	301
■ Thromboelastography: near-patient testing	301



Hereditary coagulation disorders

Hereditary deficiencies of each of the coagulation factors have been described. Haemophilia A (factor VIII deficiency), haemophilia B (Christmas disease, factor IX deficiency) and von Willebrand disease (VWD) are the most frequent; the others are rarer.

Haemophilia A

Haemophilia A is the most common of the hereditary clotting factor deficiencies. The prevalence is of the order of 30–100 per million population. The inheritance is sex-linked (Fig. 26.1) but up to one-third of patients have no family history and these cases result from recent mutation.

Molecular genetics

The factor VIII gene is situated near the tip of the long arm of the X chromosome. It is extremely large and consists of 26 exons. The factor VIII protein includes a triplicated region $A_1A_2A_3$, a duplicated region C_1C_2 and a heavily glycosylated B domain which is removed when factor VIII is activated by thrombin. The protein is synthesized in endothelial cells.

The defect is an absence or low level of plasma factor VIII. Approximately half of the patients have missense or frameshift mutations or deletions in the factor VIII gene. In others a characteristic 'flip-tip' inversion is seen in which the factor VIII gene is broken by an inversion at the end of the X chromosome (Fig. 26.2). This mutation leads to a severe clinical form of haemophilia A.

Clinical features

Infants may develop profuse post-circumcision haemorrhage or joint and soft tissue bleeds and excessive bruising when they

start to be active. **Recurrent painful haemarthroses and muscle haematomas dominate the clinical course of severely affected patients and, if inadequately treated, lead to progressive joint deformity and disability (Figs 26.3, 26.4, 26.5 and 26.6).** Local pressure can cause entrapment neuropathy or ischaemic necrosis. Prolonged bleeding occurs after dental extractions. Spontaneous haematuria and gastrointestinal haemorrhage, sometimes with obstruction resulting from intramucosal bleeding, can also occur. The clinical severity of the disease correlates inversely with the factor VIII level (Table 26.1). Operative and post-traumatic haemorrhage are life-threatening both in severely and mildly affected patients. Although not common, spontaneous intracerebral haemorrhage occurs more frequently than in the general population and is an important cause of death in patients with severe disease.

Haemophilic pseudotumours are large encapsulated haematomas with progressive cystic swelling from repeated haemorrhage. They are best visualized by MRI (Fig. 26.5b). They may occur in fascial and muscle planes, large muscle groups and in the long bones, pelvis and cranium. The latter result from repeated subperiosteal haemorrhages with bone destruction and new bone formation.

As a result of human immunodeficiency virus (HIV) present in concentrates made from human plasma during the early 1980s, over 50% of haemophiliacs treated in the USA or Western Europe became infected with HIV. AIDS was a significant cause of death until the introduction of effective anti-viral therapy.

Many patients were infected with hepatitis C virus before testing of donors and blood products became possible. This has resulted in chronic hepatitis, cirrhosis and hepatoma. Hepatitis B transmission may also be a risk. Liver transplantation cures the haemophilia.

Laboratory findings

See Table 26.2.

The following tests are abnormal:

- 1 Activated partial thromboplastin time (APTT).
- 2 Factor VIII clotting assay.

The platelet function analysis-100 (PFA-100) and prothrombin time (PT) are normal.

Carrier detection and antenatal diagnosis

Carriers are detected with DNA probes. A known specific mutation can be identified or restriction fragment length polymorphisms within or close to the factor VIII gene allow the mutant allele to be tracked. Chorionic biopsies at 8–10 weeks' gestation provide sufficient fetal DNA for analysis. Antenatal diagnosis is also possible following the demonstration of low levels of factor VIII in fetal blood obtained at 16–20 weeks' gestation from the umbilical vein. This method is only used if DNA analysis is uninformative (1% of carriers).




Figure 26.1 A typical family tree in a family with haemophilia. Note the variable levels of factor VIII activity in carriers (*) because of random inactivation of X chromosome (Lyonization). The percentages show the degree of factor VIII activity as a percentage of normal.




Figure 26.2 The mechanism of the flip-tip inversion leading to disruption of the factor VIII gene. **(Left)** The orientation of the factor VIII gene is shown with the three copies of gene A (F8A) in this region (one within an intron 22 and two near the telomere). **(Middle)** During spermatogenesis at meiosis, the single X pairs with the Y chromosome in the homologous regions. The X chromosome is longer than the Y and there is nothing to pair with most of the long arm of X. The chromosome undergoes homologous recombination between the A genes. **(Right)** The final result is that the factor VIII gene is disrupted. cen, centromeric end; tel, telomere; the arrows indicate the direction of transcription from the A gene.




Figure 26.3 Haemophilia A: acute haemarthrosis of the right knee joint with swelling of the suprapatellar region. There is wasting of the quadriceps muscles, particularly on the left.



Figure 26.4 Haemophilia A showing severe disability. The left knee is swollen with posterior subluxation of the tibia on the femur. The ankles and feet show residual deformities of talipes equinus, with some cavus and associated toe clawing. There is generalized muscle wasting. The scar on the medial side of the left lower thigh is the site of a previously excised pseudotumour.



(a)



(b)

Figure 26.5 (a) Haemophilia A: massive haemorrhage in the area of the right buttock. (b) 15-year-old boy with sudden left hip pain and haemophilia A. Magnetic resonance imaging (MRI) axial image, T2-weighted, revealing large left spontaneous haematoma (yellow arrow) in left gluteus maximus muscle compared with normal right side (red cross). Source: Courtesy of Dr P. Wylie.

Treatment

Most patients in developed countries attend specialized haemophilia centres where there is a multidisciplinary team dedicated to their care. Bleeding episodes are treated with factor VIII replacement therapy, and spontaneous bleeding is usually controlled if the patient’s factor VIII level is raised to 30–50% of normal. Guidelines exist for the plasma level to be achieved for different types of haemorrhage. For major surgery, serious post-traumatic bleeding or when haemorrhage is occurring at a dangerous site, the factor VIII level should be elevated to 100% and then maintained above 50% when acute bleeding has stopped, until healing has occurred. On average, factor VIII infusion produces a plasma increment of 20 U/L for each unit infused/kg body weight.

Recombinant factor VIII and plasma-derived purified factor VIII preparations, which are heat and solvent-detergent




Figure 26.6 Haemophilia A: X-ray of the knee joints shows destruction and narrowing of the left joint space.

treated, are available for clinical use and have never transmitted viral infections.

1-Diamino-8-D-arginine vasopressin (DDAVP; desmopressin) provides an alternative means of increasing the plasma factor VIII level in milder haemophiliacs. Following the intravenous administration of this drug, there is a two- to four-fold rise maximum at 30–60 minutes in the patient’s own factor VIII by release from endothelial cells. DDAVP may also be taken subcutaneously or nasally: this has been used as

Table 26.1 Correlation of coagulation factor activity and disease severity in haemophilia A or B.

Coagulation factor activity (percentage of normal)	Clinical manifestations
<1	Severe disease Frequent spontaneous bleeding into joints, muscles, internal organs from early life Joint deformity and crippling if not adequately prevented or treated
1–5	Moderate disease Bleeding after minor trauma Occasional spontaneous episodes
>5	Mild disease Bleeding only after significant trauma, surgery

Table 26.2 Main clinical and laboratory findings in haemophilia A, factor IX deficiency (haemophilia B, Christmas disease) and von Willebrand disease.

	Haemophilia A	Factor IX deficiency	von Willebrand disease
Inheritance	Sex-linked	Sex-linked	Dominant (incomplete)
Main sites of haemorrhage	Muscle, joints, post-trauma or postoperative	Muscle, joints, post-trauma or postoperative	Mucous membranes, skin cuts, post-trauma or postoperative
Platelet count	Normal	Normal	Normal
PFA-100	Normal	Normal	Prolonged
Prothrombin time	Normal	Normal	Normal
Partial thromboplastin time	Prolonged	Prolonged	Prolonged or normal
Factor VIII	Low	Normal	May be moderately reduced
Factor IX	Normal	Low	Normal
VWF	Normal	Normal	Low or abnormal function (see Table 26.3)
Ristocetin-induced platelet aggregation	Normal	Normal	Impaired
VWF, von Willebrand factor.			

immediate treatment for mild haemophilia after accidental trauma or haemorrhage. DDAVP has an antidiuretic action and should be avoided in the elderly; fluid restriction is advised after its use.

Local supportive measures used in treating haemarthroses and haematomas include resting the affected part, application of ice and the prevention of further trauma.

Prophylactic treatment

The availability of factor VIII concentrates that may be stored in domestic refrigerators has dramatically altered haemophilia treatment. At the earliest suggestion of bleeding, the haemophilic child may be treated at home. This advance has reduced the occurrence of crippling haemarthroses and the need for inpatient care. Severely affected patients are now reaching adult life with little or no arthritis. After the first spontaneous joint bleed, most boys with severe haemophilia are started on prophylactic factor VIII three times a week, aiming to keep their factor VIII trough levels above 1%. This may require the placement of a vascular access device such as Port-a-Cath if venous access is difficult. A controlled trial has proven that regular prophylaxis is far superior to on-demand treatment. The half-life of factor VIII is only 8–12 hours. Phase 3 studies of long-acting derivatives of factor VIII and IX such as Fc-fusion and PEGylated proteins are being conducted with the intention of reducing the frequency of replacement therapy and maintaining haemostatic factor levels for longer periods of time.

Haemophiliacs are advised to have regular conservative dental care. Haemophiliac children and their parents often

require extensive help with social and psychological matters. With modern treatment the lifestyle of a haemophilic child can be almost normal but certain activities such as extreme contact sports are to be avoided, or undertaken with extra prophylaxis.

Gene therapy

Because it is only necessary to maintain factor levels above 1% to prevent most of the mortality and morbidity of factor VIII or IX deficiency, there is great interest in gene-based therapy. Various viral vectors (retroviral, adeno-associated) as well as non-viral vectors are being explored. Durable (more than 2 years after a single injection) increased levels of factor IX have been obtained with an adenoviral vector carrying the gene to the liver, sufficient to obviate the need for replacement therapy except for trauma or surgery. Trials of factor VIII gene therapy are also planned.

Inhibitors

One of the most serious complications of haemophilia is the development of antibodies (inhibitors) to infused factor VIII which occurs in 30–40% of severely affected patients, usually within the first 50 days of exposure. This renders the patient refractory to further replacement therapy. Immunosuppression and immune tolerance regimens have been used in an attempt to eradicate the antibody with success (at great cost) in about two-thirds of cases. Recombinant activated factor VII (VIIa) and activated prothrombin complex concentrates (FEIBA – factor VIII inhibitor bypassing activity) can be useful in the treatment of bleeding episodes.

Factor VIIa complexes with tissue factor exposed at the site of injury and produces local haemostasis. The process is independent of factor VIII or IX and is not affected by their inhibitors. Factor VIIa has a short half-life and therefore frequent doses may be needed. In the longer term, immunosuppression with cyclophosphamide, rituximab and intravenous immunoglobulin and high-dose factor VIII has also been successful.

Factor IX deficiency (haemophilia B, Christmas disease)

The inheritance and clinical features of factor IX deficiency (Christmas disease, haemophilia B) are identical to those of haemophilia A. Indeed, the two disorders can only be distinguished by specific coagulation factor assays. The incidence is one-fifth that of haemophilia A. Factor IX is coded by a gene close to the gene for factor VIII near the tip of the long arm of the X chromosome. Its synthesis is vitamin K-dependent. Carrier detection and antenatal diagnosis are performed as for haemophilia A. The principles of replacement therapy are similar to those of haemophilia A. Bleeding episodes are treated with high-purity factor IX concentrates. Because of its longer biological half-life, infusions do not have to be given as frequently as do factor VIII concentrates in haemophilia A. Recombinant factor IX is preferred, but higher doses are needed than with plasma-derived factor IX to attain the same response. Longer-acting factor IX, PEGylated or Fc fused, will entail less frequent infusions. Gene therapy is discussed above.

Laboratory findings

See Table 26.2.

The following tests are abnormal:

- 1 APTT;
- 2 Factor IX clotting assay.

As in haemophilia A, the PFA-100 and PT tests are normal.

Von Willebrand disease

In this disorder there is either a reduced level or abnormal function of von Willebrand factor (VWF) resulting from a wide variety of mainly missense mutations in different parts of the gene. VWF is produced in endothelial cells and megakaryocytes. It has two roles (see Chapter 24). It promotes platelet adhesion to subendothelium and each other at high shear rates and it is the carrier molecule for factor VIII, protecting it from premature destruction. The latter property explains the reduced factor VIII levels found in VWD. VWF has a half-life in plasma of about 16 hours.

Chronic elevation of VWF is part of the acute phase response to injury, inflammation, neoplasia or pregnancy. VWF is synthesized as a large 600-kDa protein which then forms high molecular weight multimers which are the largest molecules in blood. Three types of VWD have been described (Table 26.3). Type 2 is divided into four subtypes depending on the type of functional defect. Type 1 accounts for 65–75% and type 2 for most of the remainder.

Table 26.3 Classification of von Willebrand disease.

Type 1	Quantitative partial deficiency		
Type 2	Functional abnormality		
Type 3	Complete deficiency		
Secondary classification of type 2 VWD			
Subtype	Platelet-associated function	Factor VIII binding capacity	High MW VWF multimers
2A	Decreased	Normal	Absent
2B	Increased affinity for GPIb	Normal	Usually reduced/absent
2M	Decreased	Normal	Normal
2N	Normal	Reduced	Normal
GPIb, glycoprotein Ib; MW, molecular weight; VWD, von Willebrand disease; VWF, von Willebrand factor.			

VWD is the most common inherited bleeding disorder. Usually, the inheritance is autosomal dominant. The severity of the bleeding is highly variable, depending on mutation type and epistatic genetic effect, such as ABO blood group. Women are more badly affected than men at a given VWF level. Typically, there is mucous membrane bleeding (e.g. epistaxes, menorrhagia), excessive blood loss from superficial cuts and abrasions, and operative and post-traumatic haemorrhage. The severity is variable in the different types. Haemarthroses and muscle haematomas are rare, except in type 3 disease.

Laboratory findings

See Table 26.2.

- 1 The PFA-100 test (see p. 277) is abnormal. This has largely replaced the bleeding time test.
- 2 Factor VIII levels are often low. If low, a factor VIII/VWF binding assay is performed.
- 3 The APTT may be prolonged.
- 4 VWF levels are usually low.
- 5 There is defective platelet aggregation by patient plasma in the presence of ristocetin (VWF: Rco). Aggregation to other agents (adenosine diphosphate (ADP), thrombin or adrenaline) is usually normal.
- 6 Collagen-binding function (VWF: CB) is usually reduced (but rarely measured).
- 7 Multimer analysis is useful for diagnosing different subtypes (Table 26.3).
- 8 The platelet count is normal except for type 2B disease (where it is low).

Treatment

Options are as follows:

- 1 Local measures and antifibrinolytic agent (e.g. tranexamic acid for mild bleeding).

- 2 DDAVP infusion for those with type 1 VWD. This releases VWF from endothelial cells 30 minutes after intravenous infusion.
- 3 High-purity VWF concentrates for patients with very low VWF levels. Plasma-derived factor VIII/VWF concentrates are used. Recombinant VWF is now in clinical trials.

Hereditary deficiency of other coagulation factors

All these disorders (deficiency of fibrinogen, prothrombin, factors V, VII, combined V and VIII, factors X, XI, XIII or mutation of thrombomodulin) are rare. In all the inheritance is autosomal recessive except for factor XI deficiency where there is variable penetrance. Recombinant factor VII is available for therapy. Factor XI deficiency is seen mainly in Ashkenazi Jews and occurs in either sex. The bleeding risk shows incomplete correlation to severity of the deficiency. Bleeding only occurs after trauma such as surgery. Treatment is with fibrinolytic inhibitor, factor XI concentrate or fresh frozen plasma. Factor XIII deficiency produces a severe bleeding tendency, characteristically with umbilical stump bleeding. Plasma concentrates and recombinant preparation of factor XIII are available.

Acquired coagulation disorders

The acquired coagulation disorders (Table 26.4) are more common than the inherited disorders. Unlike the inherited disorders, multiple clotting factor deficiencies are usual.

Vitamin K deficiency

Fat-soluble vitamin K is obtained from green vegetables and bacterial synthesis in the gut. Deficiency may present in the newborn (haemorrhagic disease of the newborn) or in later life.

Deficiency of vitamin K is caused by an inadequate diet, malabsorption or inhibition of vitamin K by vitamin K antagonist drugs such as warfarin. Warfarin is associated with a decrease in the functional activity of factors II, VII, IX and X and proteins C and S, but immunological methods show normal levels of these factors. The non-functional proteins are called PIVKA (proteins formed in vitamin K absence). Conversion of PIVKA factors to their biologically active forms is a post-translational event involving carboxylation of glutamic acid residues in the N-terminal region (Fig. 26.7). Gamma-carboxylated glutamic acid binds calcium ions, inducing a reversible shape change in the N-termini of vitamin K dependent proteins. This exposes hydrophobic residues which bind to phospholipid. In the process of carboxylation, vitamin K is converted to vitamin K epoxide, which is cycled back to the reduced form by a reductase (VKORC-1). Warfarin interferes with the action of vitamin K epoxide reductase leading to a functional vitamin K deficiency.

Haemorrhagic disease of the newborn

Vitamin K-dependent factors are low at birth and fall further in breast-fed infants in the first few days of life.

Table 26.4 The acquired coagulation disorders.

Deficiency of vitamin K-dependent factors

Haemorrhagic disease of the newborn
Biliary obstruction
Malabsorption of vitamin K (e.g. tropical sprue, gluten-induced enteropathy)
Vitamin K-antagonist therapy (e.g. coumarins, indandiones)
Liver disease – complex dysregulation with synthetic failure of pro- and anticoagulant factors
Disseminated intravascular coagulation – consumption of all clotting factors and platelets

Inhibition of coagulation

Specific inhibitors (e.g. antibodies against factor VIII)
Non-specific inhibitors (e.g. antibodies found in systemic lupus erythematosus, rheumatoid arthritis which paradoxically cause thrombosis)

Miscellaneous

Diseases with M-protein production that interfere with haemostasis
L-Asparaginase
Therapy with heparin, defibrinating agents or thrombolytics
Massive transfusion syndrome

Liver cell immaturity, lack of gut bacterial synthesis of the vitamin and low quantities in breast milk may all contribute to a deficiency which causes haemorrhage, usually on the second to fourth day of life, but occasionally during the first 2 months.

Diagnosis

The PT and APTT are both abnormal. The platelet count and fibrinogen are normal with absent fibrin degradation products.

Treatment

- 1 Prophylaxis. For many years vitamin K has been given to all newborn babies as a single intramuscular injection of 1 mg. This remains the most appropriate and safest treatment. Following epidemiological evidence suggesting a possible link between intramuscular vitamin K and an increased risk of childhood tumours (which has not been substantiated), some centres recommended an oral regimen but this has never been subjected to randomized controlled trial.
- 2 In bleeding infants: vitamin K 1 mg intramuscularly is given every 6 hours with, initially, prothrombin complex concentrate if haemorrhage is severe.

Vitamin K deficiency in children or adults

Deficiency resulting from obstructive jaundice, pancreatic or small bowel disease occasionally causes a bleeding diathesis in children or adults.




Figure 26.7 The action of vitamin K in γ -carboxylation of glutamic acid in coagulation factors which are then able to bind Ca^{2+} and attach to the platelet phospholipid. Warfarin inhibits vitamin K reductase. It is metabolized in the liver and genetic variations in the reductase enzyme VKORC-1 and in the cytochrome CYP2C9 largely account for wide variations in warfarin sensitivity of individuals.

Diagnosis

Both PT and APTT are prolonged. There are low plasma levels of factors II, VII, IX and X.

Treatment

- 1 Prophylaxis: vitamin K 5 mg/day orally.
- 2 Active bleeding or prior to liver biopsy: vitamin K 10 mg slowly intravenously. Some correction of PT is usual within 6 hours. The dose should be repeated on the next 2 days after which optimal correction is usual.
- 3 Rapid correction may be achieved by infusion of prothrombin complex concentrate.

Liver disease

Multiple haemostatic abnormalities contribute to a bleeding tendency and may exacerbate haemorrhage from oesophageal varices.

- 1 Biliary obstruction results in impaired absorption of vitamin K and therefore decreased synthesis of factors II, VII, IX and X by liver parenchymal cells.
- 2 With severe hepatocellular disease, in addition to a deficiency of these factors, there are often reduced levels of factor V and fibrinogen and increased amounts of plasminogen activator.
- 3 Functional abnormality of fibrinogen (dysfibrinogenemia) is found in many patients.
- 4 Decreased thrombopoietin production from the liver contributes to thrombocytopenia.
- 5 Hypersplenism associated with portal hypertension frequently results in thrombocytopenia.
- 6 Disseminated intravascular coagulation (DIC; see below) may be related to release of thromboplastins from damaged

liver cells and reduced concentrations of antithrombin, protein C and α_2 -antiplasmin. In addition, there is impaired removal of activated clotting factors and increased fibrinolytic activity.

- 7 The net haemostatic imbalance in liver disease may be prothrombotic rather than haemorrhagic.

Disseminated intravascular coagulation

Widespread inappropriate intravascular deposition of fibrin with consumption of coagulation factors and platelets occurs as a consequence of many disorders that release procoagulant material into the circulation or cause widespread endothelial damage or platelet aggregation (Table 26.5). It may be associated with a fulminant haemorrhagic or thrombotic syndrome with organ dysfunction or run a less severe and more chronic course. The main clinical presentation is with bleeding but 5–10% of patients manifest thrombotic lesions (e.g. with gangrene of limbs).

Pathogenesis

The key event underlying DIC is increased activity of thrombin in the circulation that overwhelms its normal rate of removal by natural anticoagulants (Fig. 26.8). This can come from tissue factor (TF) release into the circulation from damaged tissues present on tumour cells or from up-regulation of TF on circulating monocytes or endothelial cells in response to pro-inflammatory cytokines (e.g. interleukin-1, tumour necrosis factor, endotoxin).

- 1 DIC may be triggered by the entry of procoagulant material into the circulation in the following situations: severe trauma, amniotic fluid embolism, premature separation of the placenta, widespread mucin-secreting adenocarcino-

Table 26.5 Causes of disseminated intravascular coagulation.*Infections*

Gram-negative and meningococcal septicaemia
Clostridium welchii septicaemia
 Severe falciparum malaria
 Viral infection – varicella, HIV, hepatitis, cytomegalovirus

Malignancy

Widespread mucin-secreting adenocarcinoma
 Acute promyelocytic leukaemia

Obstetric complications

Amniotic fluid embolism
 Premature separation of placenta
 Eclampsia; retained placenta
 Septic abortion

Hypersensitivity reactions

Anaphylaxis
 Incompatible blood transfusion

Widespread tissue damage

Following surgery or trauma
 After severe burns

Vascular abnormalities

Kasabach–Merritt syndrome
 Leaking prosthetic valves
 Cardiac bypass surgery
 Vascular aneurysms

Miscellaneous

Liver failure
 Pancreatitis
 Snake and invertebrate venoms
 Hypothermia
 Heat stroke
 Acute hypoxia
 Massive blood loss

mas, acute promyelocytic leukaemia, liver disease, severe falciparum malaria, haemolytic transfusion reaction and some snake bites.

2 DIC may also be initiated by widespread endothelial damage and collagen exposure (e.g. endotoxaemia, Gram-negative and meningococcal septicaemia, septic abortion), certain virus infections and severe burns or hypothermia. Proinflammatory cytokines and activation of monocytes by bacteria up-regulate tissue factor as well as releasing micro-particles expressing tissue factor into the circulation.

In addition to its role in the deposition of fibrin in the microcirculation, intravascular thrombin formation produces large amounts of circulating fibrin monomers, which form complexes with fibrinogen and interfere with fibrin polymerization, thus contributing to the coagulation defect. Intense fibrinolysis is stimulated by thrombi on vascular walls and the release of split products interferes with fibrin polymerization, thus contributing to the coagulation defect. The combined action of thrombin and plasmin causes depletion of fibrinogen and all coagulation factors. Intravascular thrombin also causes widespread platelet aggregation in the vessels. The bleeding problems which may be a feature of DIC are compounded by thrombocytopenia caused by consumption of platelets.

Clinical features

These are usually dominated by bleeding, particularly from venepuncture sites or wounds (Fig. 26.9a). There may be generalized bleeding in the gastrointestinal tract, the oropharynx, into the lungs, urogenital tract and in obstetric cases, vaginal bleeding may be particularly severe. Less frequently, microthrombi may cause skin lesions, renal failure, gangrene of the fingers or toes (Fig. 26.9b) or cerebral ischaemia.

Some patients may develop subacute or chronic DIC, especially with mucin-secreting adenocarcinoma.





Figure 26.8 The pathogenesis of disseminated intravascular coagulation and the changes in clotting factors, platelets and fibrin degradation products (FDPs) that occur in this syndrome.



(a)



(b)

Figure 26.9 Clinical features of disseminated intravascular coagulation: **(a)** indurated and confluent purpura of the arm; **(b)** peripheral gangrene with swelling and discolouration of the skin of the feet in fulminant disease.

Laboratory findings

See Table 26.6. In many acute syndromes the blood may fail to clot because of gross fibrinogen deficiency.

Tests of haemostasis

- 1 The platelet count is low.
- 2 Fibrinogen concentration is low.
- 3 The thrombin time is prolonged.
- 4 High levels of fibrin degradation products such as D-dimers are found in serum and urine.
- 5 The PT and APTT are prolonged in the acute syndromes.

Compensation by the liver may render some of the coagulation tests normal.

Blood film examination

In many patients there is a haemolytic anaemia ('microangiopathic') and the red cells show prominent fragmentation because of damage caused when passing through fibrin strands in small vessels (see p. 70).

Treatment

Treatment of the underlying cause is most important. The management of patients who are bleeding differs from that of patients with thrombotic problems.

Bleeding

Supportive therapy with fresh frozen plasma (Table 26.7) and platelet concentrates is indicated in patients with dangerous or extensive bleeding. Cryoprecipitate or fibrinogen concentrates provide more concentrated fibrinogen; red cell transfusions may be required.

Thrombosis

The use of heparin or antiplatelet drugs to inhibit the coagulation process is considered in those with thrombotic problems such as skin ischaemia. Fibrinolytic inhibitors should not be used because of consequent failure to lyse thrombi in organs such as the kidney.

Table 26.6 Haemostasis tests: typical results in acquired bleeding disorders.

	Platelet count	Prothrombin time	Activated partial thromboplastin time	Thrombin time
Liver disease	Low	Prolonged	Prolonged	Normal (rarely prolonged)
DIC	Low	Prolonged	Prolonged	Grossly prolonged
Massive transfusion	Low	Prolonged	Prolonged	Normal
Coumarin anticoagulants	Normal	Grossly prolonged	Prolonged	Normal
Heparin	Normal (rarely low)	Mildly prolonged	Prolonged	Prolonged
Circulating anticoagulant	Normal	Normal or prolonged	Prolonged	Normal

DIC, disseminated intravascular coagulation.

Table 26.7 Indications for the use of fresh frozen plasma (National Institutes of Health Consensus Guidelines).

Coagulation factor deficiency (PCC where specific or combined factor concentrate is not available)
Reversal of warfarin effect (only if PCC is unavailable)
Multiple coagulation defects (e.g. in patients with liver disease, DIC) (PCC are much better, plasma is virtually useless)
Massive blood transfusion with coagulopathy and clinical bleeding
Thrombotic thrombocytopenic purpura
Some patients with immunodeficiency syndromes
DIC, disseminated intravascular coagulation; PCC, prothrombin complex concentrates.

Coagulation deficiency caused by antibodies

Circulating antibodies to coagulation factors are occasionally seen with an incidence of approximately 1 per million per year rising markedly with age. Alloantibodies to factor VIII occur in 5–10% of haemophiliacs. Factor VIII autoantibodies may also result in a bleeding syndrome. These immunoglobulin G antibodies occur rarely post-partum, in certain immunological disorders (e.g. rheumatoid arthritis), in cancer and in old age. Treatment consists of a combination of immunosuppression and treatment with factor replacement, usually as human factor VIII, recombinant VIIa or activated prothrombin complex concentrate (FEIBA).

Another protein known as the **lupus anticoagulant** interferes with lipoprotein-dependent stages of coagulation and is usually detected by prolongation of the APTT test (Table 26.6). This inhibitor is detected in 10% of patients with systemic lupus erythematosus (SLE) and in patients with other




Figure 26.10 Thromboelastography (TEG): (a–e) normal trace and appearances in different pathological states. α -angle, speed of solid clot formation; A_{60} , measure of clot lysis or retraction at 60 min; k, clot formation time; MA, absolute strength of fibrin clot; r, rate of initial fibrin formation. Source: From Mallett S.V. and Cox D.J.A. (1992) *Br J Anaesth* 69: 307–13. Reproduced with permission of OUP.

autoimmune diseases who frequently have antibodies to other lipid-containing antigens (e.g. cardiolipin). The antibody is not associated with a bleeding tendency but there is an increased risk of arterial or venous thrombosis and, as with other causes of thrombophilia, an association with recurrent miscarriage (see Chapter 27).

Massive transfusion syndrome

Many factors may contribute to a bleeding disorder following massive transfusion. Blood loss results in reduced levels of platelets, coagulation factors and inhibitors. Further dilution of these factors occurs during replacement with red cells.

Management

Platelet concentrates are given to maintain a platelet count above $75 \times 10^9/L$ or $100 \times 10^9/L$ in cerebral injury or after trauma. The PT and APTT should be kept to less than 1.5 times normal with fresh frozen plasma given initially at 15 mL/kg. It is usually necessary to give 4–6 units of FFP for every 6 units of red cells transfused. FFP is started early with red cell transfusion. Some protocols include 1:1:1 for red cells, platelet packs and FFP. Cryoprecipitate or fibrinogen concentrate is given to keep fibrinogen above 1.5 g/L. Trials of fibrinogen concentrates are in progress in

obstetric emergencies and perioperative bleeding during cardiac surgery. Monitoring by thromboelastography or ROTEM (see below) is also used.


Thromboelastography: near-patient testing

Thromboelastography (TEG) and thromboelastometry (ROTEM) are techniques for a global assessment of haemostatic function of a single blood sample, in which the reaction of platelets with the protein coagulation cascade is observed from the time of the initial platelet fibrin interaction through platelet aggregation, clot strengthening and fibrin cross-linkage to eventual clot lysis. It is suited as a monitor of haemostasis in surgery (e.g. of the liver or heart) associated with haemostatic defects. Freshly drawn blood is placed in a cuvette which is oscillated, the motion being transferred to a pin which transmits deflection as torsion to a photoelectric detector with computerized data capture. As fibrin strands form, the fibrin clot affects movement of the pin. The normal trace shows the rate of initial fibrin formation, the time to formation of a clot (coagulation time), strength of the fibrin clot, clot lysis index or retraction. Typical patterns showing results in fibrinolysis, hypercoagulability, haemophilia and thrombocytopenia are shown in Figure 26.10.

- Coagulation disorders may be inherited or acquired.
- Haemophilia A is the most common inherited deficiency of a clotting factor. It is severe if factor VIII activity in plasma is <1% of normal. It presents with excess bruising or prolonged bleeding after trauma and spontaneous bleeding, usually into muscles and joints, which can result in joint deformity.
- Many older patients are infected with hepatitis C or HIV as a result of receiving contaminated blood products.
- The APTT is prolonged and PT normal.
- Antenatal diagnosis is usually carried out by polymerase chain reaction (PCR) techniques, the gene being carried on the X chromosome.
- Treatment is with recombinant or concentrates of factor VIII, or with drugs (e.g. DDAVP (desmopressin)). New long-acting recombinant preparations are being introduced.
- Factor IX deficiency has a similar pattern of inheritance and clinical manifestations.
- Gene therapy for factor VIII and factor IX deficiencies is under trial.
- Von Willebrand disease (VWD) is the most frequent inherited bleeding disorder. Haemorrhage occurs from mucous membranes, skin cuts and post-trauma. It usually has a dominant inheritance. Platelet function is abnormal and VWF levels usually low.
- Acquired coagulation disorders include those caused by vitamin K deficiency (e.g. in the newborn or with malabsorption) or caused by vitamin K antagonist therapy (e.g. warfarin).
- Other common coagulation abnormalities are those in liver disease caused by reduced synthesis of coagulation factors and in disseminated intravascular coagulation (DIC), which causes consumption of coagulation factors and platelets.
- Fresh frozen plasma is used in treatment of multiple coagulation defects, or specific defects if the appropriate concentrate is not available, and in therapy for thrombotic thrombocytopenic purpura.



Now visit www.wileyessential.com/haematology to test yourself on this chapter.


A vertical strip on the left side of the page showing a microscopic view of blood cells. It includes several red blood cells (erythrocytes) which are biconcave and reddish-pink, and a few white blood cells (leukocytes) which are larger and have more complex, multi-lobed nuclei. The background is a dark blue.

CHAPTER 27

Thrombosis 1: pathogenesis and diagnosis

Key topics

■ Arterial thrombosis	303
■ Venous thrombosis	303
■ Investigation of thrombophilia	307
■ Diagnosis of venous thrombosis	308



Thrombi are solid masses or plugs formed in the circulation from blood constituents. Platelets and fibrin form the basic structure. Their clinical significance lies in the ischaemia that results from local vascular obstruction or distant embolization. Thrombi are involved in the pathogenesis of myocardial infarction, cerebrovascular disease, peripheral arterial disease, deep vein thrombosis (DVT) and pulmonary embolism (PE).

Thrombosis, both arterial and venous, is more common as age increases and is frequently associated with risk factors such as surgery or pregnancy. The term *thrombophilia* is used to describe inherited or acquired disorders of the haemostatic mechanism that predispose to thrombosis.

Arterial thrombosis

Pathogenesis

Atherosclerosis of the arterial wall, plaque rupture and endothelial injury expose blood to subendothelial collagen and tissue factor. This initiates the formation of a platelet nidus on which platelets adhere and aggregate.

Platelet deposition and thrombus formation are important in the pathogenesis of atherosclerosis. Platelet-derived growth factor (PDGF) stimulates the migration and proliferation of smooth muscle cells and fibroblasts in the arterial intima. Regrowth of endothelium and repair at the site of arterial damage results in thickening of the vessel wall. The intrinsic pathway of fibrin formation (see Fig. 24.10) is involved in pathological thrombosis *in vivo* by contact activation on damaged blood vessels.

As well as blocking arteries locally, emboli of platelets and fibrin may break away from the primary thrombus to occlude distal arteries. Examples are carotid artery thrombi leading to cerebral thrombosis and transient ischaemic attacks, and heart valve and chamber thrombi leading to systemic emboli and infarcts (Fig. 27.1).

Clinical risk factors

The risk factors for arterial thrombosis are related to the development of atherosclerosis and are listed in Table 27.1. The identification of patients at risk is largely based on clinical assessment. A number of epidemiological studies have resulted in the construction of coronary artery thrombosis risk profiles based on gender, age, elevated blood pressure, high levels of serum cholesterol, glucose intolerance, cigarette smoking and electrocardiogram abnormalities. These profiles have allowed pre-symptomatic assessment of young and apparently fit subjects and are valuable in counselling a change in lifestyle or for recommending medical therapy in individuals at risk.

Venous thrombosis

Pathogenesis and risk factors

Virchow's triad suggests that there are three components that are important in thrombus formation:

- 1 Slowing down of blood flow;
- 2 Hypercoagulability of the blood;
- 3 Vessel wall damage.




Figure 27.1 Arteriogram showing saddle embolus at the aortic bifurcation (dotted arrow) and embolus in the left common iliac artery (solid arrow).

For venous thrombosis, increased systemic coagulability and stasis are most important, with vessel wall damage being somewhat less important than in arterial thrombosis, although it may be important in patients with sepsis, in-dwelling

Table 27.1 Risk factors for arterial thrombosis (atherosclerosis).

Positive family history
Male sex
Hyperlipidaemia
Hypertension
Diabetes mellitus
Gout
Polycythaemia
Hyperhomocysteinaemia
Cigarette smoking
ECG abnormalities
Elevated CRP, IL6, fibrinogen, lipoprotein-associated phospholipase A ₂
Lupus anticoagulant
Collagen vascular diseases
Behçet's disease
CRP, C-reactive protein; ECG, electrocardiogram.

Table 27.2 Hereditary and acquired risk factors for venous thrombosis.

<i>Hereditary haemostatic disorders</i>
Factor V Leiden
Prothrombin G20210A variant
Protein C deficiency
Antithrombin deficiency
Protein S deficiency
Dysfibrinogenaemia
Non-O ABO blood group
DVT in close relative (especially if unprovoked)
<i>Hereditary or acquired haemostatic disorders</i>
Raised plasma levels of factor VIII
Raised plasma levels of fibrinogen
Raised plasma levels of homocysteine
<i>Acquired disorders</i>
Lupus anticoagulant
Oestrogen therapy (oral contraceptive and HRT)
Heparin-induced thrombocytopenia
Pregnancy and puerperium
Surgery, especially abdominal, hip and knee surgery
Major trauma
Malignancy
Acutely ill hospitalized medical patients including cardiac or respiratory failure, infection, inflammatory bowel disorders
Myeloproliferative disease
Hyperviscosity, polycythaemia
Stroke
Pelvic obstruction
Nephrotic syndrome
Dehydration
Varicose veins
Previous superficial vein thrombosis
Age
Obesity
Paroxysmal nocturnal haemoglobinuria
Behçet's disease
HRT, hormone replacement therapy.

catheters and sites of damage to veins by previous thrombosis. Stasis allows the completion of blood coagulation at the site of initiation of the thrombus (e.g. behind the valve pockets of the leg veins in immobile patients). Table 27.2 lists a number of recognized risk factors.

Hereditary disorders of haemostasis

The prevalence of inherited disorders associated with increased risk of thrombosis is higher than that of hereditary bleeding disorders. Approximately one third of patients who suffer deep vein thrombosis (DVT) or pulmonary embolus (PE) have an identifiable heritable risk factor: rare deficiencies of antithrombin, protein C or protein S, or

common mutations affecting factor V (factor V Leiden) or prothrombin (Table 27.2). Venous thromboembolism frequently results from gene–environment interaction, so additional risk factors (surgery, immobility, oestrogen exposure) are often present in patients with heritable thrombophilia when they develop thrombosis. Although heritable thrombophilia partly explains the gene–environment interaction leading to clinical expression of disease, testing for heritable thrombophilic defects has limited clinical utility as a positive test result rarely predicts a high risk of recurrence compared to patients without identifiable abnormalities. The history of a spontaneous DVT in a close relative increases an individual's risk of DVT even if no known genetic predisposition can be identified. However, testing for heritable thrombophilia can help clinical decision making in thrombosis-prone families.

Factor V Leiden gene mutation

This is the most common inherited cause of an increased risk of venous thrombosis. It occurs in approximately 3–7% of factor V alleles in white people (Fig. 27.2). There is failure of activated protein C (APC) to prolong the activated partial thromboplastin time (APTT) test when added to plasma and the phenotype is sometimes referred to as 'activated protein C resistance'. Activated protein C normally breaks down activated factor V and so it should slow the clotting reaction and prolong the APTT. APC resistance is caused by a genetic polymorphism in the factor V gene, which makes factor V less susceptible to cleavage by APC (Fig. 27.3). The frequency of factor V Leiden in the general population in Western countries means that it cannot be regarded as a rare mutation but instead is a genetic polymorphism that is maintained in the population (Fig. 27.2). Presumably, individuals with this allele have been 'selected', probably because of a reduced bleeding tendency (e.g. post-partum). It does not increase the risk of arterial thrombosis. A small minority of patients with APC resistance do not have factor V Leiden but have other mutations of factor V.

Patients who are heterozygous for factor V Leiden are at an approximately five- to eight-fold increased risk of venous thrombosis compared to the general population, but only 10% of carriers develop thrombosis during their lifetime. Individuals who are homozygous have a 30–140-fold risk. Following venous thrombosis they have a higher risk of re-thrombosis than individuals with DVT but normal factor V.

The incidence of factor V Leiden in patients with venous thrombosis is approximately 20–40%. Genetic screening is relatively easy and is widely performed. However, even if a patient with DVT is a carrier of factor V Leiden their absolute risk of thrombosis is still very low in the absence of other risk factors. At present it is not recommended to start anticoagulation therapy in individuals with the Leiden mutation, even if homozygous, if they have no history of thrombosis.




Figure 27.2 The incidence of carriers of factor V Leiden in different countries.




Figure 27.3 The genetic basis of factor V Leiden. **(a)** Activated protein C (APC) inactivates factor Va by proteolytic cleavage at three sites in the Va heavy chain. **(b)** In the factor V Leiden mutation the Arg506Gln polymorphism leads to glutamine at position 506 with less efficient inactivation of factor V by APC and increased risk of thrombosis.

Antithrombin deficiency

Inheritance is autosomal dominant. There are recurrent venous thromboses usually starting in early adult life and arterial thrombi may occur. Antithrombin concentrates

are available and are occasionally used to prevent thrombosis during surgery or childbirth. Many molecular variants of antithrombin have been categorized and are associated with varying degrees of risk of thrombosis. Treatment with heparin, including low molecular weight heparin, is effective in the vast majority of patients and treatment of DVT and PE is the same for patients with and without antithrombin deficiency.

Protein C deficiency

Inheritance is autosomal dominant with variable penetrance. Protein C levels in heterozygote individuals are approximately 50% of normal. Characteristically, many patients develop skin necrosis as a result of dermal vessel occlusion when treated with warfarin, thought to be caused by a further reduction of protein C levels in the first day or two of warfarin therapy. Rarely, infants may be born with homozygous deficiency and characteristically present with severe disseminated intravascular coagulation (DIC) or purpura fulminans in infancy. Administration of protein C was once thought to be beneficial in patients with sepsis but it is no longer generally used.

Protein S deficiency

Protein S deficiency has been found in a number of families with a thrombotic tendency. It is a cofactor for protein C. The clinical features of protein S deficiency are similar to those of protein C deficiency, including a tendency to skin

necrosis with warfarin therapy. The inheritance is autosomal dominant.

Prothrombin allele G20210A

G20210A is an allele of the prothrombin gene that has a prevalence of 2–3% in the population. It leads to increased plasma prothrombin levels and increases thrombotic risk five-fold. It is probable that the cause of venous thrombosis with this mutation is that sustained generation of thrombin results in prolonged down-regulation of fibrinolysis through activation of thrombin-activated fibrinolysis inhibitor (see p. 275).

Hyperhomocysteinaemia

High levels of plasma homocysteine may be genetic or acquired and are associated with increased risk for both venous and arterial thrombosis. However, trials show little evidence that lowering the levels reduces these risks. Homocysteine is derived from dietary methionine and is removed by either remethylation to methionine (see Fig. 5.3) or conversion to cysteine via a *trans*-sulphuration pathway. Classic homocystinuria is a rare autosomal recessive disorder caused by deficiency of cystathione β -synthase, the enzyme responsible for *trans*-sulphuration. Vascular disease and thrombosis are major features of the disease. Methylenetetrahydrofolate reductase is involved in the remethylation of tetrahydrofolate (THF), methylTHF being responsible for methylating homocysteine to methionine (see Fig. 5.5). A common thermolabile variant of the enzyme may be responsible for mild homocysteinaemia (above 15 $\mu\text{mol/L}$). In people homozygous for this mutation and who have had a DVT, long-term folic acid therapy is indicated. Other acquired risk factors for hyperhomocysteinaemia include deficiency of vitamin B₆, drugs (e.g. ciclosporin), renal damage and smoking. The levels also increase with age and are higher in men and post-menopausal females.

Defects of fibrinogen

Defects of fibrinogen are usually clinically silent or cause excess bleeding. Thrombosis is a rare association.

ABO blood group

Non-O blood group carriers have a higher risk of venous thrombosis or embolism than O carriers. This is related to their higher plasma levels of von Willebrand factor and factor VIII.

Hereditary or acquired disorders of haemostasis

High plasma factor VIII or fibrinogen levels are also associated with venous thrombosis. The combination of multiple risk factors is associated with increased risk of thrombosis. If these are persistent they may represent a reason for extended anticoagulation.

Acquired risk factors

These may cause thrombosis in patients without another identifiable abnormality but are more likely to do so if an inherited predisposing abnormality such as factor V Leiden is also present.

Hospital-acquired thrombosis (HAT)

Hospital-acquired thrombosis is responsible for up to 50% of cases of DVT and PE. Thrombosis may occur many weeks after discharge from hospital and HAT is now regularly defined as venous thromboembolism occurring within 90 days of hospitalization. In many countries there are national strategies to reduce the incidence of HAT, such as through universal patient risk assessment on admission and administration of thromboprophylaxis to high-risk patients. Thromboprophylaxis is typically given for the duration of admission and is extended after discharge in very high-risk patients such as those having had hip or knee replacement.

Postoperative venous thrombosis

This is more likely to occur in the elderly, obese, those with a previous or family history of venous thrombosis, and in those in whom major abdominal or hip operations are performed. Elasticated stockings or mechanical methods (see p. 318) are used to reduce the risk of DVT.

Venous stasis and immobility

These factors are probably responsible for the high incidence of postoperative venous thrombosis and for venous thrombosis associated with congestive cardiac failure, myocardial infarction and varicose veins. In atrial fibrillation, thrombin generation from accumulation of activated clotting factors leads to a high-risk clot formation in the atrial appendage and consequent systemic embolization. The use of muscle relaxants during anaesthesia may also contribute to venous stasis. Venous thrombosis also has a higher frequency after prolonged aeroplane journeys.

Malignancy

Patients with carcinoma of the ovary, brain and pancreas have a particularly increased risk of venous thrombosis or its recurrence but there is an increased risk with all cancers and myeloproliferative diseases. The tumours produce tissue factor and a procoagulant that directly activates factor X. Mucin-secreting adenocarcinomas may be associated with DIC.

Inflammation

This up-regulates procoagulant factors and down-regulates anticoagulant pathways, particularly protein C. Thrombosis is particularly likely in inflammatory bowel disease, Behçet's disease, systemic tuberculosis, systemic lupus erythematosus (SLE) and diabetes.

Superficial venous thrombosis

The risk of DVT is increased in those who have had a previous superficial venous thrombosis or thrombophlebitis, especially if there are additional risk factors.

Blood disorders

Increased viscosity, thrombocytosis and enhanced platelet functional responses are possible factors for the high incidence of thrombosis in patients with polycythaemia vera and essential thrombocythaemia. Testing for the *JAK2* V617F mutation may indicate an otherwise unsuspected myeloproliferative disease in patients with hepatic or portal vein thrombosis. The *CALR* mutation is rarely associated with DVT (see p. 172). There is a high incidence of venous thrombosis in patients with paroxysmal nocturnal haemoglobinuria, including thrombi in large veins such as the hepatic vein. An increased tendency to venous thrombosis has been observed in patients with sickle cell disease, during post-splenectomy thrombocytosis and those with a paraprotein.

Oestrogen therapy

Oestrogen therapy, particularly high-dose therapy, is associated with increased plasma levels of factors II, VII, VIII, IX and X, and depressed levels of antithrombin and tissue plasminogen activator in the vessel wall. There is a high incidence of postoperative venous thrombosis in women on high-dose oestrogen therapy and full-dose oestrogen-containing oral contraceptives. The risk is much less with low-dose oestrogen contraceptive preparations. Hormone replacement therapy also increases the risk of thrombosis, largely obviated by the use of low-oestrogen preparations.

The antiphospholipid syndrome

The antiphospholipid syndrome (APS) can be defined as the occurrence of venous and arterial thrombosis and/or recurrent miscarriage in association with laboratory evidence of persistent antiphospholipid antibody. One antiphospholipid is the lupus anticoagulant (LA) which was initially detected in patients with SLE and is identified by a prolonged plasma APTT which does not correct with a 50:50 mixture of normal plasma. Paradoxically, in view of its name, it is associated with venous and arterial thrombosis. A second test dependent on limiting quantities of phospholipid (such as the dilute Russell's viper venom test) is also used in diagnosis. Whereas lupus anticoagulants are reactive in the fluid phase, other antiphospholipid antibodies, such as anticardiolipin antibodies and antibodies to β_2 -GPI-1, are identified by solid phase immunoassay. Both solid phase assays and coagulation tests for LA should be used in the diagnosis of APS.

As well as patients with SLE, antiphospholipid antibodies are also found in other autoimmune disorders particularly of connective tissues, lymphoproliferative diseases, post-viral

Table 27.3 Clinical associations of lupus anticoagulant and anticardiolipin antibodies.

Venous thrombosis: deep venous thrombosis/pulmonary embolism, renal, hepatic, retinal veins

Arterial thrombosis

Recurrent fetal loss

Thrombocytopenia

Livedo reticularis

N.B. Recurrent fetal loss may also occur in other types of thrombophilia.

infections, with certain drugs including phenothiazines and as an 'idiopathic' phenomenon in otherwise healthy subjects. Arterial thrombosis may cause peripheral limb ischaemia, stroke or myocardial infarction. Venous thrombosis includes DVT, PE and thrombosis in vessels supplying the abdominal organs. As with other causes of thrombophilia, recurrent abortion caused by placental infarction is also an association (Table 27.3). Thrombocytopenia may be present and livedo reticularis is a frequent dermal manifestation.

Treatment is with anticoagulation where indicated. It is usual to maintain an international normalized ratio (INR) of between 2.0 and 3.0 with warfarin but higher levels may be needed if previous arterial or major DVT has occurred or recurrence of thrombosis occurs on warfarin therapy. Low-dose heparin and aspirin are useful in the management of recurrent miscarriage.

Collagen vascular diseases and Behçet's syndrome are also associated with arterial and venous thrombosis, whether or not the lupus anticoagulant is present.

Investigation of thrombophilia

Decisions regarding duration of anticoagulation, such as whether to continue it lifelong or for a defined period, should be made with reference to whether the first episode of venous thrombosis was provoked or not, other risk factors, and risk of anticoagulant therapy-related bleeding, regardless of whether a heritable thrombophilia is known. Testing for heritable thrombophilia may influence decisions regarding duration of anticoagulation and for family counselling. However routine testing of all patients with an episode of venous thromboembolism is not generally valuable.

Tests that may be abnormal in patients with a tendency to venous thrombosis include:

- 1 Blood count and erythrocyte sedimentation rate – to detect elevation in haematocrit, white cell count, platelet count, fibrinogen and globulins.
- 2 Blood film examination – may provide evidence of myeloproliferative disorder; leucoerythroblastic features may indicate malignant disease.

- 3 Prothrombin time (PT) and APTT – a shortened APTT is often seen in thrombotic states and may indicate the presence of activated clotting factors. A prolonged APTT test, not corrected by the addition of normal plasma, suggests an LA or an acquired inhibitor to a coagulation factor.
- 4 Anticardiolipin and anti- β_2 -GPI antibodies.
- 5 Thrombin time (and reptilase time) – prolongation suggests an abnormal fibrinogen.
- 6 Fibrinogen assay.
- 7 DNA analysis for factor V Leiden.
- 8 Antithrombin – immunological and functional assays.
- 9 Protein C and protein S – immunological and functional assays.
- 10 Prothrombin gene analysis for the G20210A variant.
- 11 Plasma homocysteine estimation.
- 12 Test for CD59 and CD55 expression (paroxysmal nocturnal haemoglobinuria) in red cells if paroxysmal nocturnal haemoglobinuria is suspected.
- 13 Test for *JAK2* (V617F) mutation if portal or hepatic vein thrombosis.
- 14 Protein electrophoresis for paraprotein.

Table 27.4 Deep vein thrombosis: clinical assessment: the Wells score.

	Points
Active cancer (treatment ongoing or within previous 6 months or palliative)	1
Paralysis, plaster	1
Bed more than 3 days, surgery within 4 weeks	1
Tenderness along veins	1
Entire leg swollen (calf circumference >3cm compared with other leg)	1
Pitting oedema	1
Collateral veins	1
Alternative diagnosis likely	-2
<i>Low probability 0–1</i>	
<i>High probability 2 or more</i>	

Diagnosis of venous thrombosis

Deep vein thrombosis

Clinical suspicion DVT is suspected in patients with a painful swollen limb. It is more common in those with previous DVT or superficial venous thrombosis, cancer or recent confinement to bed (Table 27.2). In the leg, unilateral thigh or calf swelling or tenderness, pitting oedema and the presence of collateral superficial non-varicose veins are important signs. Homan's sign (pain in the calf on flexing the ankle) is unreliable.

Plasma D-dimer concentration The concentration of these fibrin breakdown products is raised when there is a fresh thrombosis. It is a useful assay when venous thrombosis is suspected and the Wells score can be used to predict the chance of thrombosis (Table 27.4). A negative result in emergency departments can be used to exclude DVT or PE and avoid the need for radiological imaging. The test is useful when a new thrombus is suspected at a site of scarring from a previous thrombosis and scanning gives equivocal findings. D-dimer elevation in cancer, inflammation, after surgery or trauma limits its usefulness. It is also useful when ultrasound is not possible.

Compression ultrasound This is a reliable and practical method for patients with first suspicion of DVT in the legs and other sites (Fig. 27.4a). It can be combined with spectral, colour (Fig. 27.4a) or power Doppler (duplex) scanning which improves accuracy by focusing on individual veins. It does not distinguish between acute and chronic thrombi. Persisting venous obstruction detected by ultrasonography at the completion of warfarin therapy is not typically associated

with an increased risk of recurrent thrombosis and scanning to detect residual vein occlusion (RVO) is not routine practice.

Contrast venography This test is now rarely performed. Iodinated contrast medium is injected into a vein peripheral to the suspected DVT. This permits direct demonstration by X-ray of the site, size and extent of the thrombus (Fig. 27.4b). However, it is a painful invasive technique, with a risk of contrast reaction and procedure-induced DVT.

Magnetic resonance imaging (MRI) This may also be used but is expensive. It is indicated when ultrasound might be inaccurate or difficult, such as because of obesity or patients in a plaster cast.

Impedance plethysmography is less sensitive and accurate and is falling out of use.

Pulmonary embolus

This usually presents with shortness of breath and there may be pleuritic chest pain. PE should be particularly suspected in patients with signs or previous history of DVT, immobilization for more than 2 days or recent (<4 weeks) surgery, haemoptysis or cancer. Recurrent PE may lead to pulmonary hypertension.

Chest X-ray This is often normal but may show evidence of pulmonary infarction or pleural effusion.

Electrocardiogram This is performed to determine whether there is right heart 'strain' which occurs only in relatively severe cases.

Plasma D-dimer concentration D-dimer measurement using an appropriate assay has a similar negative predictive value in patients with PE as with DVT when used in conjunction with a clinical probability rule. As such PE can be reliably excluded in a patient with a low pre-test probability score and a normal D-dimer test.

Ventilation perfusion (VQ) scintigraphy This detects areas of the lung being ventilated but not perfused.

Computed tomography (CT) pulmonary angiography Fine slices of the lung are scanned by spiral CT so that filling defects in the pulmonary arteries are visualized (Fig. 27.4c).

Magnetic resonance pulmonary angiography Gadolinium-enhanced MRI is a relatively new and expensive but accurate technique.

Pulmonary angiography This is the traditional reference method but is invasive with complications, albeit uncommon, such as arrhythmia or contrast reaction.





Figure 27.4 Diagnostic imaging of deep vein thrombosis (DVT) and pulmonary embolus (PE). **(a)** Colour power Doppler ultrasound of the right femoral vessels with compression shows normal flow in the femoral artery but absent flow in the vein because of thrombus. A normal vein would collapse with compression of the probe. Source: Courtesy of Dr Tony Young. **(b)** Femoral venogram demonstrating extensive thrombus within the right external iliac vein. Source: Courtesy of Dr I.S. Francis and Dr A.F. Watkinson. **(c)** Computed tomography (CT) pulmonary angiography: a coronal image shows bilateral filling defects (green crosses) in the central pulmonary arteries indicating pulmonary emboli. Source: Courtesy of Dr Tony Young.

SUMMARY

- Thrombosis is the formation of solid masses of platelets and fibrin in the circulation. It may be arterial or venous.
- Arterial thrombosis is mainly related to atherosclerosis of the vessel wall with risk factors such as hypertension, hyperlipidaemia, smoking and diabetes.
- Venous thrombosis is related to genetic coagulation factor abnormalities (e.g. factor V Leiden), stasis of the circulation or to an acquired increase in coagulation factors (e.g. oestrogen therapy, postoperative, pregnancy) or to unknown factors (e.g. age or obesity).
- Diagnosis of deep vein thrombosis is with serial compression ultrasound combined with Doppler (duplex) scanning, contrast venography or MRI imaging. Plasma D-dimer concentration assay may help.
- Pulmonary embolus is diagnosed by chest X-ray, electrocardiogram, ventilation perfusion scintigraphy or CT angiography.



Now visit www.wileyessential.com/haematology to test yourself on this chapter.


A vertical strip on the left side of the page showing a microscopic view of blood cells. It features several red blood cells (erythrocytes) in shades of red and pink, and a few white blood cells (leukocytes) with distinct nuclei. The background is a dark blue.

CHAPTER 28

Thrombosis 2: treatment

Key topics

■ Anticoagulant drugs	312
■ Heparin	312
■ Low molecular weight heparin	314
■ Direct-acting parenteral anticoagulants	315
■ Oral anticoagulants	315
■ Direct-acting oral anticoagulant drugs	317
■ Post-thrombotic syndrome	318
■ Mechanical methods of prophylaxis of venous thrombosis	318
■ Fibrinolytic agents	318
■ Antiplatelet drugs	318



Anticoagulant drugs

Anticoagulant drugs are used widely in the treatment of venous thromboembolic disease. Their value in the treatment of arterial thrombosis is less well established. A wide variety of orally or parenterally active drugs is available (Fig. 28.1), acting either directly or indirectly at a specific site or at multiple sites in the coagulation cascade (Fig. 28.2).

Heparin

This acidic unfractionated mucopolysaccharide of average molecular weight (MW) 15 000–18 000 is an inhibitor of blood coagulation because of its action in potentiating the activity of antithrombin (see below). As it is not absorbed from the gastrointestinal tract it must be given by injection. It is inactivated by the liver and excreted in the urine. The effective biological half-life is approximately 1 hour (Table 28.1).




Figure 28.1 Classification of the anticoagulants based on their route of administration and mode of action. Source: Adapted from DeCaterina R. *et al.* (2013) *Thrombosis and Haemostasis* 109: 4. LMWH, low molecular weight heparin; UFH, unfractionated heparin.




Figure 28.2 Sites of action of the more frequently used anticoagulants. Source: Adapted from DeCaterina R. *et al.* (2013) *Thrombosis and Haemostasis* 109: 4.

Table 28.1 Comparison of unfractionated heparin with low molecular weight heparin.

	Unfractionated heparin	Low molecular weight heparin
Mean molecular weight in kilodaltons (range)	15 (4–30)	4.5 (2–10)
Anti-Xa:anti-IIa	1:1	2:1 to 4:1
Inhibits platelet function	Yes	No
Bioavailability	50%	100%
Half-life		
intravenous	1 hour	2 hours
subcutaneous	2 hours	4 hours
Elimination	Renal and hepatic	Renal
Monitoring	APTT	Xa assay (usually not needed)
Frequency of heparin-induced thrombocytopenia	High	Low
Osteoporosis	Yes	Less frequent

APTT, activated partial thromboplastin time.

Mode of action

Heparin dramatically potentiates the formation of complexes between antithrombin and the activated serine protease coagulation factors, thrombin (IIa) and factors IXa, Xa and XIa (Fig. 28.3). This complex formation inactivates these factors irreversibly. In addition, heparin impairs platelet function.




Figure 28.3 The action of heparin. This activates antithrombin which then forms complexes with activated serine protease coagulation factors (thrombin, Xa, IXa and XIa) and so inactivates them.

Low molecular weight heparin (LMWH) preparations (MW 2000–10 000) have a greater ability to inhibit factor Xa than to inhibit thrombin and interact less with platelets than standard heparin, and so may have a lesser tendency to cause bleeding. They also have greater bioavailability and a more prolonged half-life in plasma, making once-daily administration feasible (Table 28.1). Fondaparinux is a synthetic pentasaccharide which is a specific indirect factor Xa inhibitor.

Indications

LMWH is used routinely for treatment of deep vein thrombosis (DVT), pulmonary embolus (PE) and unstable angina pectoris. Initial treatment of PE may also involve thrombolysis in severe cases. LMWH is also used widely in the prophylaxis of venous thrombosis and is the drug of choice for women requiring anticoagulation in pregnancy because heparin does not cross the placenta. Heparin is also used during cardiopulmonary bypass surgery, for maintaining the patency of indwelling venous lines and in some case of disseminated intravascular coagulation (DIC) if the manifestations are predominantly vaso-occlusive.

Administration and laboratory control

Standard unfractionated heparin

Continuous intravenous infusion provides the smoothest control of heparin therapy and is the treatment of choice where rapid reversal of anticoagulation by protamine sulphate may be required (e.g. in surgical patients or late pregnancy). It has largely been replaced for treatment of acute PE and DVT by

LMWH. In an adult, 1000–2000 units/hour with a loading dose of 5000 units is usually satisfactory.

Therapy is monitored by maintaining the APTT at 1.5–2.5 times the ULN (upper limit of the normal) value. Unfractionated heparin is better monitored by an anti-Xa assay where this is available as it is a more reliable method than the APTT for this purpose. It is usual to start warfarin or a newer oral anticoagulant within 2 days of starting heparin and to discontinue heparin when the INR has been above 2.0 on 2 successive days. For acute coronary syndromes, both unfractionated heparin and LMWH are of benefit when used with aspirin in the prevention of mural thrombosis, systemic embolization and venous thrombosis. Standard heparin is widely used for keeping intravenous lines patent.

Low molecular weight heparin

LMWH is given by subcutaneous injection and, as it has a longer half-life than standard heparin, it can be given once a day in prophylaxis, or once or twice daily for treatment (Table 28.1). There are many commercial preparations. Compared with unfractionated heparin, LMWH has a more predictable dose response which avoids the need for routine monitoring. LMWH is the treatment of choice for DVT, PE and unstable angina. Patients with uncomplicated DVTs are now managed initially with regular LMWH injections, often at home, once or twice daily according to the preparation. It is also the preferred treatment of DVT in cancer patients and in pregnancy because it does not cross the placenta. Although routine monitoring is not required, measurement of anti-Xa peak levels 4 hours after injection allows dose adjustment in selected patients (e.g. in pregnancy, renal failure, gross obesity and in children).

LMWH is used for the prevention of DVT in both medical and surgical patients. It is now mandatory in many hospitals to have a policy for prevention of DVT and PE for all hospitalized medical and surgical patients. LMWHs are the gold standard pharmaceutical agents but new oral anticoagulants (see Table 28.5) may supersede them.

Bleeding during heparin therapy

Bleeding may occur because of excessive prolonged anticoagulation or due to the antiplatelet functional effect of heparin. Intravenous heparin has a half-life of less than 1 hour and it is usually only necessary to stop the infusion. Protamine is able to inactivate heparin immediately. However, protamine itself may act as an anticoagulant when in excess.

Heparin-induced thrombocytopenia

A mild lowering of the platelet count may occur in the first 24 hours as a result of platelet clumping. This is of no clinical consequence (heparin-induced thrombocytopenia (HIT) type 1). The important HIT, type 2, may occur in up to 5% of patients who are treated with unfractionated heparin and paradoxically presents with thrombosis. It results from the binding of heparin to platelet factor 4 (PF4) followed by the generation of an IgG antibody to the heparin–PF4 complex, which leads to platelet activation, thrombocytopenia and thrombosis (Fig. 28.4). Typically, it presents as a fall of more than 50% in the platelet count 5 or more days after starting heparin treatment or earlier if heparin has been given previously. Diagnosis is difficult but assays have recently been developed to allow the detection of antibodies to immobilized heparin–PF4 complex.




Figure 28.4 Mechanism of heparin-induced thrombocytopenia (HIT). Platelet factor 4 (PF4) is released from α granules and forms a complex on the platelet surface with heparin. Immunoglobulin G antibodies develop against this complex and can activate the platelet through the platelet immunoglobulin receptor Fc- γ RII. This leads to platelet stimulation, further release of PF4 and the platelet release reaction with consequent thrombocytopenia and thrombus development.

Heparin therapy must be discontinued. Argatroban or fondaparinux are used in patients with HIT. LMWH is less likely than unfractionated heparin to cause HIT. Warfarin therapy should not be started until normalization of the platelet count has been achieved.

Osteoporosis

This occurs with long-term (more than 2 months) heparin therapy, especially in pregnancy. The drug complexes minerals from the bones but the exact pathogenesis is unknown. It is less frequent with LMWH therapy.

Direct-acting parenteral anticoagulants

Fondaparinux, a synthetic analogue of the antithrombin-binding pentasaccharide of heparin, is an indirect factor Xa inhibitor. It is given subcutaneously, has a plasma half-life of 17 hours and, like the orally active factor Xa inhibitors, does not require laboratory monitoring (by measuring factor Xa levels) except in especially obese patients, those with renal failure and in children.

Bivalirudin is used as an alternative to heparin in patients undergoing percutaneous coronary interventions.

Argatroban is a small molecule direct inhibitor of thrombin given by continuous intravenous infusion. It is used to treat patients with HIT.

Oral anticoagulants

Until recently the only available drugs had been derivatives of coumarin or phenindione. **Warfarin**, a coumarin, is most widely used. Coumarins are vitamin K antagonists (see p. 296) and treatment results in decreased biological activity of the vitamin K-dependent factors II, VII, IX and X (see Fig. 26.7). After warfarin is given, factor VII levels fall considerably within 24 hours but prothrombin has a longer plasma half-life and only falls to 50% of normal at 3 days; only then is the patient fully anticoagulated.

Principles of oral anticoagulation with VKAs

A typical starting regimen for warfarin would be 5 mg on days 1, 2 and 3. The daily dose should be monitored and adjusted by measurement of the INR and algorithms for this purpose are published. The initial dose can be 'tailor-made' using an algorithm based on clinical variables and genetic information on two genes involved in warfarin metabolism or action, cytochrome p450 (CYP2C9) and vitamin K epoxide reductase (see Fig. 26.7). In practice few units do this. The usual maintenance dosage of warfarin is 3–9 mg/day but individual responses vary greatly. Lower loading dosage is recommended for the elderly or those with liver disease.

International normalized ratio (INR)

The effect of oral anticoagulants is monitored by the PT. The INR is calculated from this and is based on the ratio of the patient's prothrombin time (PT) to a mean normal PT with

Table 28.2 Control tests for warfarin therapy. Target levels recommended by the British Society for Haematology.

Target INR	Clinical state
2.5 (2.0–3.0)	Treatment of DVT, pulmonary embolism, atrial fibrillation, recurrent DVT off warfarin; symptomatic inherited thrombophilia, cardiomyopathy, mural thrombus, cardioversion
3.0 (2.5–3.5)	Recurrent DVT while on warfarin, mechanical prosthetic heart valves, antiphospholipid syndrome (some cases)
DVT, deep venous thrombosis; INR, international normalized ratio.	

correction for the 'sensitivity' of the thromboplastin used. This is calibrated against a primary World Health Organization (WHO) standard thromboplastin. The indications and recommended ranges for INR with warfarin treatment are summarized in Table 28.2. Warfarin crosses the placenta and is teratogenic so is avoided in pregnancy.

Length of anticoagulation

It is usual to continue warfarin for 3–6 months for established DVT, PE and following xenograft heart valves. If the DVT was the first such episode, provoked by a transient major predisposing event such as surgery, and is also not extensive, then 3 months' therapy alone is usually given. However if the episode is unprovoked or associated with a minor risk factor such as flying, pregnancy or oral contraceptive, then 6 months' treatment is usually given. This is also so if the DVT is extensive or the D-dimers remain raised at 3 months.

Long-term therapy is given for recurrent venous thrombosis, for embolic complications of rheumatic heart disease or atrial fibrillation, and with prosthetic valves and arterial grafts and in selected patients with the APS. It is also given when a DVT is unprovoked and occurs at an unusual site, such as a mesenteric vein, or is massive. Long-term therapy is also usually recommended for severe deficiency of proteins C or S, or antithrombin, or for homozygous factor V Leiden or prothrombin gene mutations, or when compound heterozygotes for the two are detected.

Drug interactions

Approximately 97% of warfarin in the circulation is bound to albumin and only a small fraction of warfarin is free and can enter the liver parenchymal cells; it is this free fraction that is active. In the liver cells, warfarin is degraded in microsomes to an inactive water-soluble metabolite, which is conjugated and excreted in the bile and partially reabsorbed to be also excreted in urine. Drugs that affect the albumin binding or excretion of warfarin (or of other oral anticoagulants), or those that decrease the absorption of vitamin K, will interfere with the control of therapy (Table 28.3).

Table 28.3 Drugs and other factors that interfere with the control of coumarin (e.g. warfarin) therapy.

Potential of coumarin anticoagulants	Inhibition of coumarin anticoagulants
Drugs that increase the effect of coumarins	Drugs that depress the action of coumarins
<i>Reduced coumarin binding to serum albumin</i>	<i>Acceleration of hepatic microsomal degradation of coumarin</i>
Sulphonamides	Barbiturates
<i>Inhibition of hepatic microsomal degradation of coumarin</i>	Rifampicin
Amiodarone	Ribavirin
Diltiazem	<i>Enhanced synthesis of clotting factors</i>
Propanolol	Oral contraceptives
Antibiotics – ciprofloxacin, erythromycin, fluconazole	Hereditary resistance to oral anticoagulants
Alcohol	Pregnancy
Phenytoin	
Quinidine	
Allopurinol	
Tricyclic antidepressants	
Metronidazole	
Sulphonamides	
<i>Alteration of hepatic receptor site for drug</i>	
Thyroxine	
Quinidine	
<i>Decreased synthesis of vitamin K factors</i>	
High doses of salicylates	
Some cephalosporins, other antibiotics	
Liver disease	
Decreased synthesis of vitamin K factors	
Decreased absorption of vitamin K	
e.g. malabsorption, antibiotic therapy, laxatives	

N.B. Patients are also more likely to bleed if taking antiplatelet agents (e.g. NSAIDs, dipyridamole or aspirin); alcohol in large amounts enhances warfarin action.

Management of warfarin overdose

If the INR is in excess of 4.5 without bleeding, warfarin should be stopped for 1 or 2 days and the dosage adjusted according to the INR. The long half-life of warfarin (40 hours) delays the full impact of dose changes for 4–5 days. If the INR is very high (e.g. greater than 8) without bleeding, an oral dose of 0.5–2.5 mg vitamin K should be given. Mild bleeding usually only needs an INR assessment, drug withdrawal and subsequent dosage adjustment (Table 28.4). More serious bleeding

Table 28.4 Recommendations on the management of bleeding and excessive anticoagulation with warfarin.

INR 3.0–6.0 (target INR 2.5)	Reduce warfarin dose or stop
INR 4.0–6.0 (target INR 3.5)	Restart warfarin when INR <5.0
INR 6.0–8.0	Stop warfarin*
No bleeding or minor bleeding	Restart when INR <5.0
INR >8.0	Stop warfarin*
No bleeding or minor bleeding	Restart warfarin when INR <5.0
	If other risk factors for bleeding give 0.5–2.5 mg vitamin K orally
Major bleeding	Stop warfarin
	Give prothrombin complex concentrate (PCC) 50 units/kg, in preference
	FFP 15 mL/kg (if PCC not available)
	Give 5 mg vitamin K (i.v. or oral)

FFP, fresh frozen plasma; INR, international normalized ratio; i.v., intravenous.

* 1 mg vitamin K may be given orally to rapidly reduce the INR to the therapeutic range within 24 hours in all patients with an INR above the therapeutic range and no bleeding.

needs cessation of therapy, vitamin K therapy or the infusion of PCC (prothrombin complex concentrate). Fresh frozen plasma rarely reverses the anticoagulant effect of vitamin K antagonists as it is not usually possible to give sufficient volume. Vitamin K is the specific antidote; an oral or intravenous dose of 2.5 mg is usually effective. Higher doses result in resistance to further warfarin therapy for 2–3 weeks. Even larger doses, however, and for prolonged periods of weeks or months are needed for those who have accidentally ingested coumarin-like rodenticides ('superwarfarins') which are two logs more potent than warfarin and whose action lasts for weeks.

Management of surgery: bridging anticoagulation

For minor surgery (e.g. dental extraction) anticoagulation can be maintained and mouth rinses with tranexamic acid given. For major surgery requiring normal coagulation, warfarin should be stopped 5 days before surgery (to achieve an INR less than 1.5) and recommenced early after surgery when haemostasis has been achieved. As it takes several days for an anticoagulant effect to be re-established, thromboprophylaxis, for example with low-dose LMWH, should be given postoperatively to patients at risk of hospital-acquired thrombosis whilst they are not anticoagulated with warfarin.

In exceptional patients, such as those with metal mechanical heart valves, therapeutic-dose heparin (known as bridging therapy) should be given whilst they are not anticoagulated with warfarin. Bridging can be achieved with either unfractionated or LMWH.

Direct acting oral anticoagulants

New oral direct inhibitors of coagulation proteins offer a potential improvement over VKAs (Tables 28.5 and 28.6) by virtue of:

- Predictable dose responses;
- No need for routine monitoring;

Table 28.5 The oral direct-acting anticoagulants compared with warfarin.

	Xa inhibitor	Xa inhibitor	Xa inhibitor	IIa (thrombin) inhibitor	Vitamin K antagonist
	Rivaroxaban	Apixaban	Edoxaban	Dabigatran	Warfarin
Daily	bd	bd	od	bd	od
Half-life (h)	7–11	8–14	5–11	14–17	40
Renal excretion (%)	33	27	50	80	0
Peak effect	2–4 h	1–2 h	1–2 h	1–3 h	4–5 days
Laboratory test	PT or anti Xa*			Thrombin time* Anti IIa*	PT
Antidote	No specific antidote (prothrombin complex may be tried if necessary)				Vitamin K prothrombin complex Fresh frozen plasma
*Rarely needed PT, prothrombin time. Source: Adapted from Yeh C.H. <i>et al.</i> (2014) <i>Blood</i> 124: 1020–28.					

Table 28.6 The advantages and disadvantages of the new oral anticoagulant drugs compared to warfarin.

	Direct oral inhibitors	Warfarin
Advantages	Fixed dosing	Long experience
	No monitoring	Reversibility
	No food interactions	Safe in renal failure
	Few drug interactions	INR well validated
	Less major bleeding	Monitoring encourages compliance
	No requirement for parenteral heparins	Less gastrointestinal bleeding
Disadvantages	Compliance more important because short action	Food and drug interactions
	Renal function effects	Narrow therapeutic range
	Not in liver dysfunction	Needs monitoring and variable dose
	No reversal agents (various drugs in trials)	Slow onset (need for heparin at initiation and bridging)
	Less experience	Bleeding risk
	Bleeding risk	
	Not recommended for patients with target INR >3.0, pregnancy, children <18 years, antiphospholipid syndrome	

- Reduced need for dose adjustment;
- No food interactions;
- Few drug interactions.

However, because of their shorter action, compliance with daily dosing is more important. Renal failure affects dosage, particularly for dabigatran. A reversal agent is not easily available but trials are in progress of various compounds.

Drugs that are licensed include dabigatran (a IIa inhibitor), and rivaroxaban, apixaban and edoxaban (all Xa inhibitors) (Table 28.5). Monitoring of dosage is not needed but laboratory tests for the degree of anticoagulation are used if haemorrhage or thrombosis occurs and are also often used before an invasive procedure, if there is renal impairment, extremes of body weight, in children, to assess compliance or if other drugs which interact are being taken.

Dabigatran

Dabigatran is an oral thrombin inhibitor with a short half-life. It is used for prevention of cerebral and systemic embolism in patients with non-valvular atrial fibrillation (AF). It is given at a fixed dose without monitoring and without dose adjustment. The dose in patients with atrial fibrillation is 150 mg twice daily for those aged under 75 years. Food has no significant effect on absorption. Aspirin and non-steroidal anti-inflammatory drugs (NSAIDs) should be avoided if possible.

Ten per cent of patients suffer indigestion and these can be given a proton pump inhibitor (PPI). Dabigatran should not be prescribed when the creatinine clearance is less than 30 mL/min.

Rivaroxaban and apixaban

These are factor Xa inhibitors (Table 28.5). They are used for prevention of cerebral and systemic embolism in patients with non-valvular atrial fibrillation and treatment and prevention of DVT and PE. They are given at a fixed dose without monitoring and without dose adjustment. Rivaroxaban is given once daily at a dose of 20 mg and apixaban at 5 mg twice daily for treatment of DVT and prevention of stroke and embolism in AF. The starting doses are higher for about a week and may be reduced for long-term prophylaxis, for example for AF. There are no interactions with dietary products. They must not be prescribed for patients with an estimated glomerular filtration rate (eGFR) of less than 15 mL/min, and the dose should be reduced for patients with eGFR 15–30 mL/min. In rare cases the PT is used to assess their action. Reversal, if needed, can be achieved with prothrombin complex concentrate, but, in view of their short half-lives, it may be sufficient to control local haemorrhage until the anticoagulation effect has worn off. Edoxaban is another direct Xa inhibitor undergoing trials.

Post-thrombotic syndrome

Venous thrombi that persist can destroy venous valves and venous return is then impaired. This leads to venous hypertension which is responsible for fluid accumulation in the

extravascular space, with oedema, prominent superficial veins and, in the long-term, skin atrophy, melanin pigmentation and, in severe cases, skin ulceration. The leg feels itchy and heavy and painful particularly on standing.

Mechanical methods of prophylaxis of DVT and PE

Graduated compression stockings

These are often used postoperatively, post-partum and during long aeroplane flights to reduce the risk of DVT. After a DVT, they have been recommended for 1–2 years to reduce the risk of the post-thrombotic syndrome. Knee-high stockings have been used in the vast majority of cases, compression class II, and are worn except when the patient is recumbent. However recent evidence has questioned their benefit.

Intermittent compression devices

Intermittent pneumatic compression and mechanical foot pumps are used in high-risk patients in whom bleeding as a result of LMWH is likely. They reduce the risk of thrombosis in both medical and surgical patients.

Inferior vena cava filter

This can provide protection against pulmonary embolism when a DVT in the legs is diagnosed but anticoagulation is contraindicated (e.g. ongoing or very recent intracranial or gastrointestinal bleeding or where there is recurrent PE despite adequate anticoagulation).

Fibrinolytic agents

Two fibrinolytic agents, streptokinase and tissue plasminogen activator, are most frequently used to lyse fresh thrombi, although other agents are available. These drugs may be used systemically for patients with acute myocardial infarction, major PE with haemodynamic decompensation or iliofemoral thrombosis, and locally in patients with acute peripheral arterial occlusion.

Administration of thrombolytic agents has been simplified with standardized dosage regimens. The therapy is most effective in the first 6 hours after symptoms begin but is still of benefit up to 24 hours. Aspirin therapy is also given and the value of additional heparin therapy is under study. Laboratory tests for monitoring short-term thrombolytic therapy are unnecessary. However, certain clinical complications exclude the use of thrombolytic agents (Table 28.7).

Recombinant tissue plasminogen activator has a particularly high affinity for fibrin and this allows lysis of thrombi with less systemic activation of fibrinolysis.

Antiplatelet drugs

Antiplatelet agents are gaining an increasing role in clinical medicine. It is now clear that aspirin is valuable in the secondary prevention of vascular disease. Several other agents

Table 28.7 Contraindications to thrombolytic therapy.

Absolute contraindications	Relative contraindications
Active gastrointestinal bleeding	Traumatic cardiopulmonary resuscitation
Aortic dissection	Major surgery in the past 10 days
Head injury or cerebrovascular accident in the past 2 months	Past history of gastrointestinal bleeding
Neurosurgery in the past 2 months	Recent obstetric delivery
Intracranial aneurysm or neoplasm	Prior arterial puncture
Proliferative diabetic retinopathy	Prior organ biopsy
	Serious trauma
	Severe arterial hypertension (systolic pressure >200 mmHg, diastolic pressure >110 mmHg)
	Bleeding diathesis

Table 28.8 Suggested regimens for antiplatelet therapy in patients with an acute coronary syndrome and in those undergoing percutaneous coronary intervention (PCI).

Drug	Target/patient group	Duration
<i>Acute coronary syndrome</i>		
Aspirin	All	Life-long
Clopidogrel	All	9–12 months
Glycoprotein IIb/IIIa inhibitors:		
Abciximab	None	–
Eptifibatide	High risk	48–72 hours
Tirofiban	High risk	48–72 hours
<i>Patients undergoing PCI</i>		
Aspirin	All	Life-long
Clopidogrel	All	9–12 months
Abciximab	High risk	12 hours after PCI
Eptifibatide	High risk	18–24 hours after PCI
Tirofiban	None	–

are being used for different indications (Table 28.8). The sites of action of the antiplatelet drugs are illustrated in Fig. 28.5.

Aspirin Aspirin inhibits platelet cyclo-oxygenase irreversibly, thus reducing the production of platelet thromboxane A_2 . Its action lasts for the whole of the lifespan of the platelet.




Figure 28.5 Sites of action of antiplatelet drugs. Aspirin acetylates the enzyme cyclo-oxygenase irreversibly. Sulphinpyrazone inhibits cyclo-oxygenase reversibly. Dipyridamole inhibits phosphodiesterase, increases cyclic adenosine monophosphate (cAMP) levels and inhibits aggregation. Clopidogrel and GP blockers cause inhibition of adenosine uptake by red cells which allows adenosine accumulation in plasma and so stimulates platelet adenylatecyclase. Prostacyclin (epoprostenol) stimulates adenylatecyclase. The lipid-soluble β -blockers inhibit phospholipase. Calcium-channel antagonists block the influx of free calcium ions across the platelet membrane. Dextran coat the surface interfering with adhesion and aggregation. GP, glycoprotein.

Low-dose therapy (e.g. 75 mg/day) has a lesser risk of gastrointestinal bleeding and is most commonly used in patients who have a history of coronary artery or cerebrovascular disease. It is combined with clopidogrel for a year after coronary stenting or angioplasty. It may also be useful in preventing thrombosis in patients with thrombocytosis and has a mild action more generally in reducing the risk of DVT in subjects with normal platelet counts. Aspirin is contraindicated in patients with gastrointestinal or genitourinary bleeding, retinal bleeding, peptic ulcer, haemophilia or uncontrollable hypertension.

Clopidogrel This adenosine diphosphate (ADP) receptor antagonist inhibits ADP dependent platelet aggregation (Fig. 28.5). It is activated by cytochrome p450 in the liver and in some subjects this activation is reduced so the drug is ineffective. It is used at a dose of 75 mg daily for reduction of ischaemic events in patients with ischaemic stroke, myocardial infarction or peripheral vascular disease. It is used after coronary artery stenting or angioplasty and in patients requiring long-term antiplatelet therapy who are intolerant or allergic to aspirin. **Prasugrel** has the same action but is more effective with greater risk of bleeding.

Ticragelor inhibits platelet aggregation at a different site. It does not require hepatic activation and is an allosteric reversible inhibitor, which is an advantage if bleeding occurs or surgery is needed urgently. It needs to be given twice daily which reduces compliance.

Dipyridamole (Persantin®) This drug is a phosphodiesterase inhibitor thought to elevate cyclic adenosine monophosphate levels in circulating platelets, which decreases their sensitivity to activating stimuli. Dipyridamole has been shown to reduce thromboembolic complications in patients with prosthetic heart valves and to improve the results in coronary bypass operations.


Glycoprotein IIb/IIIa inhibitors: abciximab, eptifibatide, tirofiban These are monoclonal antibodies that inhibit the platelet GPIIb/IIIa receptor. They are used in conjunction with heparin, aspirin and clopidogrel for the prevention of ischaemic complications in high-risk patients undergoing percutaneous transluminal coronary angioplasty. They can be used once only.

SUMMARY

- Anticoagulant drugs are used to prevent or treat venous or arterial thrombosis. They are also used prophylactically systemically and to maintain patency of intravascular lines.
- Heparin can be given in the unfractionated form. Much more frequently low molecular weight heparin is given subcutaneously. It is used if anticoagulation is needed in pregnancy.
- Warfarin is the most frequently used oral anticoagulant, the dose usually aimed to raise the international normalized ratio (INR) to between 2.0 and 3.0. There are frequent drug interactions that affect the dose.
- New anticoagulants include the factor X_a inhibitors, rivaroxaban and apixaban, and the factor II (prothrombin) inhibitor, dabigatran. They have the advantage of fixed dosage, monitoring is not usually needed and there are fewer drug interactions.
- Thrombi, if fresh, may be dissolved by fibrinolytic agents (e.g. streptokinase) or recombinant tissue plasminogen activator.
- Antiplatelet drugs – aspirin, clopidogrel, prasugrel and dipyridamole – are used to treat arterial disorders.



Now visit www.wileyessential.com/haematology to test yourself on this chapter.


A vertical strip on the left side of the page showing a microscopic view of blood cells. It features several large, red, biconcave disc-shaped red blood cells, some smaller yellowish-white leukocytes, and a few small, dark purple platelets, all set against a dark blue background.

CHAPTER 29

Haematological changes in systemic diseases

Key topics

■ Anaemia of chronic disorders	322
■ Haematological problems in the elderly	322
■ Malignant diseases	322
■ Rheumatoid arthritis	323
■ Renal failure	325
■ Congestive heart failure	326
■ Liver disease	326
■ Hypothyroidism	327
■ Infections	327
■ Non-specific monitoring of systemic disease	330



Anaemia of chronic disorders (ACD)

Many of the anaemias seen in clinical practice occur in patients with systemic disorders and are the result of a number of contributing factors. The anaemia of chronic disorders (ACD) (also discussed on p. 37) is of central importance and occurs in patients with a variety of chronic inflammatory and malignant diseases (Table 29.1). Usually, both the erythrocyte sedimentation rate (ESR) and C-reactive protein (CRP) are raised. It may be complicated by additional haematological changes due to the disease. The serum iron and total iron binding capacity (transferrin) are both low, serum ferritin is normal or raised. The characteristic features and pathogenesis are described in Chapter 3.

The anaemia is corrected by the successful treatment of the underlying disease. It does not respond to iron therapy despite the low serum iron. Responses to recombinant erythropoietin therapy may be obtained (e.g. in rheumatoid arthritis or cancer). In many conditions the anaemia is complicated by anaemia from other causes (e.g. iron or folate deficiency, renal failure, bone marrow infiltration, hypersplenism or endocrine abnormality).

Haematological problems in the elderly

Anaemia

The WHO defines anaemia as a haemoglobin less than 130 g/L in men and less than 120 g/L in women. Using these criteria, the incidence of anaemia is substantial in the elderly, e.g. more than 25% in men and more than 20% in women over 85 years old. The incidence is higher in black people, increases with age and predicts for shorter survival. In the USA about 10% of subjects over 65 years old are anaemic and this is associated with increased hospitalization, disability and mortality.

The main causes are ACD, iron deficiency and renal disease, but about a third are unexplained, possibly some with myelodysplasia. Clones with molecular mutations characteristic of myelodysplasia are increasingly present with advancing age in the bone marrow, without morphological changes (see Chapter 16). The

elderly have reduced marrow reserve and develop more severe and prolonged anaemia (and neutropenia and thrombocytopenia) after chemotherapy than younger subjects.

Thrombosis

There is also an increased incidence of arterial and venous thrombosis with increasing age. This is partly due to increase in plasma levels of some of the clotting factors and reduced fibrinolysis as well as for arterial thrombosis to atheromatous plaques. Elderly subjects tend to be more sensitive than younger patients to anticoagulants and need careful monitoring to avoid haemorrhage.

Malignant diseases (other than primary bone marrow diseases)

Anaemia

Contributing factors include ACD, blood loss and iron deficiency, marrow infiltration (Fig. 29.1) often associated with a leucoerythroblastic blood film (p. 94), folate deficiency, haemolysis and marrow suppression from radiotherapy or chemotherapy (Table 29.2).

Microangiopathic haemolytic anaemia (p. 70) occurs with mucin-secreting adenocarcinoma (Fig. 29.2), particularly of the stomach, lung and breast. Less common forms of anaemia with malignant disease include autoimmune haemolytic anaemia with malignant lymphoma and rarely with other tumours; primary red cell aplasia with thymoma or lymphoma; and myelodysplastic syndromes secondary to chemotherapy. There is also an association of pernicious anaemia with carcinoma of the stomach.

The anaemia of malignant disease may respond partly to erythropoietin but care must be taken not to accelerate tumour growth (see Chapter 2). Folic acid should only be given if there is definite megaloblastic anaemia caused by the deficiency; it might 'feed' the tumour.

Polycythaemia

Secondary polycythaemia is occasionally associated with renal, hepatic, cerebellar and uterine tumours (see p. 172).

White cell changes

Leukaemoid reactions (p. 94) may occur with tumours showing widespread necrosis and inflammation. Hodgkin lymphoma is associated with a variety of white cell abnormalities, including eosinophilia, monocytosis and leucopenia. In non-Hodgkin lymphoma, malignant cells may circulate in the blood (see p. 274).

Platelet and blood coagulation abnormalities

Patients with malignant disease may show either thrombocytosis or thrombocytopenia. Disseminated tumours, particularly

Table 29.1 Causes of anaemia of chronic disorders.

Chronic inflammatory diseases

Infectious (e.g. pulmonary abscess, tuberculosis, osteomyelitis, pneumonia, bacterial endocarditis)

Non-infectious (e.g. rheumatoid arthritis, systemic lupus erythematosus and other connective tissue diseases, sarcoid, Crohn's disease, cirrhosis)

Malignant disease

(e.g. carcinoma, lymphoma, sarcoma, myeloma)




Figure 29.1 Metastatic carcinoma in bone marrow aspirates: (a) breast; (b) stomach; (c) colon; bone marrow trephine biopsies: (d) prostate; (e) stomach; (f) kidney.

mucin-secreting adenocarcinomas, are associated with disseminated intravascular coagulation (DIC; see p. 297) and generalized haemostatic failure. Activation of fibrinolysis occurs in some patients with carcinoma of the prostate. Occasional patients with malignant disease have spontaneous bruising or bleeding caused by an acquired inhibitor of one or other coagulation factor, most frequently factor VIII, or to a paraprotein interfering with platelet function.

Cancer patients have a high incidence (estimated at 15%) of venous thromboembolism. This is increased by surgery and some drugs, e.g. thalidomide. It is most common in ovarian, brain, pancreatic and colon cancers. It may be difficult to manage with oral anticoagulation because of bleeding,

interruptions with chemotherapy and thrombocytopenia, anorexia or vomiting. Liver disease and drug interactions can cause further complications so daily low molecular weight heparin injections may be preferable.

Rheumatoid arthritis (and other connective tissue disorders)

In patients with rheumatoid arthritis, the anaemia of chronic disorders is proportional to the severity of the disease. It is complicated in some patients by iron deficiency caused by gastrointestinal bleeding related to therapy with salicylates, non-steroidal anti-inflammatory agents or corticosteroids. Bleeding

Table 29.2 Haematological abnormalities in malignant disease.

Haematological abnormality	Tumour or treatment associated
<i>Pancytopenia</i>	
Marrow hypoplasia	Chemotherapy, radiotherapy
Myelodysplasia	Chemotherapy, radiotherapy
Leucoerythroblastic	Metastases in marrow
Megaloblastic	Folate deficiency
	B ₁₂ deficiency (carcinoma of stomach)
<i>Red cells</i>	
ACD	Most forms
Iron deficiency anaemia	Especially gastrointestinal, uterine
Pure red cell aplasia	Thymoma
Immune haemolytic anaemia	Lymphoma, ovary, other tumours
Microangiopathic haemolytic anaemia	Mucin-secreting carcinoma
Polycythaemia	Kidney, liver, cerebellum, uterus
<i>White cells</i>	
Neutrophil leucocytosis	Most forms
Leukaemoid reaction	Disseminated tumours, those with necrosis
Eosinophilia	Hodgkin disease, others
Monocytosis	Various tumours
<i>Platelets and coagulation</i>	
Thrombocytosis	Gastrointestinal tumours with bleeding, others
Disseminated intravascular coagulation	Mucin-secreting carcinoma, prostate
Activation of fibrinolysis	Prostate
Acquired inhibitors of coagulation	Most forms
Paraprotein interfering with platelet function	Lymphomas, myeloma
Tumour cell procoagulants – tissue factor and cancer procoagulant (density activates factor X)	Especially ovarian, pancreas, brain, colon




Figure 29.2 Peripheral blood film in metastatic mucin-secreting adenocarcinoma of the stomach showing red cell polychromasia and fragmentation and thrombocytopenia. The patient had disseminated intravascular coagulation.

into inflamed joints may also be a factor. Marrow hypoplasia may follow therapy with gold. In Felty's syndrome, splenomegaly is associated with neutropenia (Fig. 29.3). Anaemia and thrombocytopenia may also be present.



(a)



(b)

Figure 29.3 Felty's syndrome: (a) the typical deformities of rheumatoid arthritis of the hand and (b) splenomegaly.

In systemic lupus erythematosus (SLE) 50% of patients are leucopenic with reduced neutrophil and lymphocyte counts often associated with circulating immune complexes. Renal impairment and drug-induced gastrointestinal blood loss also contribute to the ACD. Autoimmune haemolytic anaemia (typically with immunoglobulin G (IgG) and the C3 component of complement on the surface of the red cells) occurs in 5% of patients and may be the presenting feature of the syndrome. There may be autoimmune thrombocytopenia in 5% of patients. The lupus anticoagulant is described on p. 307. This circulating anticardiolipin interferes with blood coagulation by altering the binding of coagulation factors to platelet phospholipid and predisposes to both arterial and venous thrombosis and recurrent abortions. Tests for anti-nuclear factor (ANF) and anti-DNA antibodies are usually positive.

Patients with temporal arteritis and polymyalgia rheumatica have a markedly elevated ESR, pronounced red cell rouleaux in the blood film and a polyclonal immunoglobulin response. These and other collagen vascular disorders are associated with ACD.

Renal failure

Anaemia

A normochromic anaemia is present in most patients with chronic renal failure. Generally, there is a 20 g/L fall in haemoglobin level for every 10 mmol/L rise in blood urea. There is impaired red cell production as a result of defective erythropoietin secretion (see Fig. 29.5). Variable shortening of red cell lifespan occurs and, in severe uraemia, the red cells show abnormalities including spicules (spurs) and 'burr' cells (Fig. 29.4).




Figure 29.4 Peripheral blood film in chronic renal failure showing red cell acanthocytosis and numerous 'burr' cells.




Figure 29.5 Liver disease: peripheral blood film showing: (a) macrocytosis and target cells; and (b) marked acanthocytosis and echinocytosis in Zieve's syndrome.

Increased red cell 2,3-DPG levels in response to the anaemia and hyperphosphataemia result in decreased oxygen affinity and a shift of the haemoglobin–oxygen dissociation curve to the right (see p. 17), which is augmented by uraemic acidosis. The patient's symptoms are therefore relatively mild for the degree of anaemia.

Other factors may complicate the anaemia of chronic renal failure (Table 29.3): ACD, iron deficiency from blood

Table 29.3 Haematological abnormalities in renal failure.

Anaemia

Reduced erythropoietin production
ACD
Iron deficiency
blood loss (e.g. dialysis, venesection, defective platelet function)
Folate deficiency
chronic haemodialysis without replacement therapy

Abnormal platelet function

Thrombocytopenia abnormal platelet function
immune complex-mediated (e.g. systemic lupus erythematosus, polyarteritis nodosa)
some cases of acute nephritis and following allograft
haemolytic uraemic syndrome and thrombotic
thrombocytopenic purpura

Thrombosis

Some cases of the nephrotic syndrome

Polycythaemia

In renal allograft recipients
Rarely in renal cell carcinoma, cysts, arterial disease

loss during dialysis or caused by bleeding because of defective platelet function, and folate deficiency in some chronic dialysis patients. Patients with polycystic kidneys usually have retained erythropoietin production and may have less severe anaemia for the degree of renal failure.

Treatment

Erythropoietin corrects the anaemia in patients on dialysis or in chronic renal failure, providing that iron and folate deficiency, and infections have been corrected. The dosage of erythropoietin usually required is 50–150 units/kg three times a week with a target haemoglobin of 120 g/L. Longer-acting preparations are increasingly used. Maintenance by 75 units/kg/week subcutaneously is typical. Complications of therapy have been initial transient flu-like symptoms, hypertension, clotting of the dialysis lines and, rarely, fits. A poor response to erythropoietin suggests iron or folate deficiency, infection or hyperparathyroidism. Intravenous iron is often needed to correct iron deficiency shown by low serum ferritin and percentage saturation of total iron-binding capacity (TIBC), and increased percentage of hypochromic red cells in the blood.

Platelet and coagulation abnormalities

A bleeding tendency with purpura, gastrointestinal or uterine bleeding occurs in 30–50% of patients with chronic renal failure and is marked in patients with acute renal failure. The bleeding may be out of proportion to the degree of thrombocytopenia and has been associated with abnormal platelet or vascular function, which can be reversed by dialysis. Correction of the anaemia with erythropoietin also improves the bleeding tendency. Immune complex-mediated thrombocytopenia occurs in some patients with acute nephritis, SLE and polyarteritis nodosa and also following renal allografts. Renal allografts may also lead to polycythaemia in 10–15% of patients.

The haemolytic uraemic syndrome and thrombotic thrombocytopenic purpura are discussed on p. 285. Patients with the nephrotic syndrome have an increased risk of venous thrombosis.

Congestive heart failure

Anaemia is present in 30–50% of patients with congestive heart failure due to chronic kidney disease, haemodilution, diabetes and release of cytokines increasing hepcidin synthesis (reducing iron absorption and recycling of iron from macrophages) and reducing erythropoietin secretion. Treatment with oral or intravenous iron may improve left ventricular ejection fraction, lower CRP and brain natriuretic peptide (BNP) levels, reduce fatigue, and increase exercise capacity and quality of life, even in those without iron deficiency.

Table 29.4 Haematological abnormalities in liver disease.

Liver failure ± obstructive jaundice ± portal hypertension

Refractory anaemia – usually mildly macrocytic, often with target cells; may be associated with:
Blood loss and iron deficiency
Alcohol (± ring sideroblastic change)
Folate deficiency
Haemolysis (e.g. Zieve's syndrome, Wilson's disease, immune, hypersplenism from portal hypertension)

Bleeding tendency
Deficiency of vitamin K-dependent factors; also of factor V and fibrinogen
Thrombocytopenia, immune platelet function defects
Functional abnormalities of fibrinogen
Increased fibrinolysis
Portal hypertension – haemorrhage from varices

Viral hepatitis Aplastic anaemia

Tumours
Polycythaemia
Neutrophil leucocytosis and leukaemoid reactions

Liver disease

The haematological abnormalities in liver disease are listed in Table 29.4. Chronic liver disease is associated with anaemia that is mildly macrocytic and often accompanied by target cells, mainly as a result of increased cholesterol in the membrane (Fig. 29.5a). Contributing factors to the anaemia may include blood loss (e.g. bleeding varices) with iron deficiency, dietary folate deficiency and direct suppression of haemopoiesis by alcohol.

Haemolytic anaemia may occur in patients with alcohol intoxication (Zieve's syndrome) (Fig. 29.5b) and in Wilson's disease (caused by copper oxidation of red cell membranes), and autoimmune haemolytic anaemia is found in some patients with chronic immune hepatitis. Haemolysis may also occur in end-stage liver disease because of abnormal red cell membranes resulting from lipid changes. Viral hepatitis is associated with aplastic anaemia (see Chapter 22).

The acquired coagulation abnormalities associated with liver disease are described on p. 297. There are deficiencies of vitamin K-dependent factors (II, VII, IX and X) and, in severe disease, of factor V and fibrinogen. Thrombocytopenia may occur from hypersplenism or from immune complex-mediated platelet destruction. Abnormalities of platelet function may also be present. Dysfibrinogenaemia with abnormal fibrin

polymerization may occur as a result of excess sialic acid in the fibrinogen molecules. A consumptive coagulopathy may be superimposed. These haemostatic defects may contribute to major blood loss from bleeding varices caused by portal hypertension.

Hypothyroidism

A moderate anaemia is usual and may be caused by lack of thyroxine. T_3 and T_4 potentiate the action of erythropoietin. There is also a reduced oxygen need and thus reduced erythropoietin secretion. The anaemia is often macrocytic and the mean corpuscular volume (MCV) falls with thyroxine therapy. Autoimmune thyroid disease, especially myxoedema or Hashimoto's disease, is associated with pernicious anaemia. Iron deficiency may also be present, particularly in women with menorrhagia.

Infections

Haematological abnormality is usually present in patients with infections of all types (Table 29.5). The effect of inflammation as a prothrombotic stimulus is also discussed on p. 298.

Bacterial infections

Acute bacterial infections are the most common cause of neutrophil leucocytosis. Toxic granulation, Döhle bodies and metamyelocytes may be present in the blood (see Chapter 8). Leukaemoid reactions with a white cell count $\alpha\beta\omega\epsilon$ $50 \times 10^9/L$ and granulocyte precursors in the blood may occur in severe infections, particularly in infants and young children. Mild anaemia is common if the infection is prolonged. Severe haemolytic anaemia occurs in bacterial septicaemias, particularly those caused by Gram-negative organisms, where there is usually

Table 29.5 Blood abnormalities associated with infections.

Haematological abnormality	Infection associated
<i>Anaemia</i>	
Anaemia of chronic disorders	Chronic infections especially tuberculosis
Aplastic anaemia	Viral hepatitis
Transient red cell aplasia	Human parvovirus
Marrow fibrosis	Tuberculosis
Immune haemolytic anaemia	Infectious mononucleosis, <i>Mycoplasma pneumoniae</i>
Direct red cell damage or microangiopathic	Bacterial septicaemia (associated DIC), <i>Clostridium perfringens</i> , malaria, bartonellosis
Hypersplenism	Viruses – haemolytic uraemic syndrome and TTP Chronic malaria, tropical splenomegaly syndrome, leishmaniasis, schistosomiasis
<i>White cell changes</i>	
Neutrophil leucocytosis	Acute bacterial infections
Leukaemoid reactions	Severe bacterial infections particularly in infants
Eosinophilia	Tuberculosis
Monocytosis	Parasitic diseases (e.g. hookworm, filariasis, schistosomiasis, trichinosis)
Neutropenia	Recovery from acute infections
Lymphocytosis	Chronic bacterial infections: tuberculosis, brucellosis, bacterial endocarditis, typhoid
Lymphopenia	Viral infections – HIV, hepatitis, influenza Fulminant bacterial infections (e.g. typhoid, miliary tuberculosis) Infectious mononucleosis, toxoplasmosis, cytomegalovirus, rubella, viral hepatitis, pertussis, tuberculosis, brucellosis HIV infection <i>Legionella pneumonophila</i>
<i>Thrombocytopenia</i>	
Megakaryocytic depression, immune complex-mediated and direct interaction with platelets	Acute viral infections particularly in children (e.g. measles, varicella, rubella, malaria, severe bacterial infection)
Prothrombotic state	All with prolonged inflammation

DIC, disseminated intravascular coagulation; HIV, human immunodeficiency virus; TTP, thrombotic thrombocytopenic purpura.

associated DIC (see p. 297). DIC dominates the clinical picture in certain infections (e.g. bacterial meningitis). The acute phase response to infections is accompanied by a rise in coagulation factors and a fall in natural anticoagulants.

Clostridium perfringens organisms produce an α toxin, a lecithinase acting directly on the circulating red cells (Fig. 29.6). Haemolysis in bartonellosis (Oroya fever) is caused by direct red cell infection. *Mycoplasma pneumoniae* infections are associated with autoimmune haemolytic anaemia of the 'cold' type (see p. 69). With severe acute bacterial infections there may be thrombocytopenia. Chronic bacterial infections are associated with the ACD. In tuberculosis, additional factors in the pathogenesis of anaemia include marrow replacement and fibrosis associated with miliary disease and reactions to antituberculous therapy (e.g. isoniazid is a pyridoxine antagonist and may cause sideroblastic anaemia). Disseminated tuberculosis is associated with leukaemoid reactions and patients with involvement of bone marrow may show leucoerythroblastic changes in the peripheral blood film (see Fig. 8.9).

Viral infections

Acute viral diseases are often associated with a mild anaemia. An immune haemolytic anaemia with an anti-i autoantibody is associated with infectious mononucleosis (see p. 69). Viral infections, as well as syphilis, have been associated with paroxysmal cold haemoglobinuria (see p. 69). Viruses have also been linked to the pathogenesis of the haemolytic uraemic syndrome, thrombotic thrombocytopenic purpura (see Chapter 24) and the haemophagocytic syndrome (see p. 69). Aplastic anaemia

may occur with viral A or more usually non-A, non-B, non-C hepatitis. Transient red cell aplasia is associated with human parvovirus infection and this may result in severe anaemia (see Chapter 6).

Acute thrombocytopenia is frequent in rubella, morbilli and varicella infections. Rubella, cytomegalovirus (CMV) and other viral infections may cause a reactive lymphocytosis similar to that found in infectious mononucleosis. CMV infections in infants are associated with massive hepatosplenomegaly. In bone marrow transplant recipients or other immunosuppressed patients, CMV infections may cause pancytopenia as well as other severe disorders (see Chapter 23).

Human immunodeficiency virus (HIV) infection

This is associated with a wide range of haematological changes. These are caused by marrow defects and immune cytopenias directly resulting from HIV infection, the effects of opportunistic infections or lymphoma and the side-effects of drugs used to treat HIV itself or drugs for the complicating infection or lymphoma.

Anaemia is common and more severe as the disease progresses. It is usually multifactorial in origin: ACD, marrow dysplasia and drug therapy, especially zidovudine. Serum vitamin B₁₂ is often low, most likely because of intestinal malabsorption, but the anaemia does not respond to vitamin B₁₂ therapy. Blood transfusions or recombinant erythropoietin injections are needed.

Thrombocytopenia and neutropenia may be immune or secondary to marrow dysfunction. The marrow may be hypercellular with prominent plasma cells and lymphocytes, normocellular, hypocellular or fibrotic. Dysplastic features are common, with ineffective thrombopoiesis or granulocyte formation accounting at least in part for the cytopenia. The myelodysplasia does not show the chromosome abnormalities found in classic myelodysplasia and does not appear to be pre-leukaemic. Thrombocytopenia is treated if necessary by corticosteroids, high-dose gammaglobulin infusions or by other therapies for immune thrombocytopenia (see p. 283), or by antiretroviral therapy.

Increased plasma cells in the marrow and polyclonal increase in immunoglobulins are frequent. A paraprotein is present in 5–10% of cases, but appears benign. Non-Hodgkin lymphoma, in over 90% high grade, both systemically and in the central nervous system, occurs in HIV-infected individuals with over 100 times the frequency expected in the general population. Diffuse large B-cell lymphoma is the most common, with 20% confined to the central nervous system. A substantial minority are Burkitt lymphoma. Hodgkin lymphoma, usually of poor prognosis type, is also increased in frequency. EBV infection appears to underlie this as well as Burkitt lymphoma.

Treatment of lymphomas in the setting of HIV is with standard regimens. Continuation of the necessary antiretroviral therapy exaggerates the tendency to cytopenia induced by chemotherapy, so prophylaxis against opportunistic infections is important. The bone marrow may indeed reveal the presence of opportunistic infection (Fig. 29.7).




Figure 29.6 Peripheral blood film in a patient with haemolytic anaemia in clostridial septicaemia showing red cell contraction and spherocytosis.




Figure 29.7 Human immunodeficiency virus (HIV) infection: bone marrow trephine biopsy. **(a)** Granuloma showing positivity with Ziehl-Nielsen stain. **(b)** Higher power shows large numbers of acid-fast bacilli.

Malaria

Some degree of haemolysis is seen in all types of malarial infection (see Chapter 6). The most severe abnormalities are found in *Plasmodium falciparum* infections (Fig. 29.8). In the worst cases, DIC occurs and intravascular haemolysis is marked with haemoglobinuria. This may be associated with quinine therapy ('blackwater fever'). Thrombocytopenia is commonly found in acute malaria. Patients with chronic malaria have an anaemia of chronic disorders; hypersplenism may contribute to the anaemia and result in moderate thrombocytopenia and neutropenia. Tropical splenomegaly is probably a chronic immune reaction to malaria (see Chapter 10). Dyserythropoiesis in the marrow, folate deficiency and protein-calorie malnutrition may contribute to anaemia.

Toxoplasmosis

Toxoplasmosis in children and adults is associated with lymphadenopathy and large numbers of atypical lymphocytes in




Figure 29.8 Malaria: peripheral blood in severe *Plasmodium falciparum* infection showing: **(a)** many ring forms and a meront; and at higher magnification: **(b)** a meront; and **(c)** a gametocyte.

the blood. Congenital disease may cause a syndrome resembling hydrops fetalis with severe anaemia, an hydropic infant with gross hepatosplenomegaly and thrombocytopenia.

Kala-azar (visceral leishmaniasis)

The visceral form of leishmaniasis is associated with pancytopenia, hepatosplenomegaly and lymphadenopathy. Bone marrow or splenic aspirates may show large numbers of parasitized macrophages (Fig. 29.9).

Other parasitic diseases

Chronic schistosomiasis (bilharzia) affects over 200 million people worldwide. It is one of the most frequent causes of iron deficiency due to bleeding from the bowel or bladder. Hypersplenism follows splenic enlargement associated with portal hypertension due to liver infestation. In the acute phase of both African and South American trypanosomiasis, organisms are found in the peripheral blood (Fig. 29.10). Microfilariae of bancroftian filariasis and loiasis are also detected during blood film examination (Fig. 29.11). In many parasitic diseases there is eosinophilia.




Figure 29.9 Kala-azar: bone marrow aspirates showing macrophages containing Leishman–Donovan bodies.



Figure 29.10 African trypanosomiasis: blood film showing *Trypanosoma brucei*.

Osteopetrosis

Osteopetrosis is a rare genetic disorder in which there is an increase in bone mass with skeletal abnormality and bone marrow failure. It is due to defects in a variety of genes affecting osteoclast function. The bones are brittle, and there is extramedullary haemopoiesis with enlargement of the liver and spleen. The only cure is by stem cell transplantation.

Non-specific monitoring of systemic disease

The inflammatory response to tissue injury includes changes in plasma concentrations of proteins known as acute phase proteins. These include fibrinogen, other clotting factors, complement components, CRP (p. 332), haptoglobin, serum amyloid A (SAA) protein, ferritin and others. The rise in these liver-derived proteins is part of a wider response, which includes fever, leucocytosis and increased immune reactivity. The acute phase response is mediated by cytokines (e.g. IL-1 and TNF; see Fig. 8.4) released from macrophages and other cells. Quantitative measurements of acute phase proteins are valuable indicators of the presence and extent of inflammation and of its response to treatment. When short-term (less than 24 hours) changes in the inflammatory response are expected, CRP is the test of choice (Table 29.6). Long-term changes in the acute phase proteins are monitored by either the ESR or plasma viscosity. These tests are influenced by plasma proteins, which are either slowly responding acute phase reactants (e.g. fibrinogen) or are not acute phase proteins (e.g. immunoglobulins).

Erythrocyte sedimentation rate

This commonly used but non-specific test measures the speed of sedimentation of red cells in plasma over a period of 1 hour. The speed is mainly dependent on the plasma concentration




Figure 29.11 Peripheral blood films showing microfilariae of: **(a)** *Wuchereria bancrofti*; and **(b)** *Loa loa*.

of large proteins (e.g. fibrinogen and immunoglobulins). The normal range in men is 1–5 mm/hour and in women 5–15 mm/hour, but there is a progressive increase with age. The ESR is raised in a wide variety of systemic inflammatory and neoplastic diseases and in pregnancy. It is useful for

diagnosing and monitoring temporal arteritis and polymyalgia rheumatica and for monitoring patients with Hodgkin lymphoma. High values (>100 mm/hour) have a 90% predictive value for serious disease, including infections, collagen vascular disease or malignancy (particularly myeloma). A raised ESR is

Table 29.6 Advantages and disadvantages of the tests used to monitor the acute phase response.

Advantages	Disadvantages
<i>CRP*</i> Specific test of acute phase protein Fast response (6 hours) to change in disease activity High sensitivity – owing to large incremental change Can be measured on stored serum Small sample volumes Automated analysis	More than one protein required to measure acute (CRP) and chronic inflammation Costly when assayed in small numbers Sophisticated equipment and antisera required
<i>ESR and plasma viscosity</i> Useful in chronic disease ESR inexpensive, easy, no electrical power required Plasma viscosity—result obtained quickly (15 min) Plasma viscosity not affected by anaemia	Not sensitive to acute changes (<24 hours) Not specific for acute phase response Slow to change with alteration in disease activity and insensitive to small changes in activity Fresh samples (<2 hours) required for ESR
ESR, erythrocyte sedimentation rate. * C-reactive protein (CRP) is normally present in low concentrations (<5 mg/L). Levels are not influenced by anaemia, pregnancy or heart failure. During severe acute infection the plasma concentration may rise 100-fold.	

associated with marked rouleaux formation of red cells in the peripheral blood film (see Fig. 21.7). Changes in the ESR can be used to monitor the response to therapy.

Lower than expected readings occur in polycythaemia vera because of the high red cell concentration. Higher than expected values may occur in severe anaemia because of the low red cell concentration.

Plasma viscosity

Plasma viscosity is affected by the concentration of plasma proteins of large molecular size, especially those with pronounced axial asymmetry: fibrinogen and some immunoglobulins. Normal values at room temperature are usually in the range of 1.50–1.70 mPa/s. Lower levels are found in neonates because of lower levels of proteins, particularly fibrinogen. Viscosity increases only slightly in the elderly as fibrinogen increases. There is no difference in values between men and women. Other advantages over

the ESR test include independence from the effects of anaemia and results that are available within 15 minutes.

C-reactive protein

Phylogenetically CRP is a crude 'early' immunoglobulin which initiates the inflammatory reaction. CRP–antigen complexes can substitute for antibody in the fixation of C1q and trigger the complement cascade, initiating the inflammatory response to antigens or tissue damage. Subsequent binding of C3b on the surface of microorganisms opsonizes them for phagocytosis.

After tissue injury, an increase in CRP, SAA protein and other acute phase reactants may be detected within 6–10 hours. Immunoassays of CRP are widely used for early detection of acute inflammation or tissue injury and for the monitoring of remission (e.g. response of infection to an antibiotic).


Table 29.6 lists the advantages and disadvantages of the tests used to assess the acute phase response.

SUMMARY

- Chronic inflammation or malignant disorders cause anaemia with low serum iron and iron-binding capacity, normal or raised serum ferritin, an inadequate response to erythropoietin and reduced red cell lifespan. The degree of anaemia relates to the severity of the underlying disease. It does not respond to iron therapy.
- This anaemia may be complicated by other causes of anaemia, e.g. iron or folate deficiencies, renal failure, bone marrow infiltration, haemolysis, hypersplenism.
- Polycythaemia is a much less frequent complication of systemic diseases, e.g. renal.
- White cell changes are also frequent in systemic diseases. These include neutrophil leucocytosis, especially in bacterial infections, leucoerythroblastic or leukaemoid reactions, and, in viral and connective tissue diseases, neutropenia.
- Eosinophilia occurs with certain infections, particularly parasitic and allergic disease.
- Monocytosis is associated with chronic bacterial infections (e.g. tuberculosis, brucellosis).
- Lymphocytosis is a feature of viral infections and some bacterial infections, e.g. *Bordetella pertussis*.
- Platelets may be increased or low in malignant, infectious and other systemic diseases. Disseminated intravascular coagulation is a major cause of thrombocytopenia and fall in coagulation factors.
- C-reactive protein can be used for non-specific monitoring of systemic disease in the short term (hours or days) and erythrocyte sedimentation rate (or plasma viscosity) over weeks or months.



Now visit www.wileyessential.com/haematology to test yourself on this chapter.




CHAPTER 30

Blood transfusion

Key topics

■ Blood donor	334
■ Red cell antigens and blood group antibodies	334
■ Hazards of allogeneic blood transfusion	338
■ Techniques in blood group serology	339
■ Cross-matching and pre-transfusion tests	340
■ Complications of blood transfusion	340
■ Reduction of blood product use	342
■ Blood components	343
■ Preparations from human plasma	344
■ Acute blood loss	345



Blood transfusion consists of the 'safe' transfer of blood components (Fig. 30.1) from a donor to a recipient. In the UK all blood banks are inspected by the Medicines and Healthcare Regulatory Agency (MHRA). All adverse events involving blood products must be reported to the Serious Adverse Blood Reactions and Events (SABREA) scheme. Errors in the transfusion process including SABRE reportable events are reported to the Serious Hazards of Transfusion (SHOT) scheme.

Blood donor

In most countries blood donors contribute on a voluntary basis and this is generally preferable in terms of product safety. The measures to protect donors and for donor selection are listed in Table 30.1.

Red cell antigens and blood group antibodies

The clinical significance of blood groups in blood transfusion is that individuals who lack a particular blood group antigen may produce antibodies reacting with that antigen which may lead to a transfusion reaction. Approximately 400 red blood cell group antigens have been described. The different blood group antigens vary greatly in their clinical significance with the ABO and Rh (formerly Rhesus) groups being the most important. Some other systems are listed in Table 30.2.

Blood group antibodies

The ABO blood group antigens are unusual in that naturally occurring antibodies occur in the plasma of subjects who lack the corresponding antigen, even if they have not been transfused or been pregnant (Table 30.3). The most important of these natural antibodies are anti-A and anti-B. They are usually immunoglobulin M (IgM), and react optimally at cold temperatures (4°C) so, although reactive at 37°C, are called cold antibodies.

Immune antibodies develop in response to the introduction – by transfusion or by transplacental passage during pregnancy – of red cells possessing antigens that the subject lacks. These antibodies are commonly IgG, although some IgM antibodies may also develop, usually in the early phase of an immune response. Immune antibodies react optimally at 37°C (warm antibodies). Only IgG antibodies are capable of transplacental passage from mother to fetus and the most important immune antibody is the Rh antibody, anti-D.

ABO system

The protein that defines the ABO antigens is a glycosyl-transferase that is encoded from a single gene for which there are three major alleles, A, B and O. The A and B alleles catalyse addition of different carbohydrate residues (*N*-acetyl galactosamine for group A and d-galactose for group B) to a basic antigenic glycoprotein or glycolipid




Figure 30.1 The preparation of blood components from whole blood. FFP, fresh frozen plasma; SAGM, saline-adenine-glucose-mannitol. * Cryoprecipitate is mainly a source of fibrinogen. Cryosupernatant is used for plasma exchange in thrombotic thrombocytopenic purpura. Leucodepletion – see text.

Table 30.1 Measures used to protect the donor and for donor selection.*Donor selection*

Age 17–70 years (maximum 65 at first donation)

Weight above 50 kg (7 st 12 lb)

Haemoglobin >134 g/L for men, >120 g/L for women

Minimum donation interval of 12 weeks (16 weeks advised) and three donations per year maximum

Apheresis for platelets or plasma up to 24 times in 12 months

Pregnant and lactating women excluded because of high iron requirements; donation deferred for 9 months post pregnancy

Exclusion of those with:

Known cardiovascular disease, including hypertension

Significant respiratory disorders

Epilepsy and other CNS disorders

Gastrointestinal disorders with impaired absorption

Previous blood transfusions in the UK

Intravenous drug users

Insulin-dependent diabetes

Chronic renal disease

Cancer

Ongoing medical investigation or clinical trials

Exclusion of any donor returning to occupations such as driving bus, plane or train, heavy machine or crane operator, mining, scaffolding, etc. because delayed faint would be dangerous

Defer for 12 months after body piercing or tattoo, paid sex or homosexual sex, after acupuncture

Defer for 2 months after live vaccinations such as measles, mumps

Defer if travel history suggests risk of infection

CNS, central nervous system.

with a terminal sugar l-fucose on the red cell, known as the H substance (Fig. 30.2). The O allele is non-functional and so does not modify the H substance. Although there are six common genotypes, the absence of a specific anti-H prevents the serological recognition of more than four phenotypes (Table 30.4). The A allele actually itself has two variants, A_1 and A_2 , which further complicates the issue but these are of minor clinical significance. A_2 cells react more weakly than A_1 cells with anti-A, and patients who are A_2B can be wrongly grouped as B.

Table 30.2 Donor testing in England and Wales.

1 Blood group, Rh status (D,C,E,c,e), K

2 Screen for red cell alloantibodies

3 *Microbiological tests*

Human immunodeficiency virus (HIV) 1 and 2; antibody and RNA

Hepatitis B virus (HBV) – antibody and RNA

Hepatitis C virus (HCV) – antibody and RNA

Human T-cell leukaemia viruses (HTLV) – antibody

Cytomegalovirus (CMV) – antibody, for immunosuppressed recipients

Malaria – antibody screening of potentially exposed donors

Chagas' disease – antibody screening of potentially exposed donors

Bacteria – all donations tested for antibody to syphilis

N.B. At the current time there is no reliable test for detecting prions in blood products.

The A, B and H antigens are present on most body cells including white cells and platelets. In the 80% of the population who possess secretor genes, these antigens are also found in soluble form in secretions and body fluids (e.g. plasma, saliva, semen and sweat).

Naturally occurring antibodies to A and/or B antigen (usually IgM but occasionally IgG) are found in the plasma of subjects whose red cells lack the corresponding antigen (Table 30.4; Fig. 30.3).

Rh system

The Rh blood group locus is composed of two related structural genes, *RhD* and *RhCE*, which encode the membrane proteins that carry the D, Cc and Ee antigens. The *RhD* gene may be either present or absent, giving the Rh D+ or Rh D– phenotype, respectively. Alternative RNA splicing from the *RhCE* gene generates two proteins, which encode the C or c, and the E or e antigens (Fig. 30.4). A shortened nomenclature for Rh phenotype is commonly used (Table 30.5).

Rh antibodies rarely occur naturally and are therefore immune antibodies that result from previous transfusion or pregnancy. Anti-D is responsible for most of the clinical problems associated with the system and a simple subdivision of subjects into Rh D+ and Rh D– using anti-D antibody is sufficient for routine clinical purposes. Anti-C, anti-c, anti-E and anti-e are occasionally seen and may cause both transfusion reactions and haemolytic disease of the newborn. Anti-d does not exist. Rh haemolytic disease of the newborn is described in Chapter 31.

Table 30.3 Clinically important blood group systems.

Systems	Frequency of antibodies	Cause of haemolytic transfusion reaction	Cause of haemolytic disease of newborn
ABO	Almost universal	Yes (common)	Yes (usually mild)
Rh	Common	Yes (common)	Yes
Kell	Occasional	Yes (occasional)	Anaemia not haemolysis
Duffy	Occasional	Yes (occasional)	Yes (occasional)
Kidd	Occasional	Yes (occasional)	Yes (occasional)
Lutheran	Rare	Yes (rare)	No
Lewis	Occasional	Yes (rare)	No
P	Occasional	Yes (rare)	Yes (rare)
MN	Rare	Yes (rare)	Yes (rare)
Li	Rare	Unlikely	No

Table 30.4 The ABO blood group system.

Phenotype	Genotype	Antigens	Naturally occurring antibodies	Frequency (UK) (%)
O	OO	O	Anti-A, anti-B	46
A	AA or AO	A	Anti-B	42
B	BB or BO	B	Anti-A	9
AB	AB	AB	None	3




Figure 30.2 Structure of ABO blood group antigens. Each consists of a chain of sugars attached to lipids or proteins which are an integral part of the cell membrane. The H antigen of the O blood group has a terminal fucose (fuc). The A antigen has an additional *N*-acetyl galactosamine (galnac), and the B antigen has an additional galactose (gal). glu, glucose.




Figure 30.3 (a) The ABO grouping in a group A patient. The red cells suspended in saline agglutinate in the presence of anti-A or anti-A + B (serum from a group O patient). (b) Routine grouping in a 96-well microplate. Positive reactions show as sharp agglutinates; in negative reactions the cells are dispersed. Rows 1–3, patient cells against antisera; rows 4–6, patient sera against known cells; rows 7–8, anti-D against patient cells.




Figure 30.4 Molecular genetics of the Rh blood group. The locus consists of two closely linked genes, *RhD* and *RhCcEe*. The *RhD* gene codes for a single protein which contains the RhD antigen whereas *RhCcEe* mRNA undergoes alternative splicing to three transcripts. One of these encodes the E or e antigen whereas the other two (only one is shown) contain the C or c epitope. A polymorphism at position 226 of the *RhCcEe* gene determines the Ee antigen status whereas the C or c antigens are determined by a four amino acid allelic difference. Some individuals do not have an *RhD* gene and are therefore RhD–.

Other blood group systems

Other blood group systems are less frequently of clinical importance. Although naturally occurring antibodies of the P, Lewis and MN system are not uncommon, they usually only react at low temperatures and hence are of no clinical consequence. Immune antibodies against antigens of these systems are detected infrequently. Many of the antigens are of low

antigenicity and others (e.g. Kell), although comparatively immunogenic, are of relatively low frequency and therefore provide few opportunities for isoimmunization except in multiply transfused patients.

Molecular techniques, now available for blood group testing and extended red cell genotyping, are likely to be automated and become increasingly used.

Table 30.5 The most common Rh genotypes in the UK population.

CDE nomenclature	Short symbol	Frequency in white people (%)	Rh D status
cde/cde	Rr	15	Negative
CDe/cde	R ₁ r	31	Positive
CDe/CDe	R ₁ R ₁	16	Positive
cDE/cde	R ₂ r	13	Positive
CDe/cDE	R ₁ R ₂	13	Positive
cDE/cDE	R ₂ R ₂	3	Positive
Other genotypes		9	Positive (almost all)

Hazards of allogeneic blood transfusion

A large number of measures are taken to protect the recipient (Table 30.6).

Infection

Donor selection and testing of all donations are designed to prevent transmission of diseases (Tables 30.1 and 30.2). The main risk is from viruses that have long incubation periods, especially where these are asymptomatic. Recent viral infections can be transmitted in the pre-symptomatic viraemic phase, if blood has been collected during that short period (Table 30.7).

Hepatitis

Donors with a history of hepatitis are deferred for 12 months. If there is a history of jaundice, they can be accepted if markers for HBV and HCV are negative.

Human immunodeficiency virus (HIV)

This can be transmitted by cells or plasma. Male homosexuals, bisexuals, intravenous drug users and prostitutes are excluded, as are their sexual partners and partners of haemophiliacs. Inhabitants of large areas of sub-Saharan Africa and South-East Asia where HIV infection is particularly common are also excluded. Very rarely transmission may occur when the donor is incubating the infection but is not yet positive for the antigen–antibody test used ('window period transmission').

Human T-cell leukaemia viruses

Human T-cell leukaemia virus type I (HTLV-I) is associated with adult T-cell leukaemia or tropical spastic paraparesis. Human T-cell leukaemia virus type II (HTLV II) has no known association with any clinical condition. Screening for both is mandatory in the UK despite the low prevalence of approximately 1 in 50 000 untested donors.

Cytomegalovirus

Post-infusion cytomegalovirus (CMV) infection is usually sub-clinical but may cause an infectious mononucleosis syndrome. Immunosuppressed individuals are at risk of pneumonitis and a potentially fatal disease. These are premature babies (less than 1500 g), stem cell and other organ transplant recipients, patients who have received alemtuzumab (anti-CD52) and pregnant women (where the fetus is at risk). For such recipients, CMV-negative blood or blood components must be given, if they are CMV negative.

Other infections

Syphilis is more likely to be transmitted by platelets (stored at room temperature) than blood (stored at 4°C). However, all donations are tested. Malarial parasites are viable in blood stored at 4°C, so in endemic areas all recipients are given antimalarial drugs. In non-endemic areas, donors are carefully vetted for travel to tropical areas and in some centres

Table 30.6 Measures to protect recipient.

Donor selection (see Table 30.1)
Donor deferral/exclusion (see Table 30.1)
Stringent arm cleaning
Microbiological testing of donations (Table 30.2)
Immunohaematological testing of donations
Discarding the first 20–30 mL of blood collected
Leucodepletion of cellular products
Post-collection viral inactivation of FFP
Monitoring and testing for bacterial contamination
Pathogen inactivation of cellular components
Safest possible sources of donor for plasma products
FFP, fresh frozen plasma.

Table 30.7 Infectious agents reported to have been transmitted by blood transfusion.

Viruses	
Hepatitis viruses	Hepatitis A virus (HAV) Hepatitis B virus (HBV) Hepatitis C virus (HCV) Hepatitis D virus (HDV) (requires co-infection with HBV)
Retroviruses	Human immunodeficiency virus (HIV) 1 + 2 (+ other subtypes) Human T-cell leukaemia virus (HTLV) I + II
Herpes viruses	Human cytomegalovirus (CMV) Epstein–Barr virus (EBV) Human herpesvirus 8 (HHV-8)
Parvoviruses	Parvovirus B19
Miscellaneous viruses	GBV-C—previously referred to as hepatitis G virus (HGV) Transfusion transmitted virus (TTV) West Nile virus
Bacteria	
Endogenous	<i>Treponema pallidum</i> (syphilis) <i>Borrelia burgdorferi</i> (Lyme disease) <i>Brucella melitensis</i> (brucellosis) <i>Yersinia enterocolitica</i> / <i>Salmonella</i> spp.
Exogenous	Environmental species—staphylococcal spp./ <i>Pseudomonas</i> / <i>Serratia</i> spp.
Rickettsiae	<i>Rickettsia rickettsii</i> (Rocky Mountain spotted fever) <i>Coxiella burnettii</i> (Q fever)
Protozoa	
	<i>Plasmodium</i> spp. (malaria) <i>Trypanosoma cruzi</i> (Chagas' disease) <i>Toxoplasma gondii</i> (toxoplasmosis) <i>Babesia microti/divergens</i> (babesiosis) <i>Leishmania</i> spp. (leishmaniasis)
Prions	
	New variant Creutzfeldt–Jacob disease (nvCJD)

tests for malarial antibodies are performed. Chagas' disease is a significant problem with blood transfusion in Latin America. Bacterial infections resulting from skin commensals are most frequently transmitted by platelets stored for more than 3 days.

Prions

The risk of new variant Creutzfeldt–Jacob disease (nvCJD) is considered a threat to blood safety only in the UK. Plasma for fractionation and fresh frozen plasma for infants or children is obtained from the USA. It is unknown how many people could be infected with nvCJD. There are three rare reports of possible transmission by blood transfusion, so recipients of blood or

blood components are now excluded as blood donors in the UK. No screening tests for prions are yet available.

Techniques in blood group serology

The most important technique is based on the agglutination of red blood cells. Saline agglutination is important in detecting IgM antibodies, usually at room temperature or at 4°C (e.g. anti-A, anti-B; Fig. 30.3). Addition of colloid to the incubation or proteolytic enzyme treatment of red cells increases the sensitivity of the indirect antiglobulin test (see below), as does low ionic strength saline (LISS). These latter methods can detect a range of IgG antibodies.

The antiglobulin (Coombs') test is a fundamental and widely used test in both blood group serology and general immunology. Antihuman globulin (AHG) is produced in animals following the injection of human globulin, purified complement or specific immunoglobulin (e.g. IgG, IgA or IgM). Monoclonal preparations are also now available. When AHG is added to human red cells coated with immunoglobulin or complement components, agglutination of the red cells indicates a positive test (Fig. 30.5).

The antiglobulin test may be either direct or indirect. The **direct antiglobulin test (DAT)** is used for detecting antibody or complement already on the red cell surface where sensitization has occurred *in vivo*. The AHG reagent is added to washed red cells and agglutination indicates a positive test. A positive test occurs in haemolytic disease of the newborn, autoimmune or drug-induced immune haemolytic anaemia and haemolytic transfusion reactions.

The **indirect antiglobulin test (IAT)** is used to detect antibodies that have coated the red cells *in vitro*. It is a two-stage procedure: the first step involves the incubation of test red cells with serum; in the second step, the red cells are washed and the AHG reagent is added. Agglutination implies that the original serum contained antibody which has coated the red cells *in vitro*. This test is used as part of the routine antibody screening of the recipient's serum prior to transfusion and for detecting blood group antibodies in a pregnant woman.

Most of the above methods were originally developed for tube techniques. These were replaced by 96-well microplates but most laboratories now use gel-based technology (Fig. 30.6).

Cross-matching and pre-transfusion tests

A number of steps are taken to ensure that patients receive compatible blood at the time of transfusion.




Figure 30.5 The antiglobulin test for antibody or complement on the surface of red blood cells (RBC). The antihuman globulin (Coombs') reagent may be broad spectrum or specific for immunoglobulin G (IgG), IgM, IgA or complement (C3).




Figure 30.6 Patient antibody screening using the microcolumn (gel) system: 10 tests with two controls (tube 11 is the positive control and tube 12 the negative control) are shown. The patient's serum is tested against screening cells with known red cell phenotype. Tubes 1, 3, 5–8 and 10 show positive results. The patient's serum contained anti-Fy^a. Source: Courtesy of Mr G. Hazlehurst.

From the patient

- 1 The ABO and Rh blood group is determined.
- 2 Serum is screened for important antibodies by an indirect antiglobulin test on a large panel of antigenically-typed group O red cells.

If a red cell alloantibody is discovered in the recipient, donor blood is selected lacking the relative antigen. The most common antibodies are against Rh D, C, c, E, e and K.

From the donor

An appropriate ABO and Rh unit is selected. Donor (blood) testing is described on p. 335.

The cross-match

The techniques that may be used are described in Table 30.8.

Electronic cross-match

In this, a patient has group and antibody screen performed on two separate occasions. If both antibody screens are negative and no blood has been transfused between the test, ABO and Rh compatible blood is issued directly without further laboratory testing.

Complications of blood transfusion

Haemolytic transfusion reactions

Haemolytic transfusion reactions may be **immediate or delayed** (Table 30.9). Immediate life-threatening reactions associated with massive intravascular haemolysis are the result of complement-activating antibodies of IgM or IgG classes, usually with ABO specificity. Reactions associated with extravascular haemolysis (e.g. immune antibodies of the Rh system are which are unable

Table 30.8 Techniques used in compatibility testing. Donor cells tested against recipient serum and agglutination detected visually or microscopically after mixing and incubation at the appropriate temperature.

For detecting clinically significant IgM antibodies

Saline 37°C

For detecting immune antibodies (mainly IgG)

Indirect antiglobulin test at 37°C

Low ionic strength saline at 37°C

Enzyme-treated red cells at 37°C

Ig, immunoglobulin.

Table 30.9 Complications of blood transfusion.

Early	Late
Haemolytic reactions: immediate (IgM) or delayed (IgG)	Transmission of infection (see Table 30.7)
Reactions caused by infected blood	Transfusional iron overload (see Chapter 4)
Allergic reactions to white cells, platelets or proteins	Immune sensitization, e.g. to red cells, platelets or Rh D antigen
Pyrogenic reactions (to plasma proteins or caused by HLA antibodies)	Transfusion-associated graft-versus-host disease
Circulatory overload	
Bacterial contamination	
Air embolism	
Thrombophlebitis	
Citrate toxicity	
Hyperkalaemia	
Hypocalcaemia (infants, massive transfusion)	
Clotting abnormalities (after massive transfusion)	
Transfusion-related acute lung injury (TRALI)	
Post-transfusion purpura	
Anaphylaxis (in IgA deficient subjects)	
CMV, cytomegalovirus; HIV, human immunodeficiency virus; HLA, human leucocyte antigen.	

to activate complement) are generally usually less severe but may still be life-threatening. The cells become coated with IgG and are removed in the reticuloendothelial system. In mild cases, the only signs of a transfusion reaction may be a progressive unexplained anaemia with or without jaundice. In some cases where the pre-transfusion level of an antibody was too low to be detected in a cross-match, a patient may be reimmunized by transfusion of incompatible red cells and this will lead to a delayed transfusion reaction with accelerated clearance of the red cells. There may be rapid appearance of anaemia with mild jaundice.

Clinical features of a major haemolytic transfusion reaction

Haemolytic shock phase This may occur after only a few millilitres of blood have been transfused or up to 1–2 hours after the end of the transfusion. Clinical features include urticaria, pain in the lumbar region, flushing, headache, precordial pain, shortness of breath, vomiting, rigours, pyrexia and a fall in blood pressure. If the patient is anaesthetized this shock phase is masked. There is increasing evidence of red cell destruction, and haemoglobinuria, jaundice and disseminated intravascular coagulation (DIC) may all become apparent. Moderate leucocytosis (e.g. $15\text{--}20 \times 10^9/\text{L}$) is usual.

The oliguric phase In some patients with a haemolytic reaction there is renal tubular necrosis with acute renal failure.

Diuretic phase Fluid and electrolyte imbalance may occur during the recovery from acute renal failure.

Investigation of an immediate transfusion reaction

If a patient develops features suggesting a severe transfusion reaction, the transfusion should be stopped and investigations for blood group incompatibility and bacterial contamination of the blood must be initiated.

- Most severe reactions occur because of clerical errors in the handling of donor or recipient blood specimens. Therefore it must be established that the identity of the recipient (from the patient's wristband) is the same as that on the compatibility label and that this corresponds with the actual unit being transfused.
- The unit of donor blood and post-transfusion samples of the patient's blood should be sent to the laboratory who will:
 - repeat the group on pre- and post-transfusion samples and on the donor blood, and repeat the cross-match;
 - perform a direct antiglobulin test on the post-transfusion sample;
 - check the plasma for haemoglobinaemia;
 - perform tests for DIC; and
 - examine the donor sample directly for evidence of gross bacterial contamination and set up blood cultures from it at 20°C and 37°C. If the clinical picture is suggestive of bacterial infection, blood cultures must be taken from the patient and broad-spectrum intravenous antibiotics started.

- 3 A post-transfusion sample of urine must be examined for haemoglobinuria.
- 4 Further samples of blood are taken 6 hours and/or 24 hours after transfusion for a blood count and bilirubin, free haemoglobin and methaemalbumin (see p. XXX) estimations.
- 5 In the absence of positive findings, the patient's serum is examined 5–10 days later for red cell or white cell antibodies.

Management of patients with major haemolysis

The principal object of initial therapy is to maintain the blood pressure and renal perfusion. Intravenous dextran, plasma or saline and furosemide are sometimes needed. Hydrocortisone 100 mg intravenously and an antihistamine may help to alleviate shock. In the event of severe shock, support with intravenous adrenaline 1:10 000 in small incremental doses may be required. Further compatible transfusions may be required in severely affected patients. If acute renal failure occurs this is managed in the usual way, if necessary with dialysis until recovery occurs.

Other transfusion reactions

Hyperhaemolysis syndromes Some patients, particularly with sickle cell anaemia, haemolyse donor blood even though no alloantibodies to red cells can be detected. The haemolysis appears to be due to overactivity of the recipient's macrophages. It is prevented by infusions of gammaglobulin and corticosteroid therapy.

Febrile reactions because of white cell antibodies Human leucocyte antigen (HLA) antibodies (see below and Chapter XX) are usually the result of sensitization by pregnancy or a previous transfusion. They produce rigors, pyrexia and, in severe cases, pulmonary infiltrates. They are minimized by giving leucocyte-depleted (i.e. filtered) packed cells (see below).

Febrile or non-febrile non-haemolytic allergic reactions These are usually caused by hypersensitivity to donor plasma proteins and, if severe, can result in anaphylactic shock. The clinical features are urticaria, pyrexia and, in severe cases, dyspnoea, facial oedema and rigors. Immediate treatment is with antihistamines and hydrocortisone. Adrenaline is also useful. Washed red cells or frozen red cells may be needed for further transfusions if the majority of plasma-removed blood (e.g. saline, adenine, glucose, mannitol (SAGM) blood) causes reactions.

Post-transfusion circulatory overload The management is that of cardiac failure. These reactions are prevented by a slow transfusion of packed red cells or of the blood component required, accompanied by diuretic therapy.

Transfusion of bacterially contaminated blood This is very rare but may be serious. It can present with circulatory collapse. It is a particular problem with platelet packs that are stored at 20–24°C.

Graft-versus-host disease (GVHD) This may occur when live lymphocytes are transfused to an immunocompromised patient. It is prevented by irradiation of the blood products for susceptible recipients (Table 30.10). It is uniformly fatal.

Transfusion related acute lung injury (TRALI) This presents within 6 hours of an infusion as pulmonary

Table 30.10 Indications for irradiated blood products.

Recipients of allogeneic stem cell transplantation
Recipients of autologous stem cell transplantation
Bone marrow or stem cell or autologous donor: for 7 days prior to harvest and during harvest
Recipients of solid organ transplants when receiving immune suppressive therapy
Patients with Hodgkin lymphoma
Patients with acute leukaemia
Patients treated with fludarabine, bendamustine, other purine antagonists
Patients treated with anti-CD52
Inherited immunodeficiency
Intrauterine and neonatal transfusions

infiltrates with chest symptoms depending on severity. It is caused by positive transfer of leucoagglutins HLA antibodies in donor plasma which cause endothelial and epithelial injury. Most of the donors are multiparous women. Management is supportive.

Post-transfusion purpura This is a rare problem of severe thrombocytopenia 7–10 days after transfusion of a platelet-containing product, usually red cells. It is caused by an antibody in the recipient (resulting from previous transfusion or pregnancy) which is usually directed against a platelet-specific antigen HPA-1a (PI^A). Both the transfused and recipient platelets are destroyed by the immune complexes. It is usually self-limiting but immunoglobulin or plasma exchange may be needed.

Viral transmission Post-transfusion hepatitis may be caused by one of the hepatitis viruses, although CMV and Epstein–Barr virus (EBV) have also been implicated. Post-transfusion viral hepatitis, HTLV or HIV infection are very rarely seen because of routine screening of all blood donations.

Other infections Toxoplasmosis, malaria and syphilis may be transmitted by blood transfusion. Transfusion-transmitted nvCJD has probably occurred in three cases in the UK.

Post-transfusional iron overload Repeated red cell transfusions over many years, in the absence of blood loss, cause deposition of iron initially in reticuloendothelial tissue at the rate of 200–250 mg/unit of red cells. After 50 units in adults, and lesser amounts in children, the liver, myocardium and endocrine glands are damaged with clinical consequences. This becomes a major problem in thalassaemia major and other severe chronic refractory anaemias (see Chapter 4).

Reduction of blood product use

In the light of transfusion risks, and limited resources, appropriate use of blood components is of ever increasing importance.

Preoperative correction of anaemia (particularly iron deficiency) and cessation of anti-platelet therapies (e.g. aspirin) where possible, together with lower trigger levels for red cell transfusions (haemoglobin 70–80 g/L in most surgical patients) can all help to reduce blood use.

In surgery the use of alternative fluid replacement, intraoperative or postoperative cell salvage, biological alternatives (e.g. erythropoietin, recombinant clotting factors, recombinant activated clotting factor VII (VIIa) or fibrin glue) all may help.

Blood components

A blood donation is taken by an aseptic technique into plastic bags containing an appropriate amount of anticoagulant, usually citrate, phosphate, dextrose (CPD). The citrate anticoagulates the blood by combining with the blood calcium. Three components are made by initial centrifugation of whole blood: red cells, buffy coat and plasma (Fig. 30.1).

Red cells are stored at 4–6°C for up to 35 days, depending on the preservative. After the first 48 hours there is a slow progressive K⁺ loss from the red cells into the plasma. In cases where infusion of K⁺ could be dangerous, fresh blood should be used (e.g. for exchange transfusion in haemolytic disease of the newborn). During red cell storage there is a fall in 2,3-diphosphoglycerate (2,3-DPG) but after transfusion 2,3-DPG levels return to normal within 24 hours. Optimum additive solutions have been developed to increase the shelf-life of plasma-depleted red cells by maintaining both adenosine triphosphate (ATP) and 2,3-DPG levels.

Platelets and plasma may also be collected by apheresis (and centrifuging).

Leucodepletion

In many countries, including Britain, blood products are now routinely filtered to remove the majority of white cells, a process known as leucodepletion. This is usually performed soon after collection and prior to processing and is more effective than filtration of blood at the bedside (Table 30.1). A blood component is defined as leucocyte-depleted if there are less than 5×10^6 /L white cells present.

Leucodepletion reduces the incidence of febrile transfusion reactions and HLA alloimmunization. It is effective at preventing transmission of CMV infection and in addition should reduce the theoretical possibility of transmission of nvCJD in countries where this has been reported.

Red cells

Packed (plasma-depleted) red cells are the treatment of choice for most transfusions (Fig. 30.7a). In older subjects, a diuretic is often given simultaneously and the infusion should be sufficiently slow to avoid circulatory overload. Iron chelation therapy, to avoid iron overload, should be considered with patients on a regular transfusion programme to avoid iron overload (see Chapter 4).




Figure 30.7 Blood components: (a) plasma-depleted red cells; (b) platelets; and (c) fresh frozen plasma.

Recombinant erythropoietin is widely used to reduce transfusion requirements (e.g. in patients with renal failure, on dialysis, cancer patients and in myelodysplasia). Factor VIIa can reduce transfusion need in patients with major haemorrhage (e.g. at surgery or after trauma).

Red cell substitutes are under development but have not yet proven clinically valuable. These synthetic oxygen-carrying substitutes are often fluorinated hydrocarbons and stromal-free pyridoxylated and polymerized haemoglobin solutions.

Autologous donation and transfusion

Anxiety over HIV and other infections has increased the demand for auto-transfusion. There are three ways of administering an autologous transfusion:

- 1 Predeposit** Blood is taken from the potential recipient in the weeks immediately prior to elective surgery.
- 2 Haemodilution** Blood is removed immediately prior to surgery once the patient has been anaesthetized and then reinfused at the end of the operation.

3 Salvage Blood lost during the operation is collected during heavy blood loss and then reinfused.

Autotransfusion is the safest form of transfusion with regard to transmission of viral disease but has a higher risk of bacterial contamination and of clerical errors. The individual involved must be fit enough to donate blood and the predicted operative replacement transfusion should be 2–4 units. Larger replacement transfusions would require blood to be collected over a longer period and red cells stored in the frozen state, which is both labour intensive and expensive. The high cost of storage and initial restriction of its use to patients undergoing elective surgery means that it can benefit only a minor proportion of the total number of blood recipients. Preoperative autotransfusion is largely reserved for those patients with multiple antibodies for whom it is difficult to identify matching donor blood.

Granulocyte concentrates

These are prepared as buffy coats or on blood cell separators from normal healthy donors or from patients with chronic myeloid leukaemia. They have been used in patients with severe neutropenia ($<0.5 \times 10^9/L$) who are not responding to antibiotic therapy but it is not usually possible to give sufficient amounts. They may transmit CMV infection and must be irradiated to eliminate the risk of causing GVHD.

Platelet concentrates

These are harvested by cell separators or from individual donor units of blood (Fig. 30.7b). They are stored at room temperature. Platelet transfusion is used in patients who are thrombocytopenic, or have disordered platelet function and who are actively bleeding (therapeutic use) or are at serious risk of bleeding (prophylactic use).

For prophylaxis, the platelet count should be kept above $5\text{--}10 \times 10^9/L$ unless there are additional risk factors such as sepsis, drug use or coagulation disorders for which the threshold should be higher. For invasive procedures (e.g. liver biopsy or lumbar puncture) the platelet count should be raised to above $50 \times 10^9/L$. For brain or eye surgery the count should be $>100 \times 10^9/L$.

Therapeutic use is indicated in bleeding associated with platelet disorders. In massive haemorrhage the count should be kept above $50 \times 10^9/L$ (see Chapter 26).

Platelet transfusions should be avoided in autoimmune thrombocytopenic purpura unless there is serious haemorrhage. They are contraindicated in heparin-induced thrombocytopenia, thrombotic thrombocytopenic purpura and haemolytic uraemic syndrome (see p. 285).

Refractoriness to platelet transfusions is defined by a poor platelet increment post-transfusion (less than $7.5 \times 10^9/L$ at 1 hour or less than $4.5 \times 10^9/L$ at 24 hours). The causes are either immunological (mostly HLA alloimmunization) or non-immunological (sepsis, hypersplenism, DIC, drugs). Platelets express HLA class I (but not class II) antigens and HLA-matched or cross-match-compatible platelets are needed for patients with HLA antibodies.

The need for platelet transfusions is likely to be reduced with the introduction of direct stimulators of platelet production such as romiplostim or eltrombopag.

Preparations from human plasma

Fresh frozen plasma (FFP)

Rapidly frozen plasma separated from fresh blood is stored at less than -30°C . Frozen plasma is usually prepared from single donor units, although pooled products are also available. Its main use is for the replacement of coagulation factors (e.g. when specific concentrates are unavailable) or after massive transfusions, in liver disease and DIC, after cardiopulmonary bypass surgery, to reverse a warfarin effect, and in thrombotic thrombocytopenic purpura (see p. 285). Virally inactivated forms of FFP are now available. Male donors are preferred to reduce the risk of TRALI (see p. 342).

Human albumin solution (4.5%)

This is a useful plasma volume expander when a sustained osmotic effect is required prior to the administration of blood, but it should not be given in excess. It is also used for fluid replacement in patients undergoing plasmapheresis and sometimes for fluid replacement in selected patients with hypoalbuminaemia.

Human albumin solution (20%) (salt-poor albumin)

This may be used in severe hypoalbuminaemia when it is necessary to use a product with minimal electrolyte content. Principal indications for its use are patients with nephrotic syndrome or liver failure.

Cryoprecipitate

This is obtained by thawing FFP at 4°C and contains concentrated factor VIII and fibrinogen. It is stored at less than -30°C or, if lyophilized, at $4\text{--}6^\circ\text{C}$, and was used widely as replacement therapy in haemophilia A and von Willebrand disease before more purified preparations of factor VIII became available. Its main use is in fibrinogen replacement in disseminated intravascular coagulation (DIC) or massive transfusion or hepatic failure.

Freeze-dried factor VIII concentrates

These are also used for treating haemophilia A or von Willebrand disease. The small volume makes them ideal for children, surgical cases, patients at risk from circulatory overload and for those on home treatment. Their use is declining as recombinant forms of factor VIII become widely available.

Freeze-dried factor IX–prothrombin complex concentrates

A number of preparations are available that contain variable amounts of factors II, VII, IX and X. They are mainly used for treating factor IX deficiency (Christmas disease) but are also

used in patients with liver disease or in haemorrhage following overdose with oral anticoagulants or in patients with factor VIII inhibitors. There is a risk of thrombosis.

Protein C concentrate

This is used in severe sepsis with disseminated intravascular coagulation (e.g. meningococcal septicaemia) to reduce thrombosis resulting from depletion of protein C.

Immunoglobulin

Pooled immunoglobulin is a valuable source of antibodies against common viruses. It is used in hypogammaglobulinaemia for passive protection against viral and bacterial disease. Repeated doses are needed, for example in the winter months at 3–4-week intervals. It may also be used in immune thrombocytopenia and other acquired immune disorders (e.g. post-transfusion purpura or alloimmune neonatal thrombocytopenia).

Specific immunoglobulin

This may be obtained from donors with high titres of antibody (e.g. anti-RhD, anti-hepatitis B, anti-herpes zoster or anti-rubella).

Acute blood loss and massive haemorrhage

After a single episode of blood loss, there is initial vasoconstriction with a reduction in total blood volume. The plasma volume rapidly expands and the haemoglobin and packed cell volume fall and there is a rise in neutrophils and platelets. The reticulocyte response begins on the second or third day and lasts 8–10 days. The haemoglobin begins to rise by about the seventh day but, if iron stores have become depleted, the haemoglobin may not subsequently rise to normal. Clinical assessment is needed to gauge whether blood transfusion is needed. This is usually unnecessary in adults at losses less than 500 mL unless haemorrhage is continuing, and may not be needed with losses of up to 1.5 L. Blood transfusion is not without risks and should not be undertaken lightly.

- Blood transfusion involves the safe transfer of blood components from a donor to a recipient. Most commonly this is red cells and the red cells must be matched between recipient and donor.
- Careful donor selection and microbiological testing help to protect both donor and recipient.
- Red cells contain over 400 antigens. It is the ABO and Rh systems that are most important in transfusion. Subjects lacking an antigen (e.g. group A or B) may develop a naturally occurring antibody to it, usually IgM. These antibodies in a recipient may haemolyse or opsonize donor red cells if these contain the antigen.
- Antibodies may also develop from exposure to the antigen by a transfusion or pregnancy. Cross-matching of donor red cells with recipient plasma is therefore carried out to ensure they are compatible.
- Complications of blood transfusion include haemolytic reactions, febrile reactions to white cells or proteins, circulatory overload, transmission of infections, especially viral, and, in the longer term, iron overload.
- Blood components other than red cells can also be transfused. These include platelets and protein products including fresh frozen plasma, albumin solutions, coagulation factor concentrates and immunoglobulin.




Now visit www.wiley.com/go/essentialhaematology to test yourself on this chapter.

CHAPTER 31

Pregnancy and neonatal haematology

Key topics

■ Haematology of pregnancy	347
■ Neonatal haematology	348
■ Haemolytic disease of the newborn	350



Haematology of pregnancy

Pregnancy places extreme stresses on the haematological system and an understanding of the physiological changes that result is obligatory in order to interpret any need for therapeutic intervention.

Physiological anaemia

Physiological anaemia is the term often used to describe the fall in haemoglobin (Hb) concentration that occurs during normal pregnancy (Fig. 31.1). Blood plasma volume increases by approximately 1250 mL, or 45%, above normal by the end of gestation and although the red cell mass itself increases by some 25% this still leads to a fall in Hb concentration. Values below 100 g/L in the first trimester, 105 g/L in the second and 100 g/L in the third or postpartum are abnormal and require investigation.

Iron deficiency anaemia

Up to 600 mg iron is required for the mother's increase in red cell mass and a further 300 mg for the fetus. Despite an increase in iron absorption, few women avoid depletion of iron reserves by the end of pregnancy.

In uncomplicated pregnancy, the mean corpuscular volume (MCV) typically rises by approximately 4 fL. A fall in red cell MCV is the earliest sign of iron deficiency. Later, the mean corpuscular haemoglobin (MCH) falls and finally anaemia results. Early iron deficiency is likely if the serum ferritin is below 30 µg/L together with serum iron below 10 mmol/L and should be treated with oral iron supplements. Routine iron supplementation in pregnancy is not carried out in the UK.

Folate and vitamin B₁₂ deficiency

Folate requirements are increased approximately two-fold in pregnancy and serum folate levels fall to approximately half the normal range with a less dramatic fall in red cell folate. In some parts of the world, megaloblastic anaemia during pregnancy is common because of a combination of poor diet and exaggerated folate requirements. Given the protective effect of folate against neural tube defects (NTDs), folic acid 400 µg/day (5 mg if there has been a previous NTD pregnancy) should be taken periconceptually and throughout pregnancy. Food fortification with folate is now being practised in many countries and has been associated with a fall in incidence of NTDs. Vitamin B₁₂ deficiency is rare during pregnancy although serum vitamin B₁₂ levels fall to below normal in 20–30% of pregnancies and low values are sometimes the cause of diagnostic confusion.

Thrombocytopenia

The platelet count falls by an average of 10% in an uncomplicated pregnancy. In approximately 7% of women this fall is more severe and can result in thrombocytopenia (platelet count less than $140 \times 10^9/L$). In over 75% of cases this is mild and of unknown cause, referred to as *incidental thrombocytopenia of pregnancy*. Approximately 21% of cases are secondary to a hypertensive disorder and 4% are associated with immune thrombocytopenic purpura (ITP; Fig. 31.2).

Incidental thrombocytopenia of pregnancy This is a diagnosis of exclusion usually detected at the time of delivery. The platelet count is above $70 \times 10^9/L$ and recovers within 6 weeks. No treatment is needed and the infant is not affected.




Figure 31.1 Haematological changes during pregnancy.




Figure 31.2 Causes of thrombocytopenia during pregnancy.

Thrombocytopenia of hypertensive disorders This is variable in severity but the platelet count rarely falls to below $40 \times 10^9/L$. It is more severe when associated with pre-eclampsia when the primary treatment is as rapid delivery as possible. The HELLP syndrome (haemolysis, elevated liver enzymes and low platelets) falls into this category. It is associated with prolongation of prothrombin time (PT) and activated partial thromboplastin time (APTT).

Idiopathic thrombocytopenic purpura (ITP) (see p. 282) In pregnancy, ITP represents a particular problem, both to the mother and to the fetus, as the antibody crosses the placenta and the fetus may become severely thrombocytopenic. Like other adults, pregnant women with ITP and platelet counts higher than $50 \times 10^9/L$ do not usually need treatment. Treatment is required for women with platelet counts below $10 \times 10^9/L$ and for those with platelet counts of $10\text{--}30 \times 10^9/L$ who are in their second or third trimester or who are bleeding. Treatment is with steroids, intravenous immunoglobulin G (IgG), rituximab or splenectomy as appropriate. Thrombopoietin receptor antagonists are avoided because of potential teratogenic side-effects.

At delivery, umbilical vein blood sampling or fetal scalp vein sampling to measure the fetal platelet count may be offered although their exact role is unclear. In general, caesarean section is not indicated when the maternal platelet count is above $50 \times 10^9/L$ unless the fetal platelet count is known to be less than $20 \times 10^9/L$. Platelet transfusion may be given to mothers in labour with very low platelet counts or who are actively bleeding.

Newborns of mothers with ITP should have a blood count measured for the first 5 days of life as the platelet count may progressively drop. A count greater than $50 \times 10^9/L$ is reassuring. Cerebral ultrasonography may be performed to look for intracranial haemorrhage (ICH). In newborns without

evidence of ICH, treatment with intravenous IgG is appropriate if the infant's platelet count is less than $20 \times 10^9/L$. Neonates with thrombocytopenia and ICH should be treated with steroids and intravenous IgG therapy.

Haemostasis and thrombosis

Pregnancy leads to a hypercoagulable state with consequent increased risks of thromboembolism and disseminated intravascular coagulation (DIC; see p. 297). There is an increase in plasma factors VII, VIII, X and fibrinogen with shortening of PT and APTT; fibrinolysis is suppressed. These changes last for up to 2 months into the puerperal period and the incidence of thrombosis during this period is increased. There is an association between thrombophilic conditions in the mother and with recurrent fetal loss (see p. 307). This is presumed to result from placental thrombosis and infarction.

Treatment of thrombosis

Warfarin has no role in management. It crosses the placenta and in addition is associated with embryopathy, especially between 6 and 12 weeks' gestation. Low molecular weight heparin is now the treatment of choice because it can be given once daily and is less likely than unfractionated heparin to cause osteoporosis.

Neonatal haematology

Normal blood count

The cord blood Hb varies between approximately 165 and 170 g/L and is influenced by the timing of cord clamping (Fig. 31.3). The reticulocyte count is initially high (2–6%) but




Figure 31.3 Typical profile of the blood count in the neonatal period.

falls to below 0.5% at 1 week as erythropoiesis is suppressed in response to the marked increase in the oxygenation of tissues after birth. This is associated with a progressive fall in Hb to approximately 100–110 g/L at 8 weeks, from which point it

recovers to 125 g/L at around 6 months. The lower limit of normal during childhood is 110 g/L. Preterm infants have a more dramatic fall in Hb to 70–90 g/L at 8 weeks and are more prone to iron and folate deficiency in the first few months of life.

In the blood film, nucleated red cells will be seen for the first 4 days and for up to 1 week in preterm infants. Numbers are increased in cases of hypoxia, haemorrhage or haemolytic disease of the newborn (HDN). MCV averages 119 fL but falls to adult levels by around 9 weeks. By 1 year, the MCV has fallen to around 70 fL and rises throughout childhood again to reach adult levels at puberty. Neutrophils are initially high at birth and fall to plateau at 4 days – from this point on the lymphocyte count is higher than neutrophils throughout childhood.

Anaemia in the neonate

This should be considered for Hb below 140 g/L at birth. The clinical significance of anaemia is compounded by the high (70–80%) levels of HbF at birth, as this is less effective than HbA at releasing oxygen to the tissues (see p. 17). Causes include the following (Fig. 31.4):

- Haemorrhage** Fetomaternal, twin–twin, cord, internal, placenta.




Figure 31.4 The investigation of neonatal anaemia. * The DAT test may be negative in HDN due to ABO incompatibility. DAT, direct antiglobulin test; HDN, haemolytic disease of the newborn; MCV, mean corpuscular volume.

2 Increased destruction Haemolysis (immune or non-immune) or infection.

3 Decreased production Congenital red cell aplasia, infection (e.g. parvovirus). Anti-Kell causes alloimmune anaemia of the fetus and newborn with decreased erythropoiesis.

Generally, anaemia at birth is usually secondary to immune haemolysis or haemorrhage; non-immune causes of haemolysis appear within 24 hours. Impaired red cell production is usually not apparent for at least 3 weeks. Haemolysis is often associated with severe jaundice and the causes include HDN, autoimmune haemolytic anaemia (AIHA) in the mother and congenital disorders of the red cell membrane or metabolism.

Red cell transfusion may be needed for symptomatic anaemia with Hb less than 105 g/L or a higher threshold if there is severe cardiac or respiratory disease.

Anaemia of prematurity

Premature infants have a more marked fall in Hb after birth and this is termed **physiological anaemia of prematurity**. Features include a slowly falling Hb, normal blood film and reticulocytopenia. It can be minimized by ensuring adequate iron and folate replacement and limiting phlebotomy. Erythropoietin is used in some centres.

Neonatal polycythaemia

This is defined as a venous haematocrit over 0.65 and can occur with twin–twin transfusion, intrauterine growth restriction and maternal hypertension or diabetes. If symptoms are present it should be treated with partial exchange transfusion using a crystalloid solution.

Fetomaternal alloimmune thrombocytopenia

Fetomaternal alloimmune thrombocytopenia (FMAIT) results from a process similar to that of HDN. Fetal platelets that possess a paternally inherited antigen (HPA-1a in 80%) that is not present on maternal platelets can sensitize the mother to make antibodies which cross the placenta, coat the platelets, which are then destroyed by the reticuloendothelial system, and lead to serious bleeding, including intracranial haemorrhage. Alloimmune thrombocytopenia differs from HDN in that 50% of cases occur in the first pregnancy. Its incidence is approximately 1 in 1000–5000 births.

Thrombocytopenia can lead to serious, sometimes fatal, bleeding *in utero* or after birth. Treatment is unsatisfactory. Severe postnatal cases may be treated with a platelet transfusion that is negative for the relevant antigen. Antenatal treatment may be either maternal intravenous immunoglobulin or fetal transfusion with HPA-compatible platelets.

Other causes of neonatal thrombocytopenia include perinatal infection, placental insufficiency and congenital genetic causes.

Coagulation

Standard tests need to be interpreted with caution in the neonate. The activated partial thromboplastin time (APTT) and

prothrombin time (PT) are prolonged because of reduced levels of the vitamin K-dependent factors II, VII, IX and X, which become normal at around 6 months. Neonates have an increased risk of thrombosis. This is a result of physiologically low levels of inhibitors of coagulation and the use of indwelling vascular catheters. Antithrombin (AT) and protein C levels are approximately 60% of normal for the first 3 months. Homozygous protein C deficiency is associated with fulminant purpura fulminans in early life. Therapeutic protein C concentrates are now available. Homozygous AT deficiency usually presents later in childhood but arterial and venous thrombosis may also occur in the neonate.

Haemorrhagic disease of the newborn is discussed in the next section of this chapter.

Haemolytic disease of the newborn

HDN is the result of **red cell alloimmunization** in which IgG antibodies pass from the maternal circulation across the placenta into the circulation of the fetus where they react with fetal red cells and lead to their destruction. Anti-D antibody is responsible for most cases of severe HDN although anti-c, anti-E, anti-K and a wide range of other antibodies are found in occasional cases (see Table 30.3). Although antibodies against the ABO blood group system are the most frequent cause of HDN this is usually mild.

Rh haemolytic disease of the newborn


When an Rh D-negative (p. 335) woman has a pregnancy with an Rh D-positive fetus, Rh D-positive fetal red cells cross into the maternal circulation (especially at parturition and during the third trimester) and sensitize the mother to form anti-D. The mother could also be sensitized by a previous miscarriage, amniocentesis or other trauma to the placenta or by blood transfusion. Anti-D crosses the placenta to the fetus during the next pregnancy, coats RhD-positive fetal red cells and results in reticuloendothelial destruction of these cells, causing anaemia and jaundice. If the father is heterozygous for D antigen, and mother RhD negative there is a 50% probability that the fetus will be D-positive. The fetal Rh D genotype can be established by polymerase chain reaction (PCR) analysis for the presence of Rh D in a maternal blood sample.

The main aim of management is to prevent anti-D antibody formation in Rh D-negative mothers. This can be achieved by the administration of small amounts of anti-D antibody which ‘mop up’ and destroy Rh D-positive fetal red cells before they can sensitize the immune system of the mother to produce anti-D.


Prevention of Rh immunization

At the time of booking, all pregnant women should have their ABO and Rh group determined and serum screened for antibodies at least twice during the pregnancy. All non-sensitized Rh D-negative women should be given at least 500 units (100 µg) of anti-D at 28 and 34 weeks’ gestation to reduce the risk of

sensitization from fetomaternal haemorrhage. Fetal Rh D typing from DNA in maternal blood can be used before 28 weeks. If the fetus is Rh D-negative, no further anti-D prophylaxis is needed. In addition, at birth the babies of Rh D-negative women who do not have antibodies must have their cord blood grouped for ABO and Rh. If the baby's blood is Rh D-negative, the mother will require no further treatment. If the baby is Rh D-positive, prophylactic anti-D should be administered to the mother at a minimum dose of 500 units intramuscularly within 72 hours of delivery. A **Kleihauer test** is performed. This uses differential staining to estimate the number of fetal cells in the maternal circulation (Fig. 31.5a). If the Kleihauer is positive many centres will perform flow cytometry for a more accurate estimate of the volume of fetomaternal haemorrhage (FMH) (Fig. 31.5b). The chance of developing antibodies is related to the number of fetal cells found. The dose of anti-D is increased if there is greater than 4 mL transplacental haemorrhage. Anti-D IgG (125 units) is given for each 1 mL of FMH greater than 4 mL.



(a)



(b)

Figure 31.5 (a) Kleihauer test for fetal red cells; a deeply eosin-staining cell containing fetal haemoglobin is seen at the centre. Haemoglobin has been eluted from the other red cells by an incubation at acid pH and these appear as colourless ghosts. **(b)** Determination by flow cytometry of the number of RhD fetal cells in maternal blood using fluorescent-labelling of antibody to RhD, the mother being Rhdd. Source: Courtesy Dr. W Erber.




Figure 31.6 Doppler ultrasonography of the circle of Willis in a fetus. The cursor is placed over the middle cerebral artery and an increased blood velocity correlates with anaemia. Source: From Kumar S. and Regan F. (2005) *BMJ* 330: 1255–8. Reproduced with permission of BMJ.

Sensitizing episodes during pregnancy

Anti-D IgG should be given to Rh D-negative women who have potentially sensitizing episodes during pregnancy: 250 units is given if the event occurs up to week 20 of gestation and 500 units thereafter, followed by a Kleihauer test. Potentially sensitizing events include therapeutic termination of pregnancy, spontaneous miscarriage after 12 weeks' gestation, ectopic pregnancy and invasive antenatal diagnostic procedures.

Treatment of established anti-D sensitization


If anti-D antibodies are detected during pregnancy they should be quantified at regular intervals. The clinical severity is related to the strength of anti-D present in maternal serum but is also affected by such factors as the IgG subclass, rate of rise of antibody and past history. The development of haemolytic disease in the fetus can be assessed by velocimetry of the fetal middle cerebral artery by Doppler ultrasonography, as increased velocities correlate with fetal anaemia (Fig. 31.6). If anaemia is detected, fetal blood sampling and intrauterine transfusion of irradiated Rh D-negative packed red cells may be indicated.

Clinical features of HDN

- 1 Severe disease** Intrauterine death from hydrops fetalis (Fig. 31.7a).
- 2 Moderate disease** The baby is born with anaemia and jaundice and may show pallor, tachycardia, oedema and hepatosplenomegaly. If the unconjugated bilirubin is not controlled and reaches levels exceeding $250 \mu\text{mol/L}$, bile pigment deposition in the basal ganglia may lead to **kernicterus** – central nervous system damage with generalized spasticity and possible subsequent mental deficiency, deafness and epilepsy. This problem becomes acute after birth as maternal clearance of fetal bilirubin ceases and conjugation of bilirubin by the neonatal liver has not yet reached full activity.



(a)



(b)

Figure 31.7 (a) Ultrasound features of hydrops fetalis showing skin oedema, hepatomegaly and ascites. Source: From Kumar S. and Regan F. (2005) *BMJ* 330: 1255–8. Reproduced with permission of BMJ. (b) Rh haemolytic disease of the newborn (erythroblastosis fetalis): peripheral blood film showing large numbers of erythroblasts, polychromasia and crenated cells.

3 Mild disease Mild anaemia with or without jaundice.

Investigations will reveal variable anaemia with a high reticulocyte count; the baby is Rh D-positive, the direct antiglobulin test is positive and the serum bilirubin raised. In moderate and severe cases, many erythroblasts are seen in the blood film (Fig. 31.7b); this is known as *erythroblastosis fetalis*.

Treatment

Exchange transfusion may be necessary; the indications for this include severe anaemia (Hb <100 g/L at birth) and severe or rapidly rising hyperbilirubinaemia. More than one exchange transfusion may be required and 500 mL is usually sufficient for each exchange. The donor blood should be less than 5 days old, CMV negative, irradiated, Rh D-negative and ABO compatible with the baby's and mother's serum. Phototherapy (exposure of the infant to bright light of appropriate wavelength) degrades bilirubin and reduces the likelihood of kernicterus.

ABO haemolytic disease of the newborn


In 20% of births, a mother is ABO incompatible with the fetus. Group A and group B mothers usually have only IgM ABO antibodies (p. 334). The majority of cases of ABO HDN are caused by 'immune' IgG antibodies in group O mothers. Although 15% of pregnancies in white people involve a group O mother with a group A or group B fetus, most mothers do not produce IgG anti-A or anti-B and very few babies have severe enough haemolytic disease to require treatment. Exchange transfusions are needed in only 1 in 3000 infants. The mild course of ABO HDN is partly explained by the A and B antigens not being fully developed at birth and by partial neutralization of maternal IgG antibodies by A and B antigens on other cells, in the plasma and tissue fluids.

In contrast to Rh HDN, ABO disease may be found in the first pregnancy and may or may not affect subsequent pregnancies. The direct antiglobulin test on the infant's cells may be negative or only weakly positive. Examination of the blood film shows autoagglutination, spherocytosis, polychromasia and erythroblastosis.

- Pregnancy results in multiple changes in the haematological systems.
- There is a fall in haemoglobin because of an increased plasma volume that is proportionally greater than a 25% increase in red cell mass.
- Iron deficiency is frequent; folate deficiency is associated with maternal anaemia and with neural tube defects (NTDs) in the fetus.
- Serum vitamin B₁₂ levels fall in pregnancy but recover postpartum. Platelets counts also fall on average by 10%, but in some women this physiological fall may be severe or may be caused by immune thrombocytopenia.
- Pregnancy is a hypercoagulable state with increased levels of coagulation factors and risk of thrombosis or disseminated intravascular coagulation.
- Neonates have higher haemoglobin levels than adults. Anaemia at birth is usually caused by haemorrhage or immune haemolysis.
- Haemolytic disease of the newborn is brought about by Rh D antibodies made by a Rh D-negative mother crossing the placenta. It may cause death of the fetus (hydrops fetalis) or haemolytic anaemia. It is now rare because of administration of Rh anti-D to Rh D-negative mothers at the time of exposure to Rh D-positive fetal cells or blood products.
- ABO haemolytic disease of the newborn is more frequent. It is usually mild and may occur in the first pregnancy. It is most frequently caused by group O mothers making immune IgG antibodies (which cross the placenta) against a group A or B fetus.




Now visit www.wileyessential.com/haematology to test yourself on this chapter.



Appendix

World Health Organization classification of tumours of the haematopoietic and lymphoid tissues



Myeloproliferative neoplasms

Chronic myelogenous leukaemia, *BCR-ABL1* positive
 Chronic neutrophilic leukaemia
 Polycythaemia vera
 Primary myelofibrosis
 Essential thrombocythaemia
 Chronic eosinophilic leukaemia, not otherwise specified
 Mastocytosis
 Cutaneous mastocytosis
 Systemic mastocytosis
 Mast cell leukaemia
 Mast cell sarcoma
 Extracutaneous mastocytoma
 Myeloproliferative neoplasms, unclassifiable
 Myeloid and lymphoid neoplasms associated with
 eosinophilia and abnormalities of *PDGFRA*, *PDGFRB* or
 FGFR1
 Myeloid and lymphoid neoplasms associated with *PDGFRA*
 rearrangement
 Myeloid neoplasms associated with *PDGFRB* rearrangement
 Myeloid and lymphoid neoplasms with *FGFR1*
 abnormalities

Myelodysplastic/myeloproliferative neoplasms

Chronic myelomonocytic leukaemia
 Atypical chronic myeloid leukaemia, *BCR-ABL1* negative
 Juvenile myelomonocytic leukaemia
 Myelodysplastic/myeloproliferative neoplasms, unclassifiable
 Refractory anaemia with ringed sideroblasts (RARS)
 associated with marked thrombocytosis*

Myelodysplastic syndromes

Refractory cytopenia with unilineage dysplasia
 Refractory anaemia
 Refractory neutropenia
 Refractory thrombocytopenia
 Refractory anaemia with ring sideroblasts
 Refractory cytopenia with multilineage dysplasia
 Refractory anaemia with excess blasts
 Myelodysplastic syndromes associated with isolated del(5q)
 Myelodysplastic syndromes, unclassifiable
 Myelodysplastic syndromes in children

Acute myeloid leukaemia

Acute myeloid leukaemia (AML) with recurrent genetic
 abnormalities
 AML with t(8;21)(q22;q22), *RUNX1-RUNX1T1*
 AML with inv(16)(p13;q22) or t(16;16)(p13.1;q22),
 CBFB-MYH11

Acute promyelocytic leukaemia with t(15;17)(q22;q11-12),
 PML-RARA
 AML with t(9;11)(p22;q23), *MLLT3-MLL*
 AML with t(6;9)(p23;q34), *DEK-NUP214*
 AML with inv(3)(q21;q26.2) or t(3;3)(q21;q26.2),
 RPN1-EVI1
 AML (megakaryoblastic) with t(1;22)(p13;q13),
 RBM15-MKL1
 AML with mutated *NPM1**
 AML with mutated *CEBPA**
 AML with myelodysplasia-related changes
 Therapy-related myeloid neoplasms
 AML, not otherwise categorized
 AML with minimal differentiation
 AML without maturation
 AML with maturation
 Acute myelomonocytic leukaemia
 Acute monoblastic and monocytic leukaemia
 Acute erythroid leukaemia
 Acute erythroid leukaemia, erythroid/myeloid
 Acute pure erythroid leukaemia
 Acute megakaryoblastic leukaemia
 Acute basophilic leukaemia
 Acute panmyelosis with myelofibrosis
 Myeloid sarcoma
 Myeloid proliferations related to Down's syndrome
 Transient abnormal myelopoiesis
 AML associated with Down's syndrome
 Blastic plasmacytoid dendritic cell neoplasm
 Acute leukaemias of ambiguous lineage
 Acute undifferentiated leukaemia
 Acute biphenotypic leukaemia

Precursor lymphoid neoplasms

B-lymphoblastic leukaemia/lymphoma
 B-lymphoblastic leukaemia/lymphoma, not otherwise specified
 B-lymphoblastic leukaemia/lymphoma with recurrent
 cytogenetic/molecular genetic abnormalities
 B-lymphoblastic leukaemia/lymphoma with t(9;22)
 (q34;q11.2), *BCR-ABL1*
 B-lymphoblastic leukaemia/lymphoma with t(11;23), *MLL*
 rearranged
 B-lymphoblastic leukaemia/lymphoma with t(12;21)
 (p13;q22), *TEL-AML1 (ETV6-RUNX1)*
 B-lymphoblastic leukaemia/lymphoma with hyperdiploidy
 B-lymphoblastic leukaemia/lymphoma with hypodiploidy
 (hypodiploid ALL)
 B-lymphoblastic leukaemia/lymphoma with t(5;14)
 (q31;q32), *IL3-IGH*
 B-lymphoblastic leukaemia/lymphoma with t(1;19)
 (q23;p13.3), *E2A-PBX1 (TCF3-PBX1)*
 T-lymphoblastic leukaemia/lymphoma

Mature B-cell neoplasms

Chronic lymphocytic leukaemia/small lymphocytic lymphoma
 B-cell prolymphocytic leukaemia
 Splenic marginal zone lymphoma
 Hairy cell leukaemia
 Splenic lymphoma/leukaemia, unclassifiable
 Splenic diffuse red pulp small B-cell lymphoma*
 Hairy cell leukaemia variant*
 Lymphoplasmacytic lymphoma
 Waldenström's macroglobulinaemia
 Heavy-chain diseases
 α Heavy-chain disease
 γ Heavy-chain disease
 μ Heavy-chain disease
 Plasma cell myeloma
 Solitary plasmacytoma of bone
 Extraosseous plasmacytoma
 Extranodal marginal zone B-cell lymphoma of mucosa-associated lymphoid tissue (MALT lymphoma)
 Nodal marginal zone B-cell lymphoma
 Paediatric type nodal marginal zone lymphoma
 Follicular lymphoma
 Paediatric type follicular lymphoma
 Primary cutaneous follicle centre lymphoma
 Mantle cell lymphoma
 Diffuse large B-cell lymphoma (DLBCL), not otherwise specified
 T-cell/histiocyte-rich large B-cell lymphoma
 DLBCL associated with chronic inflammation
 Epstein–Barr virus (EBV)-positive DLBCL of the elderly
 Lymphomatoid granulomatosis
 Primary mediastinal (thymic) large B-cell lymphoma
 Intravascular large B-cell lymphoma
 Primary cutaneous DLBCL, leg type
 Anaplastic lymphoma kinase (ALK) positive large B-cell lymphoma
 Plasmablastic lymphoma
 Primary effusion lymphoma
 Large B-cell lymphoma arising in HHV8-associated multicentric Castlemann disease
 Burkitt lymphoma
 B-cell lymphoma, unclassifiable, with features intermediate between DLBCL and Burkitt lymphoma
 B-cell lymphoma, unclassifiable, with features intermediate between DLBCL and classic Hodgkin lymphoma

Mature T-cell and NK-cell neoplasms

T-cell prolymphocytic leukaemia
 T-cell large granular lymphocytic leukaemia
 Aggressive natural killer (NK) cell leukaemia
 Systemic EBV-positive T-cell lymphoproliferative disease of childhood (associated with chronic active EBV infection)
 Hydroa vaccineforme-like lymphoma

Adult T-cell leukaemia/lymphoma
 Extranodal NK/T-cell lymphoma, nasal type
 Enteropathy-associated T-cell lymphoma
 Hepatosplenic T-cell lymphoma
 Subcutaneous panniculitis-like T-cell lymphoma
 Mycosis fungoides
 Sézary syndrome
 Primary cutaneous anaplastic large-cell lymphoma
 Primary cutaneous aggressive epidermotropic CD8-positive cytotoxic T-cell lymphoma*
 Primary cutaneous $\gamma\delta$ T-cell lymphoma
 Primary cutaneous small/medium CD4-positive T-cell lymphoma*
 Peripheral T-cell lymphoma, not otherwise specified
 Angioimmunoblastic T-cell lymphoma
 Anaplastic large-cell lymphoma, ALK positive
 Anaplastic large-cell lymphoma, ALK negative*

Hodgkin lymphoma

Nodular lymphocyte-predominant Hodgkin lymphoma
 Classic Hodgkin lymphoma
 Nodular sclerosis classic Hodgkin lymphoma
 Lymphocyte-rich classic Hodgkin lymphoma
 Mixed cellularity classic Hodgkin lymphoma
 Lymphocyte-depleted classic Hodgkin lymphoma

Histiocytic and dendritic cell neoplasms

Histiocytic sarcoma
 Langerhans' cell histiocytosis
 Langerhans' cell sarcoma
 Interdigitating dendritic cell sarcoma
 Follicular dendritic cell sarcoma
 Dendritic cell tumour, not otherwise specified
 Indeterminate dendritic cell tumour
 Fibroblastic reticular cell tumour

Post-transplant lymphoproliferative disorders

Early lesions
 Reactive plasmacytic hyperplasia
 Infectious mononucleosis-like
 Polymorphic post-transplant lymphoproliferative disorder
 Monomorphic post-transplant lymphoproliferative disorder (B and T/NK cell types)*
 Classic Hodgkin lymphoma-type post-transplant lymphoproliferative disorder

*These represent provisional entities or provisional subtypes of other neoplasms. They are provisional because there are insufficient data to support their being a definite entity, significant controversies about their defining features and/or uncertainty about whether they are unique or closely related to other definite entities.

Index

Page numbers in *italics* denote figures, those in **bold** denote tables.

- A blood group **336**
- AB blood group **336**
- abciximab **319**, 320
- ABO blood group system 334–5, **336**, **336**, **337**
 - antibodies 334, **336**
 - haemolytic disease of newborn 352
 - and venous thrombosis 306
- ABVD regimen 211
- acanthocytes 23
- acanthocytosis 325
- aciclovir 259
- acquired haemolytic anaemias 68–71
 - chemical and physical agents 71
 - immune 68–70, **68**, **69**, **70**
 - infections 71
 - march haemoglobinuria 70
 - red cell fragmentation syndromes 70, **70**
- actin 19
- activated partial thromboplastin time (aPTT) 276, **276**
 - coagulation disorders **294**, **299**
- activated protein C (APC) 275, 275, 304
- activation domain 8
- acute hyperviscosity syndrome 221
- acute lymphoblastic leukaemia (ALL) 186–96
 - BCR-ABL1 positive 194–5
 - classification 188, **188**, **189**
 - clinical features 188, **188**, **189**
 - CNS-directed therapy 193
 - cytogenetics/molecular genetics 189–91, **190**, **191**
 - immunological markers **189**
 - incidence and pathogenesis 187, **187**
 - intensification (consolidation) therapy 192–3
 - investigations 188–9, **189**, **189**
 - maintenance therapy 193–4
 - minimal residual disease 192, **193**
 - prenatal origin **187**
 - prognosis **194**, **196**, **195**
 - relapse **193**, **194**
 - remission induction 192
 - specific therapy 192, **192**, **194**
 - supportive therapy 192
 - toxicity of therapy 194
- acute myeloid leukaemia (AML) 145–55
 - blood film **150**
 - classification **146**, 147–8
 - clinical features 148, **148**, **149**
 - cytogenetics/molecular genetics **146**, 149, **151**, **151**, **152**
 - incidence 147
 - investigations 148–9, **149**
 - older patients 153
 - outcome 154, **155**
 - pathogenesis 147, **147**
 - prognosis **151**, 153
 - relapse 154
 - stem cell therapy 153
 - treatment 149, 151–3, **152–4**
- acute promyelocytic leukaemia 149
 - treatment 152–3
- ADAMTS13 267, 285, **285**, **286**
- adaptive immune system 88
- adhesion molecules 8
- adriamycin **142**
 - BEACOPP regimen 211
- adult T-cell leukaemia/lymphoma 203, 204, **226**
- African iron overload 43
- age effects *see* older patients
- AIDS *see* HIV/AIDS
- Alder's anomaly 93
- alemtuzumab **143**
 - aplastic anaemia 246
 - CLL 201
- alkylating agents 140, **141**, **142**
 - see also individual drugs*
- ALL *see* acute lymphoblastic leukaemia
- all-trans retinoic acid (ATRA) **141**, 144, 152
 - ATRA syndrome 153
- allele-specific priming 85
- allergic reaction to blood transfusion 342
- allogeneic stem cell transplantation 236, 255–62
 - chimaerism analysis 256
 - complications 256–61, **258**
 - donor leucocyte infusions 261, **262**
 - graft failure 260
 - graft-versus-leukaemia effect 261, **261**
 - HLA system 255–6, **255**, **256**
 - post-transplant lymphoproliferative diseases 261–2, **262**
- alloimmune haemolytic anaemias 70
- alternative complement pathway 109, **109**
- amegakaryocytic thrombocytopenia 244
- aminocaproic acid 280
- γ -aminolaevulinic acid 16, 29
- AML *see* acute myeloid leukaemia
- amplification enzyme system 109
- amplifications 130
- amyloidosis 23–9, **239**, **239**
- anaemia 19–25, **20**
 - aplastic 91, 243–7
 - blood film 22–3, **23**
 - bone marrow in 23–4, **24**
 - of chronic disorders 37, **37**, **38**, **322**, **322**
 - classification **22**
 - clinical features 20–1, **21**
 - congenital dyserythropoietic 249
 - global incidence 19–20
 - haemolytic *see* haemolytic anaemias
 - HIV/AIDS 328
 - hypochromic 27–40, **28**
 - ineffective erythropoiesis 24–5, **25**
 - iron deficiency 32–7
 - laboratory findings 33, **38**
 - lead poisoning 39
 - leucocyte and platelet counts 22
 - macrocytic 49, 58–9, **58**
 - malignant disease 322, **323**
 - megaloblastic *see* megaloblastic anaemia
 - microangiopathic haemolytic 70, **322**
 - neonate 349–50, **349**
 - older patients 322
 - pernicious 52–3, **53**
 - pregnancy 347, **347**
 - of prematurity 350
 - red cell indices 21
 - refractory **178**
 - renal failure 325, **325**
 - sideroblastic 38–9, **38**, **38**, **39**
 - see also specific types*
- anaplastic large cell lymphoma 226
- androgens in aplastic anaemia 246
- aneuploid cells 127
- angioimmunoblastic lymphadenopathy 226
- angular cheilosis **34**, **54**
- ankyrin 19
- anthracyclines **141**, **142**

- anti-emetics in haematological malignancies 137
- antibiotics 139
 - cytotoxic **142**, 143
- antibody-dependent cell-mediated cytotoxicity 106
- anticardiolipin antibodies **307**
- anticoagulants 312–18, **312**
 - direct-acting parenteral 315
 - heparin **299**, 312–15, **313**
 - oral 315–18, **315–17**
 - vitamin K antagonists 315–18, **315–17***see also specific drugs*
- antigen-presenting cells (APCs) 109
- antigen-receptor gene rearrangements 107, **107**, **108**
- antigenic specificity 103
- antiglobulin test 65, **340**, **340**
- antihaemophilic factor *see* factor VIII
- antihuman globulin 340
- antimetabolites 141, **141**, **142**, 143
- antiphospholipid syndrome 307
- antiplatelet drugs 287, 318–20, **319**, **319**
- antithrombin deficiency 305
- antithymocyte globulin 246
- antiviral drugs 139
- apixaban **317**, 318
- aplastic anaemia 91, 243–7, **243**
 - causes **243**
 - clinical features 245–6, **245**
 - congenital (Fanconi anaemia) 244, **244**
 - diagnosis 246
 - idiopathic acquired 244
 - laboratory findings 246
 - pathogenesis 244
 - secondary causes 245
 - treatment 246–7
- apoptosis 9–10, **9**
- argatroban 315
- arsenic **143**, 144
- arterial thrombosis 303, **303**, **303**
- Ashwell-Morell receptor 266
- asparaginase **141**, **143**, 144
- aspergillosis 139–40, **139**, **140**
- aspirin 319, **319**
 - platelet function disorders 287, **288**
 - polycythaemia vera 171
- atherosclerosis *see* arterial thrombosis
- ATRA syndrome 153
- atrophic gastritis 33, 35, 37, 53
- Auer rods **150**
- autoimmune gastritis 35, 52
- autoimmune haemolytic anaemias 68–70, **69**
 - classification **68**
 - cold 69–70
 - warm 68–9, **69**
- autoimmune (idiopathic) thrombocytopenic purpura 282–3, **282**
 - see also* idiopathic thrombocytopenic purpura
- autologous blood transfusion 343–4
- autologous stem cell transplantation 254, **255**
- automated blood cell counter **12**
- autotransfusion 343–4
- azacytidine **142**
 - myelodysplasia 184
- azathioprine **141**
- B** blood group **336**
- B cells 103–4
 - functional aspects **105**
 - immunodeficiency **113**
- B-cell chronic lymphoid leukaemias 198–203
- B-cell prolymphocytic leukaemia 202, **202**
- B-cell receptor 104, **105**
- bacterial infections
 - in cancer patients 138–9
 - haematological changes 327–8
 - and haematological malignancies 125
 - transfusion-related **339**, **342**
- band neutrophils 89, **89**
- Barr bodies 94
- basket cells 23
- basophil leucocytosis (basophilia) 97
- basophils **12**, 88, 88, 90
 - blood count **20**, **88**
- BCL-2 **9**
- BCR-ABL inhibitors **142**, 143
- BCR-ABL1 mutation 161, 187, 194–5
- BEACOPP regimen 211
- Bence-Jones protein 232
- bendamustine **142**
- benign ethnic neutropenia 95
- Bernard-Soulier syndrome 266, 287
- bilirubin 61
- biopsy of bone marrow 24, **24**, **25**
- bisphosphonates 237
- bivalirudin 315
- blackwater fever 329
- bleeding disorders 278–89
 - abnormal bleeding 279, **279**
 - thrombocytopenia *see* thrombocytopenia
 - vascular 279–80, **279**
- bleeding time 277
- bleomycin **141**, **142**, 143
 - BEACOPP regimen 211
- blood cells 12, **12**, **12**, **13**
 - count *see* blood count
 - see also* red cells; white cells
- blood coagulation *see* coagulation
- blood components 343–5, **343**
- blood count **20**, **88**, **276**
 - see also specific cell types*
- blood disorders and venous thrombosis 307
- blood donors 334, **335**
 - see also* blood transfusion
- blood film 276
 - acute myeloid leukaemia **150**
 - adult T-cell leukaemia/lymphoma **203**
 - ALL **190**
 - anaemia 22–3, **23**
 - autoimmune haemolytic anaemia **69**
 - B-cell prolymphocytic leukaemia **202**
 - β-thalassaemia major **79**
 - CLL **199**
- blood film CML, **159**, **163**
 - diffuse large B-cell lymphoma **224**
 - eosinophilic leucocytosis **97**
 - G6PD deficiency **67**
 - hairy cell leukaemia **203**
 - hereditary spherocytosis **65**
 - iron deficiency 34, **34**, **35**
 - large granular lymphocytic leukaemia **203**
 - liver disease **325**
 - lymphocytosis **112**
 - malignant disease **323**, **324**
 - mantle cell lymphoma **223**
 - megaloblastic anaemia **55**
 - microangiopathic haemolytic anaemia **70**
 - multiple myeloma **235**
 - myelodysplasia **183**
 - myelofibrosis **174**
 - non-Hodgkin lymphoma **218**
 - renal failure **325**
 - sickle cell anaemia **84**
 - splenic atrophy **117**
 - thrombocythaemia **173**
 - thrombotic thrombocytopenic purpura **286**
- blood groups
 - antibodies 334, **336**
 - determination of **340**, **340**
 - serology 339–40
 - see also individual blood groups*
- blood loss *see* haemorrhage
- blood products
 - haematological malignancies 136–7
 - irradiated **342**
 - post-stem cell transplant 259
 - reduction of use 342–3
 - see also* blood transfusion

- blood salvage 344
- blood transfusion 333–45, 334
 - autologous 343–4
 - bacterial contamination 342
 - blood group serology 339–40
 - complications 340–2, **341**
 - cross-matching and pre-transfusion tests 340, 340, **341**
 - hazards of 338–9, **338**, **339**
 - iron overload 44, 44, 45, 342
 - massive transfusion syndrome 287
 - post-transfusion purpura 284, 342
 - protection of recipient **338**
- bone morphogenetic protein (BMP) 30
- bone marrow 2, 2
 - ALL 188
 - anaemia 23–4, 24
 - biopsy 24, 24, **25**
 - failure 245
 - HIV/AIDS 329
 - iron stores 34, 35
 - myelodysplasia 180
 - pluripotent stem cells 3
 - red cell aplasia 247
 - stroma 4, 4
 - transplantation 251
 - trepine biopsy 24, 24, **25**
- Borrelia burgdorferi* **339**
- bortezomib 141, 143, **143**
 - mantle cell lymphoma 223
 - multiple myeloma 237
 - VMP regimen 235
- bosutinib **142**, **160**
- bridging anticoagulation 316–17
- Brucella melitensis* **339**
- Burkitt lymphoma 224–5, 225
 - cytogenetics **217**
 - immunophenotype **217**
- burr cells 325
- C-reactive protein 332
- C282Y mutation 43
- CALR (calreticulin) mutation 166, 167, **172**
- Candida* infections in cancer patients 140
- CAR (chimeric antigen receptors) 194
- carfilzomib **143**
 - multiple myeloma 235, 237
- caspases 9
- CD2 **189**
- CD3 **189**
- CD7 **189**
- CD10 **189**
- CD11c **147**
- CD13 **147**
- CD14 **147**
- CD19 **189**, 194
- CD20 antibodies 201
- CD22 **189**
- CD33 **147**
- CD39 270
- CD41 **147**
- CD42 **147**
- CD52 antibodies 201
- CD61 **147**
- CD64 **147**
- CD79a **189**
- CD117 **147**
- Cdks 8
- CDT regimen 235
- cell cycle 8
- central nervous system lymphoma 224, 225
- central venous catheters 136, 136
- centroblasts 111
- centromere 128, 129
- cerebral lymphoma 224, 225
- checkpoints 8
- Chédiak-Higashi syndrome 93, 94, 95
- chelation therapy 45–7, 46
- chemical exposure
 - acquired haemolytic anaemias 71
 - haematological malignancies 124
- chemokines 92–3
- chemotaxis 92
 - defects 93
- children
 - blood count **88**
 - Burkitt lymphoma 224–5, 225
 - haemolytic uraemic syndrome 286
 - iron requirements **33**
 - juvenile myelomonocytic leukaemia 184
 - myelodysplasia **178**
 - sites of haemopoiesis **2**
 - see also* neonates
- chimaerism 256
- chimeric antigen receptors (CAR) 194
- chlorambucil **142**
- 2-chlorodeoxyadenosine **142**
- CHOP regimen 223
- Christmas disease *see* haemophilia B (Christmas disease)
- Christmas factor *see* factor IX
- chromosomes 127–9, 128, 129
- chronic disorders, anaemia of 37, **37**, **38**, 322, **322**
- chronic eosinophilic leukaemia 164
- chronic lymphocytic leukaemia (CLL) 198–202
 - classification **198**
 - clinical features 198–9, 198, 199
 - course 202
 - cytogenetics **217**
 - immunophenotype **200**, **217**
 - laboratory findings 199–200, 199
 - molecular tests 200, **200**
 - pathogenesis 198
 - prognosis **202**
 - staging 200, **201**
 - treatment 200–2
- chronic lymphoid leukaemias 197–204
 - B-cell 198–203
 - diagnosis 198
 - T-cell 203–4
 - see also specific types*
- chronic myeloid leukaemia (CML) 156–64, **157**
 - accelerated phase disease 162–3
 - atypical 184
 - clinical features 159
 - genetic mutations **166**
 - laboratory findings 159, 159
 - Philadelphia chromosome 157–8
 - prognostic scores 159
 - stem cell therapy 162, 163
 - treatment 160–2, 160–3, **160**, **162**
- chronic myelomonocytic leukaemia 180, 184
- chronic neutrophilic leukaemia 164
- ciclosporin
 - aplastic anaemia 246
 - CLL 201
- cidofovir 259
- classical complement pathway 109, 109
- clg **189**
- CLL *see* chronic lymphocytic leukaemia
- clofarabine **142**
- clonal abnormalities **128**
- clonal progression 125, 127, 127
- clopidogrel 287, 288, **319**, 320
- Clostridium perfringens* 328, 328
- CNS *see* central nervous system
- coagulation 270–3
 - amplification 272–3, 272
 - in vivo* 270–1
 - initiation 271
 - physiological limitation 274
 - see also* haemostasis
- coagulation cascade 270, 271
- coagulation disorders 290–301
 - acquired 296–301, **296**
 - circulating antibodies 300–1
 - disseminated intravascular coagulation 287, 297–300, 298
 - haemophilia A 291–5

coagulation disorders (*continued*)

- haemophilia B (Christmas disease) **293**, **294**, 295
- hereditary 291–6, 291
- liver disease 297
- malignant disease 322–3, **324**
- massive transfusion syndrome 287, **299**, 301
- renal failure 326
- von Willebrand disease **294**, 295–6, **295**
- coagulation factors **271**, **273**
 - assays 277
 - circulating antibodies 300–1
 - inhibitors 275
 - vitamin K deficiency 296–7, 297
 - see also specific factors*
- cobalamin 49
- cold autoimmune haemolytic anaemias 69–70
- colony-stimulating factor (CSF) 90
- colorectal carcinoma 36
- complement 109, 109
- compression paraplegia 237
- compression ultrasound 308
- computed tomography *see* CT
- conditioning of stem cells 253
- congenital dyserythropoietic anaemia 249
- congestive heart failure 326
- conjunctival pallor 21
- connective tissue disorders 279
 - haematological changes 324
 - see also specific conditions*
- contrast venography 308
- Coombs' test 65, 340, 340
- cords of spleen 117, 117
- corticosteroids **142**, 143
 - idiopathic thrombocytopenic purpura 283
- coumarins *see* warfarin
- Coxiella burnettii* **339**
- CRAB acronym 231
- Creutzfeldt-Jacob disease 339, **339**
- cross-matching of blood 340, 340
- cryoprecipitate 344
- CT
 - multiple myeloma 233
 - non-Hodgkin lymphoma 219, 220
- CT pulmonary angiography 309, 309
- cyclin-dependent protein kinases *see* Cdk
- cyclins 8
- cyclophosphamide **142**
 - BEACOPP regimen 211
 - CHOP regimen 223
 - CLL 200
 - R-CVP regimen 222

cytogenetics

- ALL 189–91, 190, 191
- AML **146**, 149, 151, **151**, 152
- Burkitt lymphoma **217**
- CLL **217**
- diffuse large B-cell lymphoma **217**
- follicular lymphoma **217**
- hairy cell leukaemia **217**
- lymphoma **217**
- lymphoplasmacytic lymphoma **217**
- MALT lymphoma **217**
- mantle cell lymphoma **217**
- myelodysplasia **179**
- cytokinesis 8
- cytomegalovirus infection
 - haematological changes 328
 - post-stem cell transplant 259
 - transfusion-related 338
- cytosine arabinoside 141, **142**
- cytotoxic drugs 140–3, 141, **142–3**
- D-dimer 277
 - deep vein thrombosis 308
 - pulmonary embolus 309
- dabigatran **317**, 318
- dasatinib **142**, **160**
- daunorubicin **142**
- DDAVP 267, 293–4, 296
- decitabine **142**, 184
- deep vein thrombosis 308, **308**
 - prophylaxis 318
 - see also* thrombosis
- deferisirox 45, 46
- deferiptone 45–6, 46
- deferroxamine 46
- deletions 129, 130
- demethylation agents 141, **142**, 144
- dendritic cells 97, **98**, 109
 - neoplasms 226
- deoxycoformycin **142**
- deoxyhaemoglobin 17
- 1-diamino-8-D-arginine vasopressin *see* DDAVP
- Diamond-Blackfan syndrome 244, 247, **248**
- differentiation syndrome 153
- diffuse large B-cell lymphoma 223–4, 224
 - cytogenetics **217**
 - immunophenotype **217**
- dihydrofolate reductase 51, 52
- diploid cells 127
- dipyridamole 287, 320
- direct antiglobulin test (DAT) 340
- disseminated intravascular coagulation 287, 297–300, 298

clinical features 298–9, 299

- laboratory findings 299, **299**
- pathogenesis 297–8, **298**
- treatment 299–300, **300**
- divalent metal transporter 1 *see* DMT-1
- DMT-1 29, 30
- DNA methylation 179
- DNA microarrays 131, 132
- DNA-binding domain 8
- Döhle bodies 93
- Donath-Landsteiner antibody 69–70
- donor leucocyte infusions 162, 253, 261, 262
- driver mutations 125, 147
- drug interactions, oral anticoagulants 315, **316**, **317**
- drug-induced disorders
 - aplastic anaemia **243**
 - haematological malignancies 124
 - immune haemolytic anaemias 70, 70
 - neutropenia 95
 - platelet function 287, 288
 - thrombocytopenia 281, **281**, 284, 284
- Duffy blood group system **336**
- duplications 129, 130
- dyskeratosis congenita 244
- Eastern Cooperative Oncology Group (ECOG) Performance Status 136, **136**
- echinocytes 23
- eculizumab 247
- edoxaban **317**
- Ehlers-Danlos syndrome 279
- electrocardiogram in pulmonary embolus 308
- elliptocytes 23
- elliptocytosis 19
 - hereditary **64**, 65, 65
- eltrombopag 246, 283, 284
- Embsen-Meyerhof pathway 18, 18
- defects 67–8
- endothelial cells 273, 274
- enteropathy-associated T-cell lymphomas 226
- eosin-5-maleimide 65
- eosinophils **12**, 88, 88, 90
 - blood count **20**, **88**
- eosinophilic granuloma 98
- eosinophilic leucocytosis (eosinophilia)
 - causes 96–7, 97, **97**
 - myeloid neoplasms **166**
- epigenetic changes 129, 130–1
- epigenetics 8–9, **8**, 130, 178
- Epstein-Barr virus (EBV) 111–12
 - antibody 113
 - post-stem cell transplant 259

- eptifibatide **319**, **320**
 erythroblasts **13**
 erythrocyte sedimentation rate **330**, **332**
 erythrocytes *see* red cells
 erythrocytosis (primary polycythaemia) **168**, **168**
 erythropoiesis **11–26**
 assessment of **25**
 ineffective **24–5**, **25**
 erythropoietin **5**, **6**, **13–16**, **15**, **326**
 clinical uses **16**
 and haemoglobin production **15**
 indications for therapy **16**
 thrombocythaemia **172**
 essential thrombocythaemia *see* thrombocythaemia
 etoposide **141**, **142**
 BEACOPP regimen **211**
ETV6-RUNX1 translocation **187**
 extramedullary haemopoiesis **118**
 extravascular haemolysis **63**
- 18F-fluorodeoxyglucose *see* FDG
 factor I **270**, **273**, **273**
 factor II **271**, **273**
 factor IIa inhibitors **317**
 factor III **271**, **272**, **273**, **275**
 factor V **271**, **273**
 deficiency **296**
 factor V Leiden mutation **304–5**, **305**
 factor VII **271**, **273**
 deficiency **296**
 freeze-dried concentrates **344**
 recombinant activated **294**
 factor VIII **271**, **273**, **293**
 autoantibodies **300**
 deficiency *see* haemophilia A
 inhibitors **294**
 raised, in venous thrombosis **306**
 recombinant **293**
 factor IX **271**, **273**
 deficiency *see* haemophilia B (Christmas disease)
 factor IX-prothrombin complex
 concentrates **344–5**
 factor X **271**, **273**
 deficiency **296**
 factor Xa inhibitors **317**
 factor XI **271**, **273**
 deficiency **296**
 factor XII **271**
 factor XIII **271**, **273**, **273**
 deficiency **296**
 familial amyloidosis **238**
 familial polycythaemia **75**
- Fanconi anaemia **244**, **244**, **245**
 FDG-PET in Hodgkin lymphoma **209–10**, **210**
 FEIBA **300**
 Felty's syndrome **324**
 ferriredutase **31**
 ferritin
 hyperferritinaemia **42**
 iron in **28**
 regulation **29**
 serum levels **20**, **35**, **36**
 ferroportin **31**, **32**
 fertility in cancer patients **137**
 fetomaternal alloimmune thrombocytopenia **350**
 fetus/fetal
 haemoglobin **16**, **73**, **74**, **81**
 sites of haemopoiesis **2**
 see also neonates
 fibrin (factor I) **270**, **273**, **273**
 fibrin-stabilizing factor *see* factor XIII
 fibrinogen **271**, **272–3**
 defects **306**
 quantitation **276**
 raised, in venous thrombosis **306**
 fibrinolysis **275–6**, **276**
 tests of **277**
 fibrinolytic agents **318**
 contraindications **319**
 Fitzgerald factor (high molecular weight kininogen) **271**
 Fletcher factor (prekallikrein) **271**
 flow cytometry **131**, **132**, **133**
 ALL **189**
 platelet function **277**
 fludarabine **141**, **142**
 CLL **200**
 fluorescence *in situ* hybridization (FISH)
 131, **131**
 FOG-1 **14**
 folate **50**, **51–2**
 absorption, transport and function **51**
 red cell **20**, **57**
 reduction **52**
 serum **20**, **57**
 structure **51**
 folate deficiency **53**
 causes **53**
 clinical features **53–6**, **54–6**, **54**
 diagnosis **56–7**, **57**
 pregnancy **347**
 tests for cause **57**, **57**
 treatment **57–8**, **57**, **58**
 follicular lymphoma **222–3**, **222**
 cytogenetics **217**
 immunophenotype **200**, **217**
- fondaparinux **315**
 foscarnet **259**
 fresh frozen plasma (FFP) **344**
 fungal infections in cancer patients **139–40**, **139**, **140**
- G1 phase **8**
 G2 phase **8**
 ganciclovir **259**
 gastritis
 atrophic **35**, **37**, **53**
 autoimmune **35**, **52**
 GATA-2 **4**, **14**
 Gaucher's disease **99**, **99**, **100**
 gene segments **107**, **107**
 gene sequencing **131**
 gene therapy in haemophilia A **294**
 genetic markers **133–4**, **133**, **134**
 genetics
 ALL **189–91**, **190**, **191**
 AML **146**, **149**, **151**, **151**, **152**
 CML **166**
 haematological malignancies **124–7**, **125–7**, **126**
 pre-implantation genetic diagnosis **85**
 germinal centre **111**
 giant cavernous haemangioma **279**
 Gilbert's disease **64**
 glandular fever *see* infectious mononucleosis
 Glanzmann's thrombasthenia **266**, **287**
 glossitis **32**, **54**
 glucose-6-phosphate dehydrogenase
 deficiency **66–7**, **66**, **67**
 glutathione deficiency **67**
 glycophorin **147**
 glycoprotein IIb/IIIa inhibitors **319**, **320**
 golf ball cells **77**
 graduated compression stockings **318**
 graft-versus-host disease **256**, **258**
 acute **257**, **258**, **258**
 chronic **258**
 post-transfusion **342**
 graft-versus-leukaemia effect **261**, **261**
 granulocyte colony-stimulating factor (G-CSF) **5**, **6**, **90**
 clinical uses **91**
 effects of **91**
 granulocyte-macrophage colony-stimulating factor (GM-CSF) **90**
 granulocytes **88**, **89–90**, **89**
 concentrates **344**
 see also specific types
 granulopoiesis **90–1**, **90**, **91**
 growth factor receptors **6–8**, **7**
 growth factors **4–6**, **5**, **5**, **6**

- haem 16, 30, 32
- haem enzymes **28**
- haemangioblasts 2
- haematinics, normal values **20**
- haematocrit **20**
- haematological malignancies 122–34, 123, 135–44
 - aetiology 124–5, 124
 - diagnostic methods 131–3, 131–3
 - ECOG status 136, **136**
 - genetic abnormalities associated with 129–31, 129, 130
 - genetic markers 133–4, 133, 134
 - genetics 124–7, 125–7, **126**
 - incidence 123, 123
 - specific therapies 140–4, 141, **142–3**
 - supportive therapy 136–40
 - see also specific therapies*
- haematoma **279**
- haematopoietin receptor superfamily 6
- haemochromatosis **42**, 43–4, 43
- haemoglobin 2, 13, 16–17, **16**, 73
 - and erythropoietin production 15
 - fetal-adult switch 74
 - function 17, 17
 - genetic disorders of 74–86, **75**
 - hereditary synthesis disorders 68
 - iron in **28**
 - molecular aspects 73–4, 73, 74
 - myelofibrosis **174**
 - normal values **20**, **20**
 - oxygen dissociation curve 17, 20
 - prenatal diagnosis of genetic disorders 85, 86
 - synthesis 16–17, 16, 17, 73
 - thrombocythaemia **172**
 - see also anaemia*
- haemoglobin A **16**, 73
- haemoglobin A2 **16**, 73
- haemoglobin Barts **75**, 76, 76
- haemoglobin C 80, 84, 85
- haemoglobin Constant Spring 76
- haemoglobin D 81, 85
- haemoglobin E 81, 85
- haemoglobin F **16**, 73, 81, 85
- haemoglobin H 77, 80
- haemoglobin Lepore 77, 81
- haemoglobin S 17, 80, 85
 - see also sickle cell anaemia*
- haemoglobinaemia 63
- haemoglobinuria 63, 64
 - march 70
 - paroxysmal cold 69–70
 - paroxysmal nocturnal 62, 246, 247, 247
- haemolysis **75**
- haemolytic anaemias 60–71
 - acquired 68–71
 - autoimmune *see* autoimmune haemolytic anaemias
 - classification 62, **62**
 - clinical features 62, 63
 - hereditary 64–8
 - intravascular/extravascular haemolysis 63, **63**, 64
 - laboratory findings 63
- haemolytic disease of newborn 350–2, 351, 352
 - ABO incompatibility 352
- haemolytic transfusion reactions 340–2, **341**
- haemolytic uraemic syndrome 285, 285, 286
 - Rh incompatibility 350–2, 351, 352
- haemophagocytic lymphohistiocytosis 98, 98
- haemophilia A 291–5
 - antenatal diagnosis 291
 - carrier detection 291
 - clinical features 291, 292, 293
 - disease severity **293**
 - gene therapy 294
 - laboratory findings 291, **294**
 - molecular genetics 291, 291, 292
 - prophylaxis 294
 - treatment 293–4
- haemophilia B (Christmas disease) 295
 - disease severity **293**
 - laboratory findings **294**, 295
- haemopoiesis 1–10
 - extramedullary 118
 - regulation of 4, 7, 91
 - sites of 2, 2, **2**
 - stem and progenitor cells 2–4, 3
- haemopoietic growth factors 4–6, 5–7, **5**, **6**
- aplastic anaemia 247
- haemopoietic progenitors 2, 3
- haemorrhage
 - massive 345
 - retinal 21
- haemorrhagic cystitis 260
- haemorrhagic disease of newborn 296
- haemosiderin 28
- haemosiderinuria 63, 247
- haemostasis 265, 273–5
 - disorders of 306
 - haematological malignancies 137
 - platelet plug 274–5
 - pregnancy 348
 - tests of 276–7, **276**
 - vasoconstriction 273–4
 - see also blood coagulation*
- Hageman (contact) factor *see* factor XII
- hairy cell leukaemia 202–3, 203
 - cytogenetics **217**
 - immunophenotype **200**, **217**
- Hand-Schüller-Christian disease 98
- haploid cells 127
- haploidentical stem cell transplantation 256
- haptoglobins 61
- Heinz bodies 24, 67, 71
- Helicobacter pylori* 33
 - chronic idiopathic thrombocytopenic purpura 282
 - haematological malignancies 125
 - MALT lymphoma 221–2
 - pernicious anaemia 52
- hemojuvelin 30, 43
- HEMPAS 249
- Henoch-Schönlein syndrome 280, 280
- heparin 312–15, **313**
 - administration and laboratory control 313–14
 - bleeding during therapy 314
 - haemostasis tests **299**
 - indications 313
 - low molecular weight **313**, 314
 - mode of action 313, 313
 - osteoporosis 315
 - unfractionated 313–14, **313**
- heparin-induced thrombocytopenia 314–15, 314
- hepatitis, transfusion-related 338, **339**
- hepcidin 29–30, 31, 43
- hereditary haemolytic anaemias 64–8
 - elliptocytosis **64**, 65, 65
 - glucose-6-phosphate dehydrogenase deficiency 66–7, 66, 67
 - South-East Asian ovalocytosis 65
 - spherocytosis 64–5, **64**, 65
- hereditary haemorrhagic telangiectasia 279, 280
- herpes viruses 139
 - post-stem cell transplant 259
 - transfusion-related **339**
- heterophile antibodies 112–13
- hexose monophosphate shunt 18
- HFE* gene 43
- high molecular weight kininogen (Fitzgerald factor) **271**
- histiocytes 97, **98**
- histiocytic cell neoplasms 226
- HIV/AIDS
 - cerebral lymphoma 224, 225
 - haematological changes 328, 329
 - transfusion-related 338, **339**

- HLA system 255–6, 255, **256**
 inheritance 257
 and stem cell transplantation 256
- HLA-restricted T cells 109
- Hodgkin lymphoma 205–12
 advanced stage 211
 clinical features 206, 206, 207
 diagnosis and histological classification
 207, 207, 208, **208**
 early stage 211
 FDG-PET 209–10, 210
 haematological and biochemical findings
 206–7
 history and pathogenesis 206
 late effects 212
 prognosis **210**, 212
 Reed-Sternberg cells 133, 206, 207
 relapse 211–12
 response to treatment 211
 staging 208–9, 209, **209**
 treatment 210–12, **210**, 211, 212
 WHO classification **208**
- Howell-Jolly bodies 24, 117, 120
- human albumin solution (4.5%) 344
- human albumin solution (20%) 344
- human leukocyte antigen *see* HLA
- human T-cell leukaemia virus, transfusion-related 338
- Hurler's syndrome 94
- hydrops fetalis **75**, 76, 76, 352, 352
- hydroxocobalamin **57**
- hydroxodaunorubicin, CHOP regimen 223
- hydroxycarbamide 85, 141, **142**
 polycythaemia vera 170, 171
- hydroxyurea *see* hydroxycarbamide
- hypercalcaemia 237
- hyperdiploid cells 127
 ALL 190, 191
- hyperferritinaemia **42**
- hyperglobulinaemia 287
- hyperhaemolysis syndrome 342
- hyperhomocysteinaemia 306
- hypersplenism 119–20, **120**
- hypertensive disorders, thrombocytopenia
 of 348
- hyperviscosity syndrome 240, 240
- hypochromic anaemias 27–40, 28
 anaemia of chronic disorders 37, **37**, **38**
 differential diagnosis **38**, 39
 iron deficiency *see* iron deficiency
 anaemia
 lead poisoning 39
 sideroblastic anaemia 38–9, 38, **39**
- hypodiploid cells 127
 ALL 190, 191
- hypomethylating agents 184
- hyposplenism 120, **120**
 infection prevention 121, **121**
- hypothyroidism 327
- ibritumomab **143**
- ibrutinib **106**, 141, **142**
 CLL 201
 mantle cell lymphoma 223
- idarubicin **142**
- idelasib **106**, **142**, 201
- idiopathic acquired aplastic anaemia 244
- idiopathic thrombocytopenic purpura (ITP)
 282–3
 acute 283
 chronic 282–3
 clinical features 282–3
 diagnosis 283
 pathogenesis 282, 282
 pregnancy 348
 treatment 283
 warm autoimmune haemolytic anaemia
 association 68–9, 69
see also autoimmune thrombocytopenic
 purpura
- Igs *see* immunoglobulins
- imatinib 141, **142**, 143, **160**
- immune function, splenic control 117
- immune haemolytic anaemias
 alloimmune 70
 autoimmune 68–70, 69
 drug-induced 70, 70
- immune response 109–11, 110, 111
- immunity
 adaptive 88
 innate 88
- immunodeficiency 113–14, **113**
see also HIV/AIDS
- immunoglobulins 106–7, 106, **106**
 gene rearrangements 107, 107, 108
 idiopathic thrombocytopenic purpura
 283
 IgA **106**
 IgG **106**
 IgM **106**
 isotypes 106
 pooled 345
 replacement 202
 subclasses 106
- immunohistology (immunocytochemistry)
 133, 133
- immunological markers **147**
- immunological memory 103
- immunoparesis 232
- immunosuppressants in idiopathic
 thrombocytopenic purpura 283
- indirect antiglobulin test (IAT) 340
- infections 327–30, **327**, 328–30
 bacterial 327–8
 cancer patients 138–9, 138
 haemolysis 71
 hyposplenic patients 121, **121**
 kala-azar (visceral leishmaniasis) 329,
 330
 malaria 329, 329
 parasitic 329, 330, 331
 post-stem cell transplant 258–60, 259,
 260
 and thrombocytopenia 283
 toxoplasmosis 329
 transfusion-related 338–9, **338**, **339**
 viral 328
see also HIV/AIDS
- infectious mononucleosis 111–12
- inferior vena cava filter 318
- inflammation, and venous thrombosis 306
- innate immune system 88
- interferon- α **143**, 144
 chronic myeloid leukaemia 162
 polycythaemia vera 171
- interferon- γ 5
- interleukins 5, **6**, 90
- intermittent compression devices 318
- international normalized ratio (INR) 315
- interphase 8
- interstitial pneumonitis 259, 260
- intravascular haemolysis 63, **63**, 64
- iron
 absorption 30–1, **30**, 31, 32
 body distribution/transport 28–30, **28**,
 29, 30
 daily cycle 29
 dietary 30
 oral 36–7, **37**
 parenteral 37
 requirements 31–2, **33**
 serum levels **20**, 34–5, 36
- iron chelation therapy 45–7, 46, 79
 aplastic anaemia 247
 myelodysplasia 182
- iron deficiency anaemia 32–7
 blood film 34, 34, 35
 causes 32–3, 33, **33**
 clinical features 32, 34
 investigations 35–6, 36
 laboratory findings **38**
 pregnancy 347
 red cell indices 34
 treatment 36–7, **37**

- iron overload 41–7
 - African 43
 - assessment 42, **42**, 43
 - causes **42**
 - chelation therapy 45–7, 46
 - haemochromatosis **42**, 43–4, 43
 - transfusional 44, 44, 45, 342
- iron refractory iron deficiency anaemia (IRIDA) 37
- iron regulatory protein 29
- iron response elements 29
- isochromosome 129
- ITP *see* idiopathic thrombocytopenic purpura
- JAK inhibitors 171
- JAK proteins 6, 8
- JAK/STAT pathway 6
- JAK2 mutation 166, 167, **169**, **172**
- Janus-associated kinase proteins *see* JAK proteins
- juvenile myelomonocytic leukaemia 184
- kala-azar (visceral leishmaniasis) 329, 330
- karyotype 127, 128
 - analysis 131
- Kell blood group system **336**, 337
- kernicterus 351
- Kidd blood group system **336**
- kidney
 - myeloma 235
 - renal failure 325–6, 325, **325**
- Kleihauer test 351, 351
- koilonychia 21, 32, 34
- Kupffer cells 275
- labile factor *see* factor V
- lactate dehydrogenase 286
- Langerhans' cell histiocytosis 98, 98, 226
- large granular lymphocytic leukaemia 203–4, 203
- lazy leucocyte syndrome 93
- lead poisoning 39
- lenalidomide **143**
 - CLL 201
 - mantle cell lymphoma 223
 - multiple myeloma 236, 237
 - myelodysplasia 182
- Lepore syndrome 77, 81
- lestaurtinib 171
- Letterer-Siwe disease 98
- leucocyte adhesion molecules 92
- leucocytes *see* white cells
- leucodepletion 343
- leucoerythroblastic reaction 94–5, 94, **95**
- leukaemia
 - acute lymphoblastic (ALL) 186–96
 - acute myeloid (AML) 145–55
 - acute promyelocytic 149, 152–3
 - chronic eosinophilic 164
 - chronic lymphoid 197–204
 - chronic myeloid (CML) 156–64, 184
 - chronic myelomonocytic 180, 184
 - chronic neutrophilic 164
 - classification 146, **146**
 - diagnosis 146–7, 146, **147**
 - vs. lymphoma 214
 - see also specific types*
- leukaemoid reaction 94, 322
- Lewis blood group system **336**, 337
- Li blood group system **336**
- liver disease 326–7, **326**
 - coagulation defects 297
 - haemostasis tests **299**
- Loa loa* 331
- low molecular weight heparin **313**, 314
- lupus anticoagulant 300–1
 - and venous thrombosis **307**
- Lutheran blood group system **336**
- lymph nodes 110
- lymphadenopathy 114, 114
- lymphoblastic lymphoma 225
- lymphocytes **12**, 88, 88, 102–15, 103
 - antigen receptors 105
 - B cells 103–4, 105, **105**
 - blood count **20**, **88**
 - circulation 106
 - natural killer cells 104, 106
 - T cells 103–4, 105, **105**
- lymphocytosis 111–13, **111**
 - clinical features 112
 - diagnosis 112–13, 112
 - differential diagnosis 113
 - heterophile antibodies 112–13
 - infectious mononucleosis 111–12
 - non-Hodgkin lymphoma 203
 - pleomorphic atypical 112, 112
 - treatment 113
- lymphoid organs 103, 104
 - secondary 110
- lymphoma
 - HIV/AIDS 224, 225
 - Hodgkin *see* Hodgkin lymphoma
 - non-Hodgkin *see* non-Hodgkin lymphoma
 - T-cell 225–6
 - vs. leukaemia 214
 - see also specific types*
- lymphopenia 113
- lymphoplasmacytic lymphoma
 - cytogenetics **217**
 - immunophenotype **217**
- lymphoplasmacytoid lymphoma 221, 221
- lymphoproliferative diseases, post-transplant 261–2, 262
- lysosomal storage diseases 99–101
 - Gaucher's disease 99, 99, 100
 - Niemann-Pick disease 99, 101
- M phase 8
- macrocytes 23
- macrocytic anaemia 49, 58–9, **58**
 - see also* megaloblastic anaemia
- macrophage colony-stimulating factor (M-CSF) 5, 6
- macrophages 92
- macrospherocytes 23
- magnetic resonance imaging *see* MRI
- major histocompatibility complex (MHC) 256, **256**
 - see also* HLA system
- malaria 329, 329
 - transfusion-related 338–9, **339**
- malignancy
 - dendritic cells 226
 - haematological 122–34, 123, 135–44
 - haematological changes 322–3, 323, 324, **324**
 - histiocytic cells 226
 - myelodysplastic/myeloproliferative 184, **184**
 - myeloid **166**
 - plasma cells **231**
 - and venous thrombosis 306
 - see also named diseases and tumours*
- malignant disease *see* neoplasms
- MALT lymphoma 221–2, 222
 - cytogenetics **217**
 - immunophenotype **217**
- mantle cell lymphoma 223, 223
 - cytogenetics **217**
 - immunophenotype **200**, **217**
- MAP kinase 6
- march haemoglobinuria 70
- marginal zone lymphomas 221–2, 222
- massive haemorrhage 345
- massive transfusion syndrome 287, 301
 - haemostasis tests **299**
- mastocytosis 175, 175
 - genetic mutations **166**
- matriptase 2 30
- May-Hegglin syndrome 93, 94, 281
- mean cell haemoglobin (MCH) **20**
- mean cell volume (MCV) **20**, 21
- megakaryocytes 182, 265–6, 266, 267

- megaloblastic anaemia 49, **49**, 93
 - biochemical basis 51, 52
 - clinical features 53–6, 54–6, **54**
 - laboratory findings 55–6, 55, 56
 - treatment 57–8, **57**, 58
 - see also* folate; vitamin B12
- melphalan **142**
 - MTP regimen 235
 - multiple myeloma 235
- men, iron requirements **33**
- menorrhagia 33
- 6-mercaptopurine 141, **142**
- mesenchymal stem cells 4
- methaemalbuminaemia 63
- methaemoglobinaemia 17, **75**
- methotrexate 141, **142**
- methylenetetrahydrofolate reductase 306
- MGUS 229, **231**, 232, 237–8, 238
- microangiopathic haemolytic anaemia 70, 322
- microcytes 23
- microRNAs 131
- minor histocompatibility antigens 256
- mitogen-activated protein kinase *see* MAP kinase
- mitosis 8
- mitoxantrone **142**
- mixed phenotype acute leukaemia 148
- MN blood group system **336**, 337
- molecular genetics
 - ALL 189–91, 190, 191
 - AML **146**, 149, 151, **151**, 152
 - CLL 200
- mometinib 171
- monoclonal antibodies 141, **143**
 - idiopathic thrombocytopenic purpura 283
 - non-Hodgkin lymphoma 220
 - see also* named antibodies
- monoclonal B-cell lymphocytosis (MBL) 198
- monoclonal gammopathy of undetermined significance *see* MGUS
- monoclonal immunoglobulins, associated diseases **229**
- monocyte colony-stimulating factor (M-CSF) 90
- monocytes **12**, 88, 88, 89, 89, 92
 - blood count **20**, **88**
 - disorders of function 92–4, 92, 93
- monocytosis 96, **97**
- monospot test 112
- MR pulmonary angiography 309
- MRI
 - cerebral lymphoma 224
 - deep vein thrombosis 308
- haemophilia A 292
 - non-Hodgkin lymphoma 219
- mucosa-associated lymphoid tissue lymphoma *see* MALT lymphoma
- mucosal bleeding **279**
- multiple myeloma 229–37
 - clinical features 231–2, 232–5, 239
 - diagnosis 231
 - pathogenesis 229, 230, **231**
 - prognosis 237
 - relapse 237
 - smouldering 229, 231, **231**
 - supportive care 237
 - treatment 235, 233–7, 236
- Mycoplasma pneumoniae* 328
- mycosis fungoides 226, 226
- myeloablative conditioning 253
- myeloblasts 89, 89
- myelodysplasia 91, 177–86
 - classification **178**, 179–80
 - clinical features 180, 182
 - cytogenetic abnormalities **179**
 - diagnosis 181
 - genetic abnormalities 182
 - high-risk syndromes 184
 - laboratory findings 180, 183
 - low-risk syndromes 182
 - prognostic score **179**, **180**
 - treatment 182, 184
- myelodysplastic syndromes *see* myelodysplasia
- myelodysplastic/myeloproliferative neoplasms 184, **184**
- myelofibrosis 173–5
 - clinical features 174
 - genetic mutations **166**, 167
 - laboratory findings 174, 174
 - survival score **174**
 - treatment 174–5
- myeloid growth factors 90–1, 91
- myeloma *see* multiple myeloma
- myeloma kidney 235
- myelomonocytic leukaemia
 - chronic 180, 184
 - juvenile 184
- myeloproliferative disease 165–76, 166, **166**, 287
 - JAK2 mutation 166, 167
 - mastocytosis **166**, 175, 175
 - myelofibrosis **166**, 167, 173–5
 - polycythaemia 168–72
 - thrombocythaemia **166**, 167, 172–3, **172**, 173, **173**
- Mylotarg® **143**
- myoglobin, iron in **28**
- nail bed pallor 21
- natural killer cells 104, 106
- neonates 348–52
 - anaemia 349–50, 349
 - blood coagulation 350
 - blood count **88**, 348–9, 349
 - fetomaternal alloimmune thrombocytopenia 350
 - haemolytic disease of newborn 350–2, 351, 352
 - haemorrhagic disease of newborn 296
 - hydrops fetalis **75**, 76, 76, 352, 352
 - polycythaemia 350
- neoplasms *see* malignancy
- neural tube defects 55, 55
- neutropenia 91, 95–6
 - autoimmune 96
 - benign ethnic 95
 - causes **95**
 - clinical features 96, 96
 - congenital 95
 - cyclical 95
 - diagnosis 96
 - drug-induced 95
 - HIV/AIDS 328
 - idiopathic benign 96
 - management 96
 - refractory **178**
- neutrophil leucocytosis 93, 94–5, 94, **95**
 - causes **94**
- neutrophils **12**, 88, 88, 89, 89
 - blood count **20**, **88**
 - disorders of function 92–4, 92, 93
 - precursors 89, 89
- Niemann-Pick disease 99, 101
- nilotinib **142**, 143, **160**
- nitric oxide 270
- non-Hodgkin lymphoma 213–27
 - cell of origin 214, 216
 - cellular origins 215
 - classification 214, **214**, 215
 - clinical features 216
 - cytogenetics/genetic analysis 217–18, **217**, 218
 - high-grade 214, 223–5
 - histology 216, 217
 - HIV/AIDS 328
 - imaging 219, 220
 - laboratory investigations 216–17
 - low-grade 214, 220–3
 - lymphocytosis 203
 - pathogenesis 216, **216**
 - prognostic index **218**
 - staging 218
 - T-cell lymphomas 225–6

- non-Hodgkin lymphoma (*continued*)
 treatment 220
see also specific subtypes
- non-myeloablative conditioning 253
- NOTCH signalling pathway 188
 ALL 191
- NOTCH-1 4
- nutritional support of cancer patients 137–8
- O blood group **336**
- obinutuzumab **143**, 201
- oestrogen therapy, and venous thrombosis 307
- ofatumumab **143**, 201
- older patients
 acute myeloid leukaemia 153
 haematological changes 322
- oncogenes 125
- oral anticoagulants 315–18, **315–17**
 bridging anticoagulation 316–17
 direct acting 317–18, **317**
 drug interactions 315, **316**
 INR 315
 length of anticoagulation 315
 overdose 316, **316**
see also specific drugs
- osteopetrosis 249
 haematological changes 330
- osteoporosis, heparin-induced 315
- osteosclerotic myeloma (POEMS syndrome)
 237
- ovalocytosis, South-East Asian 65
- oxyhaemoglobin 17
- P blood group system **336**, 337
- packed cell volume (PCV) **20**
- pacritinib 171
- pain in haematological malignancies 138
- pancytopenia 243, **243**
 malignant disease **324**
- Pappenheimer bodies 24, 117, 120
- paraproteinaemia 107, 229, 229
- parasitic infections 329, 330, 331
 transfusion-related **339**
- paroxysmal cold haemoglobinuria 69–70
- paroxysmal nocturnal haemoglobinuria 62,
 246, 247, 247
- parvovirus B19 248
 transfusion-related **339**
- passenger mutations 125
- Pearson's syndrome 38
- PD-1 112
- Pelger abnormality 180, 183
- Pelger-Huët anomaly 93, 94
- pencil cells 23
- pentose phosphate shunt 18
- periarteriolar lymphatic sheath 117, 117
- peripheral blood stem cell (PBSC)
 transplantation 251
- peripheral T-cell non-Hodgkin lymphoma,
 unspecified 225
- pernicious anaemia 52–3
 associations **53**
- PET
 Hodgkin lymphoma 209–10, 210
 multiple myeloma 233
 non-Hodgkin lymphoma 219, 220
- petechiae **279**
- PFA-100 test 277
 coagulation disorders **294**
- phagocytes **12**, 88
- phagocytosis 92–3, 92
 defects 93
 killing and digestion 93
- Philadelphia (Ph) chromosome 157–8, 191
- phosphatidylinositol 3 kinase *see* PI3 kinase
- PI3 kinase 6, 8
- PIVKA factors 296
- plasma cells 103
 neoplasms **231**
- plasma cell leukaemia 237
- plasma thromboplastin antecedent *see* factor
 XI
- plasma viscosity 332
- plasma volume **168**
- plasmin 275
 inactivation 276
- plasminogen 275
- platelets **12**, 265–70
 adhesion 269
 aggregation 267–8, **294**
 amplification 268
 anaemia 22
 antigens **147**, 266
 concentrates 344
 destruction 282–7
 failed production 281, **281**
 function 267–70
 inhibitors of function 270
 malignant disease 322–3, **324**
 plug 274–5
 procoagulant activity 268
 production 265–6, 266, 267
 release reaction 268
 structure 266, 269
 transfusion 283, 289
- platelet count **20**, 266
 coagulation disorders **294**, **299**
 myelofibrosis **174**
 raised **173**
 thrombocythaemia **172**
- platelet function disorders 287–9
 Bernard-Soulier syndrome 266, 287
 diagnosis 288–9, 288
 drug-induced 287, 288
 Glanzmann's thrombasthenia 266, 287
 hyperglobulinaemia 287
 renal failure 326
 storage pool diseases 287
 uraemia 287
see also myeloproliferative disease
- platelet function tests 277
- platelet-derived growth factor (PDGF) 270,
 303
- platinum derivatives 144
- pleomorphic atypical lymphocytosis 112,
 112
- pletixafor 251
- pluripotent stem cells 2–4, 3
- Pneumocystis carinii* 259, 260
- POEMS syndrome (osteosclerotic myeloma)
 237
- point mutations 129, 129
- polycythaemia 168–72
 apparent 172
 classification 168, **168**
 differential diagnosis 172
 genetic mutations **166**, 167
 malignant disease 322
 neonate 350
 primary (erythrocytosis) 168, **168**
 secondary **168**, 172
- polycythaemia vera 168–71
 clinical features 169–70, 169, 170
 congenital causes 171
 course and prognosis 171
 diagnosis 169, **169**
 genetic mutations **166**
 laboratory findings 170, 170
 treatment 170–1, 171
- polymerase chain reaction (PCR) 134, 134
- pomalidomide **143**
 multiple myeloma 235, 237
- ponatinib **160**
- positron emission tomography *see* PET
- post-capillary venules 106
- post-thrombotic syndrome 318
- post-transfusion circulatory overload 342
- post-transfusion purpura 284, 342
- post-transfusional iron overload 44, 44, 45,
 342
- post-transplant lymphoproliferative diseases
 261–2, 262
- postoperative venous thrombosis 306
- prasugrel 287, 320
- pre-implantation genetic diagnosis 85

- prednisolone
 - ALL 189
 - autoimmune haemolytic anaemia 68, 201
 - BEACOPP regimen 211
 - CHOP regimen 223
 - ITP 283, 284
 - MTP regimen 235
 - R-CVP regimen 222
- pregnancy 347–8, 347, 348
 - anaemia 347, 347
 - folate/vitamin B12 deficiency 347
 - haemostasis and thrombosis 348
 - thrombocytopenia 347–8, 348
- prekallikrein (Fletcher factor) 271
- prenatal diagnosis in haemoglobin disorders 85, 86
- primary cold agglutinin disease 69–70
- prion diseases, transfusion-related 339, 339
- procabazine, BEACOPP regimen 211
- proconvertin *see* factor VII
- progenitor cells 2–4, 3, 12–13
- prolymphocytic leukaemia
 - B-cell 202, 202
 - T-cell 203
- promyelocytes 89, 89
- prostacyclin 268, 270, 270
- protein 4.1 19
- protein C 275, 275
 - concentrate 345
 - deficiency 305
- protein S 275
 - deficiency 305–6
- prothrombin allele G20210A 306
- prothrombin (factor II) 271, 273
- prothrombin time 276, 276
 - coagulation disorders 294, 299
- proto-oncogenes 125, 125
- protozoal infections
 - and haematological malignancies 125
 - transfusion-related 339
 - see also* malaria
- pseudodiploid cells 127
- pseudopolycythaemia 168
- psychological support of cancer patients 137
- pulmonary angiography 309
- pulmonary embolus 308–9, 309
 - prophylaxis 318
- purpura 280, 280, 281
 - post-transfusion 284, 342
 - thrombotic thrombocytopenic 285, 285, 286
- pyruvate kinase deficiency 67–8
- 5q-syndrome 180
- R-CHOP regimen 223
- R-CVP regimen 222–3
- R-FC regimen 200
- R-ICE regimen 224
- radiation-induced disorders 124
- radiography
 - ALL 189
 - haemophilia A 292
 - interstitial pneumonitis 260
 - multiple myeloma 233, 235
 - pulmonary embolus 308
- radionuclide scanning 239
- radiotherapy
 - CLL 201
 - multiple myeloma 237
- rasburicase 137
- RD regimen 235
- reactive systemic AA amyloidosis 238
- recombinases 107
- red cells 12, 17–19
 - abnormalities 23
 - amplification and maturation 14
 - anaemia 21, 22
 - antigens 334, 335
 - destruction 61, 61
 - DNA content 14
 - erythrocyte sedimentation rate 330, 332
 - folate 20, 57
 - inclusions 24
 - malignant disease 324
 - membrane 18–19, 19
 - metabolism 18, 18
 - metabolism defects 66–8, 66, 67
 - packed 343, 343
 - splenic control of integrity 117
 - volume 168
- red cell aplasia 247–9, 247, 248
- red cell count 20
- red cell fragmentation syndromes 70, 70, 70
- red cell indices
 - anaemia 21, 34
 - iron deficiency 34
 - normal 20
- red pulp of spleen 117, 117
- Reed-Sternberg cells 133, 206, 207
- refractory anaemia 178
 - with excess blasts 178, 180
 - with ring sideroblasts (RARS) 178
- refractory cytopenia
 - with multilineage dysplasia (RCMD) 178
 - with unilineage dysplasia (RCUD) 178
- refractory neutropenia 178
- refractory thrombocytopenia 178
- renal failure 325–6, 325, 325
- reticulocytes 24
- reticulocyte count 20
 - anaemia 22, 22
- reticuloendothelial system 61, 92
- retinal haemorrhage 21
- trans*-retinoic acid 142
- Rh blood group system 335, 336
 - genotypes 338
 - haemolytic disease of newborn 350–2, 351, 352
 - molecular genetics 337
- rheumatoid arthritis, haematological
 - changes 323–5, 324
- Rickettsia rickettsii* 339
- ring sideroblasts 38, 38, 178, 180, 183
- rituximab 143
 - CLL 200
 - idiopathic thrombocytopenic purpura 283
 - non-Hodgkin lymphoma 220
- rivaroxaban 317, 318
- romiplostim 283
- Rosai-Dorfmann disease 99
- ROTEM (thromboelastometry) 277, 301
- rouleaux formation 234
- ruxolitinib 141, 142
 - polycythaemia vera 171
- schistosomiasis 329
- Schwachman-Diamond syndrome 95, 244, 249
- SCL 4
- scurvy (vitamin C deficiency) 280, 280
- self-renewal 2, 3, 4
- senile purpura 280
- serum free light chain assay 232, 233
- Sézary syndrome 226
- sicca syndrome 261
- sickle cells 23
- sickle cell anaemia 80, 81–5
 - antenatal diagnosis 86
 - aplastic crises 83
 - clinical features 81–4, 82–4
 - haemolytic crises 83
 - laboratory findings 84
 - molecular pathology 81
 - organ damage 83–4, 83, 84
 - treatment 84–5
 - vaso-occlusive crises 82, 83
- sickle cell trait 80, 85
- sideroblastic anaemia 38–9, 38, 39
 - laboratory findings 38
- signal transducer and activator of transcription *see* STAT
- signal transduction 6–8, 7
- signal transduction inhibitors 142, 143

- sinus histiocytosis with massive lymphadenopathy 99
- slg **189**
- small lymphocyte lymphoma 220
 - immunophenotype **217**
- smouldering myeloma 229, 231, **231**
- solitary plasmacytoma 237
- somatic hypermutation 200
- somatic mutation 107
- South-East Asian ovalocytosis 65
- spectrin 19
- spherocytosis 19, 328
 - hereditary 64–5, **64**, 65
- spleen 116–21
 - anatomy and circulation 117, *117*
 - atrophy *117*
 - cords of 117, *117*
 - enlarged *see* splenomegaly
 - extramedullary haemopoiesis 118
 - functions 117
 - hypersplenism 119–20
 - hyposplenism 120, **120**, 121, **121**
 - imaging 118, *118*, *119*
 - red pulp 117, *117*
 - white pulp 117, *117*
- splenectomy 79, 120, **120**
 - idiopathic thrombocytopenic purpura 283
 - myelofibrosis 175
- splenic pooling 287
- splenomegaly 118–19, **120**
 - Felty's syndrome 324
 - marginal zone lymphoma 222
 - myelofibrosis *174*
 - platelet distribution 286
 - polycythaemia vera *169*
- SRSF2* mutation 184
- starry sky appearance 225
- STAT 6
- stem cells
 - homing 4
 - mesenchymal 4
 - mobilization 4
 - pluripotent 2–4, 3
- stem cell transplantation 250–63
 - acute myeloid leukaemia 153
 - allogeneic *see* allogeneic stem cell transplantation
 - aplastic anaemia 246–7
 - autologous 254, 255
 - bone marrow cells 251
 - chronic myeloid leukaemia 162, *163*
 - CLL 202
 - conditioning 253
 - haploidentical 256
 - idiopathic thrombocytopenic purpura 283
 - indications **251**
 - myelodysplasia 184
 - peripheral blood stem cells 251
 - post-transplant engraftment/immunity 253–4, *254*
 - potential donors **251**
 - principles 251, **251**
 - procedure 252
 - stem cell processing 251, 253, 253
 - umbilical cord blood cells 251
- stercobilin 61
- stercobilinogen 61
- steroid purpura 280
- stomatocytes 23
- storage pool diseases 287
- streptokinase 318
- Stuart-Prower factor *see* factor X
- subacute combined degeneration of cord 53, 54, 55
- superficial venous thrombosis 307
- systemic amyloid light chain amyloid disease 238–9, **238**, 239
- systemic diseases, haematological changes 321–32
 - anaemia of chronic disorders 322, **322**
 - congestive heart failure 326
 - hypothyroidism 327
 - infections 327–30, **327**, 328–30
 - liver disease 326–7, **326**
 - neoplasms 322–3, 323
 - non-specific monitoring 330–2, **331**
 - older patients 322
 - osteopetrosis 330
 - renal failure 325–6, 325, **325**
 - rheumatoid arthritis 323–5, *324*
 - systemic lupus erythematosus (SLE) 325
- T cells 103–4
 - functional aspects **105**
 - HLA-restricted 109
 - immunodeficiency **113**
- T-cell chronic lymphoid leukaemias 203–4
- T-cell lymphomas 225–6
- T-cell prolymphocytic leukaemia 203
- T-cell receptor, gene rearrangements 107, 108
- T helper cells 110
- target cells 23
- TdT **189**
- tear drop poikilocytes 23
- telomerase 129
- telomere 128, 129, *129*
- TET2* mutation 127, 184
- tetrahydrofolate 306
- thalassaemias **75**
 - α -thalassaemia syndromes **75**, 76, 76, 77
 - β -thalassaemia major 76–9, 77, 78
 - β -thalassaemia minor 79, 81
 - classification **75**
 - $\delta\beta$ -thalassaemia 81
 - geographical distribution 75
 - haemoglobin in 73, 74
 - thalassaemia intermedia 43–4, **75**, 79–81, 79, 80, 81
 - thalassaemia major 43, 44, **75**
- thalassaemia trait **75**, 79, 81
 - laboratory findings **38**
- thalidomide **143**
 - MTP regimen 235
 - multiple myeloma 236, 237
- thrombasthenia 266, 287
- thrombin 270–1, 272
- thrombin time 276, **276**
 - coagulation disorders **299**
- thrombocythaemia 172–3
 - clinical features **172**, 173
 - course 173
 - diagnosis 172–3
 - genetic mutations **166**, 167
 - laboratory findings **172**, 173, *173*, **173**
 - prognosis and treatment 173
 - survival **172**
- thrombocytopenia 281–7
 - with absent radii 244
 - amegakaryocytic 244
 - antiphospholipid syndrome 307
 - causes **281**
 - drug-induced *281*, **281**, 284, *284*
 - fetomaternal alloimmune 350
 - haemolytic uraemic syndrome 285, 285, 286
 - heparin-induced 314–15, *314*
 - of hypertensive disorders 348
 - infection-related 283
 - liver disease 297
 - malaria 329, 329
 - massive transfusion syndrome 287, 301
 - platelet destruction 282–7
 - platelet function disorders 287–9
 - platelet production failure 281, *281*, **281**
 - post-transfusion purpura 284, 342
 - pregnancy 347–8, *348*
 - refractive **178**
 - splenic pooling 287
 - thrombomimetics 289
 - viral infections 328
- thrombocytopenic purpura, autoimmune 282–3, 282

- thromboelastography 277, 300, 301
 thromboelastometry 277, 301
 thrombolytic therapy 287, 318–20, 319, **319**
 thrombomimetics 289
 thrombophilia 307–8
 thrombopoietin (TPO) 5, **6**, 265, 268
 thrombopoietin-receptor agonists 283
 thrombosis 302–10, 311–20
 anticoagulants 312–18
 antiplatelet drugs 318–20, 319, **319**
 arterial 303, 303, **303**
 fibrinolytic agents 318
 older patients 322
 post-thrombotic syndrome 318
 pregnancy 348
 prophylaxis 318
 venous *see* venous thrombosis
 thrombotic thrombocytopenic purpura 285, 285, 286
 thromboxane A2 268, 270
 ticagrelor 320
 tirofiban **319**, 320
 tissue factor *see* factor III
 tissue factor pathway inhibitor (TFPI) 272
 tissue plasminogen activator (TPA) 275, 276, 318
 tissue plasminogen activator inhibitor (TPAI) 275
 total iron-binding capacity **20**, 34–5, 36
 toxoplasmosis 329
 trabecular arteries 117, 117
 tranexamic acid 280
 transcobalamins 49
 transcription factors 8
 transferrin 28, **28**
 regulation 29
 transferrin receptor 1 regulation 29, 30
 transforming growth factor- β 5
 transfusion reactions 340–2, **341**
 transfusion-related acute lung injury (TRALI) 342
 transfusional iron overload 44, 44, **44**, 45
 translocations 129–30, 129, 130
 treosulfan 253
Treponema pallidum **339**
 tropical splenomegaly syndrome 119
 trypanosomiasis 329, 330
 tuberculosis 328
 tumour lysis syndrome 137
 tumour necrosis factor (TNF) 5, **6**, 90, 256
 tumour-suppressor genes 125
 tyrosine kinase inhibitors 141, **142**
 chronic myeloid leukaemia 160–1, 160, **160**, 161
 response to therapy 160–2, 160–2, **162**
 see also individual drugs
 tyrosine kinases 125
 ultra-large von Willebrand factor multimers 285, 285
 umbilical cord cell transplantation 251
 uraemia 287
 vaccination of hyposplenic patients 121, **121**
 vascular bleeding disorders 279–80, **279**
 acquired 279–80
 connective tissue disorders 279
 giant cavernous haemangioma 279
 hereditary haemorrhagic telangiectasia 279, 280
 vascular endothelial growth factor (VEGF) **6**
 vasoconstriction 273–4
 VCD regimen 235
 venesection in polycythaemia vera 170
 venous stasis 306
 venous thrombosis 303–7
 acquired risk factors 306–7, **307**
 deep vein thrombosis 308, **308**
 hereditary risk factors 304–6, **304**
 pathogenesis 303–4
 postoperative 306
 pulmonary embolus 308–9, 309
 superficial 307
 ventilation perfusion scintigraphy 309
 vinblastine **142**
 vinca alkaloids 141, **142**, 143
 vincristine **142**
 BEACOPP regimen 211
 CHOP regimen 223
 R-CVP regimen 222
 viral infections
 cancer patients 139
 haematological changes 328
 and haematological malignancies 124–5
 post-stem cell transplant 259, 259, 260
 post-transfusion hepatitis 338, **339**
 transfusion-related 342
 see also HIV/AIDS
 vitamin B12 49–51, 49
 absorption 49, 50, **50**
 biochemical function 51, 51
 neuropathy 53, 54, 55
 prophylactic 58
 serum **20**, **57**
 transport 49
 vitamin B12 deficiency 52–3
 causes **52**, 53
 clinical features 53–6, 54–6, **54**
 diagnosis 56–7, **57**
 pregnancy 347
 tests for cause 57, **57**
 treatment 57–8, **57**, 58
 vitamin C deficiency (scurvy) 280, 280
 vitamin K antagonists 315–18, **315–17**
 vitamin K deficiency 296–7, 297
 von Willebrand disease 295–6
 classification **295**
 laboratory findings **294**, 295
 treatment 295–6
 von Willebrand factor 266, 267, 269, 295
 ultra-large 285, 285
 VRD regimen 235
 VTD regimen 235
 Waldenström's macroglobulinaemia 221, 221
 warfarin **317**
 drug interactions 315, **316**
 haemostasis tests **299**, **315**
 overdose 316, **316**
 vitamin K deficiency 296
 warm autoimmune haemolytic anaemias 68–9, 69
 Weibel-Palade bodies 267, 285
 white cells 87–101, 102–15
 abnormal 93
 anaemia 22
 antibodies 342
 donor leucocyte infusions 162, 253, 261, 262
 malignant disease 94, 322, **324**
 see also specific types
 white cell count **20**, **88**
 myelofibrosis **174**
 thrombocythaemia **172**
 white pulp of spleen 117, 117
 WHO classification **208**, 354–6
 Wilson's disease 71, 326
 Wiskott-Aldrich syndrome 95, 281
 women
 iron requirements **33**
 pregnancy 347–8, 347, 348
Wuchereria bancrofti 331
 X-ray *see* radiography
Yersinia enterocolitica **339**
 Zieve's syndrome 325, 326

WILEY END USER LICENSE AGREEMENT

Go to www.wiley.com/go/eula to access Wiley's ebook
EULA.